Reaction–diffusion waves in cardiovascular diseases

A.V. Panfilov^{a,f}, H. Dierckx^{a,*}, V. Volpert^{b,c,d,e}

^a Department of Physics and Astronomy, Krijgslaan 281, Ghent University, Belgium

^b Institut Camille Jordan, UMR 5208 CNRS, University Lyon 1, 69622 Villeurbanne, France

^c INRIA Team Dracula, INRIA Lyon La Doua, 69603 Villeurbanne, France

^d Peoples' Friendship University of Russia (RUDN University), 6 Miklukho-Maklaya St, Moscow, 117198, Russian Federation

^e Poncelet Center, UMI 2615 CNRS, 11 Bolshoy Vlasievskiy, 119002 Moscow, Russian Federation

^f Laboratory of Computational Biology and Medicine, Ural Federal University, Ekaterinburg, Russian Federation

HIGHLIGHTS

- Cardiovascular diseases are reviewed from joint grounds of reaction–diffusion waves.
- Atherosclerosis is modeled as chronic inflammation of blood vessel walls.
- The state-of-the-art models of thrombus formation are considered.
- Cardiac arrhythmia modeling via reaction–diffusion equations is discussed.
- Response function perturbation theory for cardiac arrhythmias is presented.

ARTICLE INFO

Article history:

Received 10 December 2018

Received in revised form 30 March 2019

Accepted 5 April 2019

Available online xxxx

Communicated by V.M. Perez-Garcia

Keywords:

Cardiovascular diseases

Reaction–diffusion waves

Atherosclerosis

Thrombosis

Arrhythmia

ABSTRACT

Cardiovascular diseases remain one of the largest causes of death worldwide. The lethal pathologies often occur due to atherosclerosis, clot formation and cardiac arrhythmia. It turns out that all these pathologies can be described using the generic framework of multi-component reaction–diffusion equations. These parabolic partial differential equations sustain waves and pattern formation that are relevant in the study of all those processes. Here we present the first review which combines the description of these fields: atherosclerosis, clot formation and cardiac arrhythmias, and includes classic and recent developments in these areas. We show how mathematical models for the underlying physiological processes have been constructed, and which generic properties follow from their analytical or numerical study. Finally, we discuss the possibility of integrative studies of cardiovascular disease which will include both clot formation and cardiac arrhythmias. Such approach will be highly relevant to atrial fibrillation, a common cardiac arrhythmia whose main complication is clot formation and stroke.

© 2019 Elsevier B.V. All rights reserved.

Contents

1. Introduction	3
1.1. Cardiovascular diseases	3
1.1.1. Atherosclerosis	3
1.1.2. Cardiac arrhythmias	3
1.2. Structure of this review	3
2. Mathematical preliminaries: wave patterns in reaction–diffusion systems	3
2.1. Waves and pulses for the scalar reaction–diffusion equation	3
2.1.1. Existence and stability of waves	4
2.1.2. Existence and instability of pulses	4
2.2. Waves and pulses for monotone systems of equations	4
2.2.1. Existence and stability of waves	5
2.2.2. Existence and instability of pulses	5
2.3. Multi-dimensional problems and other generalizations	5
2.3.1. Response of a traveling wave to small stimuli	5

* Corresponding author.

E-mail address: hans.dierckx@ugent.be (H. Dierckx).

2.3.2.	Application: velocity–curvature relation for wave fronts	6
2.4.	Beyond one spatial dimension: spiral and scroll waves	7
3.	Atherosclerosis as chronic inflammation	7
3.1.	Physiology of atherosclerosis	7
3.2.	Atherosclerosis as a reaction–diffusion wave	7
3.2.1.	ODE model of chronic inflammation	7
3.2.2.	One-dimensional waves	8
3.2.3.	Two-dimensional waves	8
3.2.4.	Cellular automata	9
3.2.5.	The influence of hemodynamics	9
4.	Blood coagulation and thrombosis	9
4.1.	Physiological mechanism of clot formation	9
4.1.1.	The coagulation cascade	9
4.1.2.	Platelet aggregation	9
4.1.3.	Initiation and termination mechanisms	10
4.1.4.	Bleeding disorders and thrombosis	10
4.2.	Modeling of blood coagulation in quiescent plasma	10
4.2.1.	Reaction–diffusion model for the coagulation cascade	11
4.2.2.	Existence and stability of waves	11
4.2.3.	Reduction to the scalar equation and the speed of wave propagation	12
4.2.4.	2D models and clot growth arrest	12
4.3.	The influence of blood flow	12
4.4.	Regimes of blood coagulation	14
4.4.1.	Initiation without propagation	14
4.4.2.	Initiation with propagation and with growth arrest	14
4.4.3.	Initiation with propagation and without growth arrest	14
5.	Basic principles of cardiac arrhythmia modeling	15
5.1.	Normal function of the heart	15
5.1.1.	Physiology of the heart	15
5.1.2.	Excitation sequence in sinus rhythm	15
5.2.	Brief overview of cardiac arrhythmias	15
5.3.	Rotor formation	15
5.4.	Electrical activity of the heart as a reaction–diffusion system	16
5.4.1.	Cell model	16
5.4.2.	Anisotropic coupling of cells	16
5.4.3.	Monodomain model	17
5.5.	Approaches beyond the mono- and bidomain models	17
6.	Dynamics of wave patterns underlying cardiac arrhythmias	18
6.1.	Numerical approach to cardiac arrhythmia management	18
6.1.1.	Numerical studies of fundamental dynamics	18
6.1.2.	Numerical modeling with direct clinical or experimental applications	18
6.2.	Spiral breakup during fibrillation explained as a 1D map	19
6.3.	Dynamics of complex structures unraveled by properties of smaller building blocks	22
6.3.1.	Motivation	22
6.3.2.	Response function theory of spiral waves	22
6.3.3.	Spiral wave response functions	23
6.3.4.	Response of a spiral wave to small perturbations	24
6.3.5.	Dynamics of filaments of scroll waves	25
6.4.	Curved-space viewpoint on anisotropic wave propagation in the heart	25
6.4.1.	Motivation	25
6.4.2.	Operational definition of distance	25
6.4.3.	Role of extrinsic and intrinsic curvature in the dynamics of excitation patterns	26
6.4.4.	Some results on the dynamics of activation patterns in the anisotropic medium	26
7.	Discussion	27
7.1.	Transition processes in physiology	27
7.1.1.	Multiple equilibria in physiological systems	27
7.1.2.	Similar models for different processes	28
7.2.	Integration of models for cardiovascular diseases	28
	Acknowledgments	29
	Appendix Elements from differential geometry	29
A.1.	Metric tensor	29
A.1.1.	Covariant differentiation	29
A.2.	Geodesics	29
A.3.	Riemann normal coordinates	29
A.4.	Intrinsic curvature	29
A.5.	Extrinsic curvature	30
	References	30

1. Introduction

1.1. Cardiovascular diseases

According to the World Health Organization the largest cause of death in the world is ischemic heart disease, followed by stroke [1]. In many cases, both these diseases are a result of reduced blood supply which can occur due to narrowing of the arteries as result of atherosclerosis or formation of clots due to internal blood coagulation. Death from ischemic heart disease occurs in 90% of cases due to cardiac arrhythmias (irregular heart beat). Also, one third of all strokes was found to occur in patients with cardiac arrhythmias [2]. Thus we see that the processes of atherosclerosis, clot formation and cardiac arrhythmias are extremely important. These three processes are very different in nature and are studied by different groups of researchers, published in different scientific journals, etc. Strikingly, atherosclerosis, clot formation and cardiac arrhythmias can be described using the same mathematical background, the so-called reaction–diffusion systems. In this review article we provide a first joint description of these processes in a unified framework and discuss possible further interaction between these fields.

1.1.1. Atherosclerosis

Involving blood vessels and heart, cardiovascular diseases (CVD) represent the main cause of mortality in the world affecting mostly elder people. One of the most common CVD is coronary artery disease (angina, myocardial infarction) caused by the limitation of blood flow to the heart because of atherosclerosis developed in the coronary artery. The lack of oxygen by heart muscle cells can lead to their death resulting in the cardiac tissue damage. Chronic narrowing of coronary artery can cause ventricular arrhythmia and fibrillation, which often lead to death. Along with coronary artery disease, stroke, peripheral artery disease, and some others also involve atherosclerosis.

Atherosclerosis is a chronic inflammation of blood vessel walls whose initiation is related to the high level of low-density lipoproteins (LDL, cholesterol) in blood. Penetrating in the blood vessel wall, LDL is oxidized (ox-LDL) becoming a pro-inflammatory agent and provoking innate immune response. White blood cells (monocytes) come to the inflammation site, penetrate the vessel wall, transform into foam cells and remain trapped there increasing inflammation even more. This self-amplifying inflammation becomes chronic, leading to thickening of blood vessel walls and narrowing the lumen with blood flow (stenosis). As indicated above, stenosis perturbs blood circulation and oxygen supply. Furthermore, atherosclerotic plaque can rupture due to the pressure from the blood flow in such a way that some pieces of it detach and possibly block blood circulation downstream leading to stroke or heart attack. Another possible consequence of plaque rupture is blood coagulation, which can also lead to vessel occlusion.

Blood coagulation is an important physiological mechanism aimed to preserve hemostasis in the case of injury. It includes biochemical reactions in blood plasma and platelet aggregation being also regulated by blood flow and blood vessel walls. Reactions of blood coagulation involve numerous biochemical substances called blood factors. Insufficient blood coagulation can lead to bleeding disorders while excessive blood coagulation to thrombosis. Thrombosis is a disease which can have many different manifestations, such as deep vein thrombosis, portal vein thrombosis, and so on. A piece of thrombus can detach and circulate in the blood, provoking a venous thromboembolism. Arterial embolism can lead to stroke or heart attack, in particular because of atherosclerosis narrowing the lumen and facilitating vessel occlusion. Arterial thrombosis often occurs due to the atheroma (atherosclerotic plaque) rupture. Arterial occlusion can also occur because of blood clots formed in the heart.

1.1.2. Cardiac arrhythmias

It is well known that there exist electrical waves in the heart, as one of the first medical exams in cardiology is the electrocardiogram. Such electrical waves trigger mechanical contraction, and if they fail to propagate or do it in an abnormal manner, contraction of the heart can be significantly impaired. Such abnormal regimes of wave propagation in the heart are called cardiac arrhythmias. The study of arrhythmias is a very difficult problem due to several reasons. First it is a very rare event. Sudden cardiac death can be initiated just by a single abnormal beat. Thus if we consider that the heart contracts about 70 times per minute, or approx. 40 million times per year, the tracing and predicting of such abnormality is a very difficult task. Also, this abnormal situation can occur on the time scales of the order of seconds, which is very fast compared to other biological processes leading to other diseases. For example, abnormal tissue growth may take years. Finally, the waves are generated by billions of cardiac cells, and the excitation of each of the cells is a result of dynamics of tens of thousands individual ionic channels. Cardiac arrhythmias are thus a typical multiscale problem. As a result, the precise mechanisms behind many arrhythmias are currently still poorly understood, and successful treatment is nowadays often experience-based. Therefore, helping patients will require an interdisciplinary effort that bridges across spatial scales, from the genetic and intracellular scales via collective dynamics and (computer) modeling to the clinical practice. In mathematical terms, cardiac tissue is often represented as a set of reaction–diffusion equations and although the work is far from finished, the analysis of these models (either numerical or semi-analytical) has contributed to a better understanding of the dynamics of heart rhythm disorders.

1.2. Structure of this review

This paper is structured as follows: first we present a general introduction to reaction–diffusion systems and basic properties of traveling waves in those systems, in Section 2. This knowledge is used in Sections 3–4 to discuss the atherosclerosis and blood coagulation processes. The second part of this review entails cardiac arrhythmias. The mathematical modeling and phenomenology of emergent patterns are presented in Section 5. Thereafter, we discuss how mathematical modeling using reaction–diffusion systems has already provided valuable insights on the dynamics of cardiac arrhythmia in Section 6. First, we discuss the relevance of numerical modeling for clinical applications (Section 6.1) and we show that, in spite of the complexity of the non-linear functions in the PDE models, generic features of electrical pattern evolution can be inferred based on dynamical systems theory, geometrical grounds and asymptotic analysis (Sections 6.2–6.4). We end with a general discussion, conclusions and outlook in Section 7.

2. Mathematical preliminaries: wave patterns in reaction–diffusion systems

2.1. Waves and pulses for the scalar reaction–diffusion equation

In this section we briefly recall the main results on waves and pulses for the scalar reaction–diffusion equations (i.e. equations in a single state variable). In the next section, they will be generalized for some classes of reaction–diffusion systems. Consider the equation

$$\frac{\partial u}{\partial t} = D \frac{\partial^2 u}{\partial x^2} + F(u), \quad (2.1)$$

where $u(x, t)$ has $x \in \mathbb{R}$. The function $F(u)$ is such that $F(0) = F(1) = 0$. The typical examples of the non-linearity are shown

Fig. 1. Typical examples of the nonlinearity in Eq. (2.1) in the monostable case (left) and the bistable case (right).

in Fig. 1. $F(u)$ can be positive in the interval $(0, 1)$ (monostable case) or it can change sign being negative for small values of u and positive for large values (bistable case). More complex functions are also encountered in some applications.

A traveling wave solution of Eq. (2.1) is a solution of the form $u(x, t) = w(x - ct)$, where the constant c is the wave speed. The function $w(\xi)$ satisfies the equation

$$Dw'' + cw' + F(w) = 0 \quad (2.2)$$

with the limits at infinity:

$$w(\infty) = 0, \quad w(-\infty) = 1. \quad (2.3)$$

The wave speed c is a priori unknown, and we should determine its values for which problem (2.2), (2.3) has a solution.

2.1.1. Existence and stability of waves

In the case of the scalar equation, wave existence is well studied (see [3] and the references therein). For the functions shown in Fig. 1, the main results can be summarized as follows. In the monostable case, (monotone) waves exist for all values of the speed greater than or equal to the minimal speed c_0 . In the bistable case, the wave exists for a unique value of speed. For general $F(u)$, these values of speed cannot be found exactly, but they can be estimated or approximated by various analytical formulas.

In the bistable case, the wave is globally asymptotically stable [4–7]. The solution of Eq. (2.1) with the initial condition $u(x, 0) = u_0(x)$, such that it is a monotonically decreasing function with the limits $u(-\infty) = 1$, $u(\infty) = 0$, was shown to converge to a traveling wave, that is,

$$\sup_x |u(x, t) - w(x - ct - h)| \rightarrow 0 \quad \text{as } t \rightarrow \infty. \quad (2.4)$$

The conditions on the function $u_0(x)$ can be less restrictive, but this is sufficient for what follows. Let us recall that solutions of problem (2.2), (2.3) are invariant with respect to translation in space. This means that the function $w(\xi + h)$ is a solution of this problem for any real h . The value of the constant h in (2.4) depends on the initial condition. This is so-called stability with shift.

In the monostable case, the waves are monotonically decreasing functions for $c \geq c_0$. They also exist for $0 < c < c_0$, but they become non-monotone. The monotone waves are asymptotically stable in some properly chosen weighted spaces, non-monotone waves are unstable.

2.1.2. Existence and instability of pulses

The pulse is a positive stationary solution $w_p(x)$ of Eq. (2.1) with the limits $w_p(\pm\infty) = 0$ at infinity. It can be easily verified that such solutions exist in the bistable case if and only if

$$\int_0^1 F(u) du > 0. \quad (2.5)$$

Such solutions do not exist in the monostable case.

Fig. 2. Schematic representation of the pulse solution $w_p(x)$ and of the solution $u(x, t)$ of Eq. (2.1) forming two waves propagating in the opposite directions. This structure forms due to the instability of the pulse solution.

Multiplying equation (2.2) by w' and integrating from $-\infty$ to ∞ , we conclude that inequality (2.5) is a necessary and sufficient condition for the wave speed c to be positive. Hence, the pulse solution exists if and only if the wave solution has a positive speed. This assertion is quite simple for the scalar equation. We will see below that it holds true for the monotone system of equations where condition (2.5) is not applicable, neither to determine the wave speed nor to assert the existence of a pulse solution.

The pulse solution is unstable as a solution of Eq. (2.1). If we consider the initial condition $u(x, 0) = w_p(x) + \epsilon(x)$, where $\epsilon(x)$ is a small perturbation, then the solution $u(x, t)$ of this equation does not converge, in general, to the stationary solution $w_p(x)$. The instability occurs since the linearized operator

$$Lv = Dv'' + F'(w_p(x))v$$

has a positive eigenvalue [3,8,9]. If $\epsilon(x) > 0$, then the solution $u(x, t)$ locally converges to 1 (in every bounded interval), and it forms two waves propagating in the opposite directions (Fig. 2). If the perturbation is negative, the solution decays and converges to 0. Thus, the pulse solution separates two classes of initial conditions with different behavior of solutions. We will see below that this conclusion remains valid for monotone systems of equations, and it has important physiological implications.

2.2. Waves and pulses for monotone systems of equations

Let us now consider the reaction–diffusion equation (2.1) in the vector case, where $\mathbf{u} = (u_1, \dots, u_n)$, $\mathbf{F} = (F_1, \dots, F_n)$, D is a diagonal matrix with positive diagonal elements d_i . Furthermore, we assume that the following condition is satisfied:

$$\frac{\partial F_i}{\partial u_j} \geq 0, \quad i \neq j, \quad i, j = 1, \dots, n. \quad (2.6)$$

If (2.6) holds, the system (2.1) is called a monotone system. Monotone systems are characterized by the fulfillment of the

maximum principle, and for them the main results on the existence and stability of waves remain valid.

2.2.1. Existence and stability of waves

Suppose that $\mathbf{F}(\mathbf{w}_+) = \mathbf{F}(\mathbf{w}_-) = \mathbf{0}$ for some $\mathbf{w}_+$ and $\mathbf{w}_-$, where $\mathbf{w}_+ < \mathbf{w}_-$. Here and below, the inequalities between vectors are understood component-wise. In the bistable case, where all eigenvalues of the matrices $\mathbf{F}'(\mathbf{w}_\pm)$ lie in the left-half plane, and for any other point $\mathbf{w}_0$ such that $\mathbf{F}(\mathbf{w}_0) = \mathbf{0}$ and $\mathbf{w}_+ \leq \mathbf{w}_0 \leq \mathbf{w}_-$, the principal eigenvalue¹ is positive, then there exists a unique traveling wave solution $\mathbf{u}(x, t) = \mathbf{w}(x - ct)$ of system (2.1) with the limits

$$\mathbf{w}(\pm\infty) = \mathbf{w}_\pm \quad (2.7)$$

at infinity. This traveling wave solution is a monotonically decreasing function of x , and it is globally asymptotically stable. The wave speed admits the following minimax representation:

$$c = \inf_{\rho \in K} \sup_{x,i} \frac{d_i \rho_i'' + F_i(\rho)}{-\rho_i'} = \sup_{\rho \in K} \inf_{x,i} \frac{d_i \rho_i'' + F_i(\rho)}{-\rho_i'}, \quad (2.8)$$

where K is the set of all monotonically decreasing vector functions $\rho(x)$ continuous with its first derivative and bounded second derivative, and having limits $\rho(\pm\infty) = w_\pm$ at infinity.

In the monostable case, the principal eigenvalue of the matrix $\mathbf{F}'(\mathbf{w}_+)$ is positive, and of the matrix $\mathbf{F}'(\mathbf{w}_-)$ negative. If there are no other zeros of the function $F(w)$ in the rectangle $w_+ \leq w \leq w_-$, then traveling waves with the limits (2.7) exist for all values of the speed greater than or equal to the minimal speed c_0 . The latter admits the first representation in (2.8). The waves are asymptotically stable in properly chosen weighted spaces [8,9].

2.2.2. Existence and instability of pulses

The existence of pulses has been demonstrated for a monotone system of two equations [10,11] and for the system of equations of blood coagulation [12] (Section 4) but not yet in the general setting. The main result can be formulated in terms of traveling waves: the pulse exists if and only if the wave speed is positive. Similar to the existence of waves, the proof of this result uses quite sophisticated mathematical methods, including the Leray-Schauder method for elliptic operators in unbounded domains and a priori estimates of solutions in some properly chosen weighted spaces.

Similar to the scalar equation, pulses for monotone systems are unstable. They separate the initial conditions for which the solution of Eq. (2.1) is growing and locally converging to $\mathbf{w}_-$ and decaying to $\mathbf{w}_+$. The convergence to the traveling wave solutions is observed numerically in this case but it is not proved.

2.3. Multi-dimensional problems and other generalizations

The results on wave existence, stability and the speed of propagation are generalized for the multi-dimensional scalar equation [13] and for monotone systems of equations [14]. Traveling waves in a two-dimensional strip with nonlinear boundary conditions arising in modeling of atherosclerosis are studied in [15–17]. We will return to this problem in Section 3. Furthermore, the wave existence is proved in the one-dimensional case for locally monotone systems [3,18]. This is a class of systems for which inequality (2.6) holds only at the sets where $F_i(\mathbf{u}) = 0$.

For a generic multi-component reaction–diffusion equation that supports traveling waves, it is nevertheless possible to derive how wave speed depends on external perturbations and the wave front's curvature. This question is discussed in the next section.

¹ Principal eigenvalue is the eigenvalue with the maximal real part. In the case of the matrices with positive off-diagonal elements, it is real, simple, and the corresponding eigenvector is positive. The same property remains valid for irreducible matrices with non-negative off-diagonal elements.

2.3.1. Response of a traveling wave to small stimuli

Let us consider wave propagation on a line with constant diffusion coefficient. Then, we may write a multi-component reaction–diffusion system as:

$$\partial_t \mathbf{u} = \hat{\mathbf{P}} \partial_x^2 \mathbf{u} + \mathbf{F}(\mathbf{u}), \quad (2.9)$$

where $\hat{\mathbf{P}}$ is a matrix containing the diffusion coefficient of each variable on its diagonal. Note that the partial differential equations that describe electrical activation of the heart can also be cast into this form; see Eq. (5.10) below.

We suppose that $\mathbf{u}_0(x, t) = \mathbf{U}_0(x - ct)$ is a stable 1D traveling wave solution to Eq. (2.9), whence

$$\hat{\mathbf{P}} \partial_\rho^2 \mathbf{u}_0 + c \partial_\rho \mathbf{u}_0 + \mathbf{F}(\mathbf{u}_0) = \mathbf{0}. \quad (2.10)$$

Here, $\rho = x - ct$ is a co-moving coordinate with the solution. Note that, since space dependency occurs via the terms $D_0 \partial_\rho^2 + c \partial_\rho$, one finds that

$$c \propto \sqrt{D}. \quad (2.11)$$

Next, suppose that at $t = 0$, the state is perturbed to $\mathbf{U}_0 + \tilde{\mathbf{U}}$, with small $\tilde{\mathbf{U}} = \mathcal{O}(\eta)$, then inserting this in Eq. (2.9) delivers in linear order in η that

$$\partial_t \tilde{\mathbf{U}} = \hat{\mathbf{L}} \tilde{\mathbf{U}}, \quad (2.12a)$$

$$\hat{\mathbf{L}} = \hat{\mathbf{P}} \partial_\rho^2 + c \partial_\rho + \mathbf{F}'(\mathbf{U}_0). \quad (2.12b)$$

Here $\mathbf{F}'(\mathbf{U}_0)$ is the Jacobian matrix of the reaction kinetics, i.e. shorthand for the matrix with elements $\partial F^m / \partial u^p$, evaluated at the reference solution $\mathbf{U}_0$.

Therefore, if $\tilde{\mathbf{U}}$ contains a contribution from an unstable eigenvector of $\hat{\mathbf{L}}$, i.e. $\hat{\mathbf{L}} \psi = \lambda \psi$ with $\Re(\lambda) > 0$, $\tilde{\mathbf{U}}$ will exponentially grow in time, rendering $\mathbf{U}_0$ unstable. In what follows, we will suppose that $\hat{\mathbf{L}}$ has no eigenvalues with strictly positive real part.

Note that since Eq. (2.9) is invariant with respect to translation in space and time, $\hat{\mathbf{L}}$ has an eigenfunction with zero eigenvalue: if $\mathbf{U}_0(\rho)$ is a solution, then so should the shifted solution $\mathbf{U}_0(\rho - \epsilon)$. Expanding in linear order in η delivers

$$\hat{\mathbf{L}} \partial_\rho \mathbf{U}_0 = \mathbf{0}, \quad \mathbf{V}_\rho = -\partial_\rho \mathbf{U}_0. \quad (2.13)$$

The minus sign in the definition of the zero mode $\mathbf{V}_\rho$ is a convention which will avoid a minus sign later on, in Eqs. (2.24).

Other eigenvalues of $\hat{\mathbf{L}}$ are not necessarily real-valued, since the operator $\hat{\mathbf{L}}$ is not self-adjoint with respect to the inner product (in n spatial dimensions):

$$\langle \mathbf{f} | \mathbf{g} \rangle = \int_{\mathbb{R}^n} \mathbf{f}^H \mathbf{g} \, d^n x. \quad (2.14)$$

In practice, we will work with functions $\mathbf{f}$ that decay exponentially at large distance and functions $\mathbf{g}$ that grow at most polynomially, such that the integrals (2.14) converge.

The adjoint operator to the operator $\hat{\mathbf{L}}$, denoted $\hat{\mathbf{L}}^\dagger$ is the operator defined by

$$\langle \mathbf{f} | \hat{\mathbf{L}} \mathbf{g} \rangle = \langle \hat{\mathbf{L}}^\dagger \mathbf{f} | \mathbf{g} \rangle \quad (2.15)$$

for all $\mathbf{f}, \mathbf{g}$ in a chosen function space. Due to our assumptions that $\mathbf{f}$ are tempered and $\mathbf{g}$ grow at most polynomially, boundary terms from integration by parts vanish and

$$\hat{\mathbf{L}}^\dagger = \hat{\mathbf{P}}^H \partial_\rho^2 - c \partial_\rho + \mathbf{F}^H(\mathbf{u}_0). \quad (2.16)$$

While the convection term can be eliminated by a change of variables, the reaction term makes $\hat{\mathbf{L}}$ generally non self-adjoint when more than one state variable is present.

Next, it is usually assumed that $\hat{\mathbf{L}}^\dagger$ also has a zero mode, $\mathbf{W}^\rho$, satisfying

$$\hat{\mathbf{L}}^\dagger \mathbf{W}^\rho = \mathbf{0}. \quad (2.17)$$

It is convenient to normalized the adjoint critical mode such that $\langle \mathbf{W}^\rho | \mathbf{V}_\rho \rangle = 1$.

The physical meaning of $\mathbf{W}^\rho$ can be assessed by considering the effect of a perturbation $\mathbf{h}(\vec{r}, t)$, say of order η , to the right-hand side of Eq. (2.9):

$$\partial_t \mathbf{u} = \hat{\mathbf{P}} \partial_x^2 \mathbf{u} + \mathbf{F}(\mathbf{u}) + \mathbf{h}. \quad (2.18)$$

Since the traveling wave solution $\mathbf{U}_0$ is stable, we can search for a solution that is the original traveling wave, but propagating at a different velocity: in a moving frame with the origin at the wave front position $X(t)$, i.e. $x = X(t) + \rho$ one expects that

$$\begin{aligned} \mathbf{u}(x, t) &= \mathbf{U}_0(\rho) + \tilde{\mathbf{U}}(\rho, t) + \mathcal{O}(\eta^2), \\ \partial_t X(t) &= c + \mathcal{O}(\eta). \end{aligned} \quad (2.19)$$

The decomposition (2.19) can be made unambiguous by demanding that

$$\langle \mathbf{W}^\rho | \tilde{\mathbf{U}} \rangle = 0 \quad (2.20)$$

since, if $\tilde{\mathbf{U}}$ possesses a small component along $\partial_\rho \mathbf{U}_0$, the wave front position can be changed to satisfy the gauge condition (2.20). Since Eq. (2.20) should hold for all times, it follows that

$$\langle \mathbf{W}^\rho | \partial_t \tilde{\mathbf{U}} \rangle = 0. \quad (2.21)$$

Now, plugging expansion (2.19) into Eq. (2.18) delivers for the left-hand side that

$$\partial_t \mathbf{U} = -\partial_\rho \mathbf{U}_0 \partial_t X + \partial_t \tilde{\mathbf{U}} = \partial_t X \mathbf{V}_\rho + \partial_t \tilde{\mathbf{U}}, \quad (2.22)$$

while for the right-hand side, it holds up to linear order in η that:

$$\hat{\mathbf{P}} \partial_\rho^2 \mathbf{U} + \mathbf{F}(\mathbf{U}) + \mathbf{h} = \hat{\mathbf{P}} \partial_\rho^2 \mathbf{U}_0 + \mathbf{F}(\mathbf{U}_0) + \mathbf{h} + \hat{\mathbf{L}}(\tilde{\mathbf{U}}) = c \mathbf{V}_\rho + \hat{\mathbf{L}}\tilde{\mathbf{U}} + \mathbf{h} + \hat{\mathbf{L}}(\tilde{\mathbf{U}}). \quad (2.23)$$

Equating both sides and taking the inner product with $\mathbf{W}^\rho$ delivers, with $\langle \mathbf{W}^\rho | \hat{\mathbf{L}}\tilde{\mathbf{U}} \rangle = \langle \hat{\mathbf{L}}^\dagger \mathbf{W}^\rho | \tilde{\mathbf{U}} \rangle = \langle \mathbf{0} | \tilde{\mathbf{U}} \rangle = 0$, it follows that

$$\partial_t X(t) = c + \langle \mathbf{W}^\rho | \mathbf{h} \rangle + \mathcal{O}(\eta^2). \quad (2.24)$$

Thus, knowledge of the critical adjoint eigenmode $\mathbf{W}^\rho$ allows us to find the modified propagation velocity under a small perturbation. One could also say that the components of $\mathbf{h}$ along the modes with negative real part of $\hat{\mathbf{L}}$ will decay in time, while the component of $\mathbf{h}$ along $\mathbf{W}^\rho$ causes a net spatial shift, leading to a modified propagation speed.

The result (2.24) is the equivalent of the impulse response for linear systems in engineering or Green's functions in physics. If one knows the response function of a system (either in full or that it is localized), it is possible to predict the dynamics of larger patterns that have the wave front as it building block; see e.g. the velocity–curvature relation (2.26) below.

Remarkably, it was found that response functions for traveling waves are localized in space, precisely at the wave front. Therefore, a traveling wave effectively behaves as a localized ‘front’. This makes it possible to regard a two- or three-dimensional traveling wave as a bundle of one-dimensional solutions, which we will use in the next paragraph to find the propagation velocity as a function of wave front curvature.

2.3.2. Application: velocity–curvature relation for wave fronts

Eqs. (2.24) can be used to find the motion of curved wave fronts. For now, we consider an isotropic medium; dealing with anisotropy is deferred to Section 6.4.

Fig. 3. Traveling wave solution (propagating to the right) together with the ‘density’ function $w(\rho)$ (shaded), showing that the linear curvature correction to the velocity is sensitive to the curvature at the wave front. Full line denotes u and dotted line v in Barkley’s model, with parameters $\hat{\mathbf{P}} = \text{diag}(1, 0)$, $a = 0.75$, $b = 0.02$, $\epsilon = 0.04$.

For wave fronts, we start from the observation that for a cylindrical outward wave front of radius r measured the diffusion term in polar coordinates can be written

$$\hat{\mathbf{P}} \Delta \mathbf{u} = \hat{\mathbf{P}} \partial_r^2 \mathbf{u} + \hat{\mathbf{P}} \frac{1}{r} \partial_r \mathbf{u}. \quad (2.25)$$

Thus, one finds Eq. (2.9) with $\mathbf{h} = \hat{\mathbf{P}} \frac{1}{r} \partial_r \mathbf{u}$, such that for small or moderate wave front curvature $K = 1/r$, the wave front velocity will become

$$\partial_t X = c - \gamma K, \quad \gamma = \langle \mathbf{W}^0 | \hat{\mathbf{P}} \partial_r \mathbf{U}_0 \rangle. \quad (2.26)$$

This derivation can be formalized to other geometries by introducing Gauss normal coordinates around the front [19]; in 3D, $K = 1/r_1 + 1/r_2$ with r_1, r_2 the principal curvatures at the wave front. Eq. (2.26) is known as the velocity–curvature relation for wave fronts in reaction–diffusion, sometimes called the eikonal equation. Notably, in most systems, $\gamma > 0$ such that an expanding wave will need more time to expand than a plane wave. In a clinical context, Eq. (2.26) is referred to as source–sink mismatch, since when $K > 0$, cardiac myocytes at the wave front on average need to excite a larger number of cells just ahead of the front. In physicist’s term, this is a pure curvature effect, as the definition of extrinsic curvature of a surface is

$$K = \lim_{A \rightarrow 0} \partial_n A / A \quad (2.27)$$

with n being the normal vector in the direction of propagation and A the surface area of a small portion of the wave front. Thus, a curvature effect is equivalent to a source–sink effect.

Note that the term γK in Eq. (2.26) originates from

$$\gamma K = \int_{\mathbb{R}} W^0 \hat{\mathbf{P}} \partial_\rho \mathbf{u}_0 d\rho = \int_{\mathbb{R}} w(\rho) K(\rho) d\rho. \quad (2.28)$$

Thus, the curvature of the wave front is weighted with a function $w = \sum_a W_a^0(\rho) D_a \partial_\rho (U_0)_a(\rho)$, with the sum running over all diffusing state variables. It is found numerically that the $\mathbf{W}^0$ are strongly localized around the wave front; Fig. 3 shows $w(\rho)$ together with $u(\rho)$ and $v(\rho)$ for Barkley’s model, i.e. $\mathbf{u} = [u \ v]^T$, $\hat{\mathbf{P}} = \text{diag}(1, 0)$ and $\mathbf{F}(\mathbf{u}) = [\epsilon^{-1} u(1 - u)(u - (v + b)/a), \ u - v]^T$. Higher order momenta of the distribution $w(\rho)$ give rise to some of the higher-order curvature corrections to the $c(K)$ relation [19].

2.4. Beyond one spatial dimension: spiral and scroll waves

We have only discussed results on traveling waves in one spatial dimension here. In two or three spatial dimensions, other distinct patterns called spiral waves may occur. Since they are mostly relevant to cardiac arrhythmias and not to atherosclerosis or thrombosis, we will discuss spiral waves extensively in Section 5.

3. Atherosclerosis as chronic inflammation

3.1. Physiology of atherosclerosis

The development of atherosclerosis is associated with various risk factors, such as inadequate nutrition, smoking, etc., although the immediate causes of initiation of its onset have so far not been precisely identified. One of the main risk factors is cholesterol (low-density lipoproteins, LDL), which penetrates into the blood vessels walls (intima), and becomes oxidized (ox-LDL). At the same time, high-density lipoproteins (HDL) prevent the oxidation of LDL taking on the oxidation process. Thus, already at the initial stage and, as we shall see later, the entire development of this disease is the counteraction of various pro- and anti-atherosclerotic factors. (See Fig. 4.)

When ox-LDL accumulates in the intima, it initiates the immune response of the organism. It activates surface receptors of the endothelial cells separating the bloodstream (lumen) and the intima. Monocytes, a type of white blood cells, roll along the inner surface of the blood vessels, where they are pushed by the red blood cells. When monocytes get into the place of the developing inflammation, they are hammered by the surface receptors, stop there and penetrate into the intima. There they are transformed into macrophages, the primary role of which is to absorb ox-LDL and other pro-inflammatory factors. Macrophage are eventually transformed into foam cells (lipid laden cells) which secrete various proinflammatory factors (IL-1, TNF- α) and lead to a self-amplifying process of chronic inflammation. Moreover, foam cells are trapped in the intima causing a gradual thickening of blood vessel walls and narrowing of the lumen. This is the most essential clinical consequence of atherosclerosis.

At the same time, the development of the inflammatory process is accompanied by the production of various anti-inflammatory factors that prevent the production of inflammatory cytokines. Besides, inflammation promotes proliferation and migration of smooth muscle cells which form a fibrous cap over the lipid deposit. It protects the inflammation site from further recruitment of monocytes. On the other hand, it modifies the blood flow geometry resulting in other important consequences. The most significant among them is a possible plaque rupture due to the pressure exerted by the bloodstream. In this case, blood coagulation can begin possibly resulting in vessel occlusion, or some solid pieces of plaque can depart with flow leading to stroke or heart attack if they block blood vessels downstream.

Thus, atherosclerosis is a chronic inflammation of blood vessel walls that leads to plaque formation and vessel remodeling. At the advanced stages, the lumen becomes narrow complicating blood circulation, and the plaque can rupture resulting in vessel occlusion and important clinical casualties.

3.2. Atherosclerosis as a reaction–diffusion wave

Theoretical modeling of atherosclerosis began with the investigation of the LDL distribution in the intima [21]. Taking into account LDL diffusion from the lumen and convection through the vessel wall, it was possible to describe the experimental data. The influence of blood pressure on the LDL distribution

was studied in [22]. It was shown that the most significant effect on the growth of LDL concentration in the intima is an increase in the permeability of the endothelial layer with increasing pressure. Oxidation of LDL in the presence of HDL was studied in [23]. A more complete model with several cell types and cytokines is suggested in [24]. The model takes into account reverse cholesterol transport (RCT) where HDL removes ox-LDL from the foam cells and transports it back to the liver. Furthermore, RCT reduces inflammation converting pro-inflammatory M1 macrophages into anti-inflammatory M2 macrophages. Development of atherosclerosis as a reaction–diffusion wave was considered in [20,25,26].

3.2.1. ODE model of chronic inflammation

The simplest model of atherosclerosis consists of two equations [25]:

$$\begin{cases} \frac{dM}{dt} = f_1(A) - \lambda_1 M \\ \frac{dA}{dt} = f_2(A)M - \lambda_2 A \end{cases}, \quad (3.1)$$

where M is the density of immune cells and A is the concentration of cytokines,

$$f_1(A) = \frac{\alpha_1 + \beta_1 A}{1 + A/\tau_1}, \quad f_2(A) = \frac{\alpha_2 A}{1 + A/\tau_2},$$

The function $f_1(A)$ models the recruitment of the immune cells from the blood flow, promoted by the inflammatory cytokines, where $\alpha_1 = f_1(0)$ corresponds to the beginning of the inflammation, that is the recruitment of monocytes due to the presence of ox-LDL, promoted by endothelial adhesion molecules, chemoattractants and growth factors. The factor $1 + A/\tau_1$ represents the effect of the fibrous cap created by smooth muscle cells. The term $f_2(A)M$ models the cytokines production rate, where $\alpha_2 A$ represents the secretion of pro-inflammatory cytokines promoted by the pro-inflammatory cytokines themselves and $1 + A/\tau_2$ represents the inhibition of the pro-inflammatory cytokines secretion mediated by the anti-inflammatory cytokines. The last terms in Eqs. (3.1) represent degradation of immune cells and cytokines.

Introducing the functions $F_1(A) = f_1(A)/\lambda_1$, $F_2(A) = \lambda_2 A/f_1(A)$, we can find the stationary points of system (3.1) from the equation $F_1(A) = F_2(A)$. One of the possible cases, where there are three stationary points, is shown in Fig. 5 (left). The points E_0 and E_r are stable while the intermediate point E_l is unstable. Hence, it is a bistable case, E_0 corresponds to the healthy state and E_r to the disease state.

If we decrease the parameter α_1 (concentration of ox-LDL), then the function $F_1(A)$ also decreases. For sufficiently small α_1 there is only one stationary point E_0 , and it is globally stable. On the other hand, for α_1 sufficiently large, there are two stationary points, E_0 becomes unstable, and E_r is stable.

Summarizing the results of this simple analysis of the ODE model, we can conclude that there are three possible situations. In the case of low bad cholesterol concentration, the healthy state is globally stable and the disease does not develop. For intermediate values of this concentration, the healthy state remains stable but only locally, and a disease state appears. The transition from healthy to the disease state occurs for sufficiently large initial perturbations. Finally, if the concentration of ox-LDL is high, the healthy state is unstable, and transition to the disease state occurs for any small perturbation.

Fig. 4. Schematic representation of the development of atherosclerosis in the blood vessel walls. Bad cholesterol (ox-LDL) penetrates there and initiates an immune response. Monocytes are stopped by activated receptors of endothelial cells and penetrate into the intima. They are transformed into macrophages and then into foam cells leading to a self-amplifying process of chronic inflammation. Reprinted from [20] with permission.

Fig. 5. Nullclines of system (3.1) (left). There are three stationary points, healthy state E_0 and disease state E_r are stable, the intermediate point E_1 is unstable. Wave propagation in the bistable case (right). The initial condition should be sufficiently large to initiate wave propagation. Otherwise the solution decays. Reprinted from [25] with permission.

3.2.2. One-dimensional waves

Let us now take into account diffusion of cytokines and random motion of cells. Then instead of the ODE model, we get the reaction–diffusion system of equations:

$$\begin{cases} \frac{dM}{dt} = D_M \frac{\partial^2 M}{\partial x^2} + f_1(A) - \lambda_1 M \\ \frac{dA}{dt} = D_A \frac{\partial^2 A}{\partial x^2} + f_2(A)M - \lambda_2 A \end{cases} \quad (3.2)$$

We will consider this system of equations on the whole real x . This assumption is justified since the inflammation zone is much smaller than the length of blood vessels. System (3.2) satisfies the monotonicity condition (Section 2.2) for positive M and A . Therefore, we can apply the results on the existence and stability of waves.

Similar to the ODE model, behavior of solutions of the reaction–diffusion system of Eqs. (3.2) depends on the values of parameters. If the concentration of ox-LDL is sufficiently low, such that the ODE system has only a healthy stationary point, then this solution remains globally asymptotically stable also as a solution of system (3.2).

In the intermediate case with two stable equilibria, the healthy state E_0 and the disease state E_r , there exists a unique (up to translation in space) traveling wave solution of system (3.2) with the corresponding limits at infinity. In general, the sign of its speed can be positive or negative. The development of atherosclerosis corresponds to the propagation of wave with a positive speed (Fig. 5, right). Furthermore, behavior of solutions depends on the initial condition. Let us recall (Section 2) that the wave

speed is positive if and only if there exists a pulse solution. If the initial condition is larger than the pulse (for all x), then the perturbation will grow and converge to the wave. If the initial condition is less than the pulse (also for all x), then the solution will decay. The former corresponds to the disease progression while the latter to the return to the healthy state. Thus, the pulse solution determines the critical conditions for the disease initiation.

Finally, if the concentration of ox-LDL is sufficiently high, such that we get the monostable case, the onset of disease occurs for any small initial perturbation. The solution converges to the disease state E_r , and the two waves propagate in the opposite direction implying growth of the atherosclerotic plaque.

3.2.3. Two-dimensional waves

The one-dimensional model considered above neglects the width of the blood vessel wall and the distribution of cells and concentrations inside the wall. In a more complete setting we consider the blood vessel wall as an infinite two-dimensional strip $-\infty < x < \infty$, $0 \leq y \leq h$. Its upper boundary $y = h$ corresponds to the endothelial layer separating the blood flow from the intima. The equations for the concentration of monocytes and inflammatory cytokines become as follows:

$$\frac{\partial M}{\partial t} = d_M \Delta M - \beta M, \quad (3.3)$$

$$\frac{\partial A}{\partial t} = d_A \Delta A + f(A)M - \gamma A. \quad (3.4)$$

The first equation does not contain the production term since the monocyte enter the domain through the boundary, and they are not supposed to proliferate. The right-hand side of Eq. (3.4) contains conventional diffusion term, production of inflammatory cytokines and their elimination. The boundary conditions

$$y = 0 : \frac{\partial M}{\partial y} = 0, \quad \frac{\partial A}{\partial y} = 0, \quad y = h : \frac{\partial M}{\partial y} = g(A), \quad \frac{\partial A}{\partial y} = 0 \quad (3.5)$$

describe the influx of cells through the boundary depending on the concentration A . Here $f(A)$ and $g(A)$ are some growing functions such that $f(0) = g(0) = 0$. This model can be reduced to the one-dimensional model (3.2) by averaging with respect to the variable y .

Behavior of solutions of problem (3.3)–(3.5) depends on the existence and stability of solutions of the problem in the cross section:

$$d_M M'' - \beta M = 0, \quad d_A A'' + f(A)M - \gamma A = 0 \quad (3.6)$$

$$M'(0) = A'(0) = 0, \quad M'(h) = g(A), \quad A'(h) = 0. \quad (3.7)$$

Here prime denotes the derivative with respect to y . It can be easily verified that this problem has a zero solution, $M_+ = A_+ = 0$. Suppose that it has a positive solution $M_-(y)$, $A_-(y)$ such that there are no other solutions between these two solutions. Furthermore, we assume that the eigenvalue problem obtained as linearization of problem (3.6), (3.7) about the solution M_+, A_+ has a positive eigenvalue, and all eigenvalues of the problem linearized about $M_-(y)$, $A_-(y)$ lie in the left-half plane. Then this is a monostable case, and the wave existence is proved for all values of the speed greater than or equal to the minimal speed [20, 26]. The latter admits a minimax representation. In the bistable case, where both eigenvalue problems have negative principal eigenvalues, the wave existence is proved under some restrictive conditions where the existence and stability of solutions of problem (3.6), (3.7) can be controlled [15–17].

3.2.4. Cellular automata

The 2D models discussed in the previous paragraph take into account the width of the vessel wall but not its change due to accumulation of foam cells. We will consider now another approach to atherosclerosis modeling based on cellular automaton [27].

The 2D computational domain represents the luminal surface of a flattened area of the wall of a large artery. It comprises an array of computational cellular automata, each numerical cell encoding the content of the arterial wall. At each point, the content of biological cells in the arterial wall beneath is simulated, and has a content of macrophages, smooth muscle cells and extra cellular lipid. Initially, there is a small quantity of macrophages randomly distributed in the computational domain. The probability of recruitment of new macrophages and of muscle cells is a given growing function of the quantity of macrophages already present in the numerical cell. Furthermore, macrophage death results in the extra-cellular lipid accumulation. Finally, the presence of smooth muscle cells downregulates the recruitment of new macrophages.

An example of numerical simulations is shown in Fig. 6. There are growing plaques viewed from above in the upper image, and a cross section of a plaque in the lower image. Plaque growth observed here is qualitatively similar to the reaction–diffusion waves in the continuous models discussed above.

3.2.5. The influence of hemodynamics

The interaction of hemodynamics with atherosclerosis is beyond the scope of this review. Let us briefly mention some important factors. At the initiation stage, blood pressure influences the penetration of LDL in the vessel wall [21,22], and hypertension is one of the important risk factors. Flow vorticity and low shear rate also stimulate the development of atherosclerosis. However, the most important effect of hemodynamics is flow – plaque interaction with a possible plaque rupture [28–32].

4. Blood coagulation and thrombosis

4.1. Physiological mechanism of clot formation

In the case of blood vessel damage, bleeding is terminated by two interdependent mechanisms: blood coagulation in plasma leading to the formation of fibrin clot and platelet aggregation. Reactions of blood coagulation in plasma propagate in space as a reaction–diffusion wave. Though platelet aggregation has a primary importance for the clot formation, their role in the reaction wave propagation is not yet well studied, and we will restrict ourselves to a brief overview of the mechanisms of platelet aggregation.

4.1.1. The coagulation cascade

The reaction of fibrin polymerization appears at the final stage of the proteolytic enzymatic cascade where the activated clotting factors act as catalysts for activation of the others [33,34]. The key enzyme of the coagulation cascade is thrombin as it catalyzes fibrinogen conversion to fibrin and the distribution of the thrombin concentration in blood plasma has crucial influence on the kinetics of the clot formation [33,35,36]. Formation of thrombin results from the prothrombin activation in the coagulation cascade. This process can be launched by the tissue factor expressed to the blood flow in case of the endothelium rupture (extrinsic pathway), or through the activation of factor XII which triggers activation of factor XI in case of the contact with the foreign surface (intrinsic pathway) [34,37,38]. Both pathways lead to the activation of factor X that contributes to the prothrombin conversion to thrombin [34]. Once the thrombin concentration reaches the threshold value, further prothrombin activation takes place due to the positive feedback loops of the coagulation cascade [34,37,39] (Fig. 7).

4.1.2. Platelet aggregation

Although models of plasma coagulation are important for the understanding of hemostasis mechanisms in veins and in vitro, the basis of arterial thrombosis is the attachment of blood platelets to the vessel wall injury [40–42]. Platelets are the principal components of the hemostatic plug formed to arrest bleeding after an injury to the vascular wall. Attachment of platelets and their accumulation into thrombus is regulated by mechanical interactions with erythrocytes and vessel wall, by numerous chemical agents such as thrombin, or ADP, or prostaglandins, or collagen, as well as by an enormous network of intracellular signaling. The complexity of platelet aggregation and activation during thrombosis makes it an challenging goal for computational modeling: the problem involves complex hydrodynamics, cell adhesion mechanics, chemical cascades and a variety of different interplays between these phenomena [43], some of which are yet to be understood. Since we will not consider platelet aggregation in the reaction–diffusion wave of blood coagulation, we limit ourselves here by this short description of the role of platelets in clot formation (see [44] for a more detailed review).

Fig. 6. Upper part:- Simulation of arterial wall showing development of fatty streaks and advanced atherosclerotic plaques. Normal arterial wall — red, macrophages — yellow, smooth muscle cells — green. Simulation run for 300 cycles with default parameters. Note smooth muscle cell-free borders of plaques. Lower part:- cross section of lesion as indicated by line in upper part. Components are stacked arbitrarily. Smooth muscle cells — blue, macrophages — green, lipid — yellow. Note central location of fibrous cap and lipid. Reproduced with permission from [27]. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

4.1.3. Initiation and termination mechanisms

The initiation of blood coagulation in case of blood vessel wall damage occurs due to the interaction of the tissue factor with factor VII (Fig. 8). They form a complex that activates factors IX and X, which, in their turn, lead to the conversion of prothrombin into thrombin. When the initial concentration of thrombin exceeds some critical quantity, its further production takes place due to the self-accelerating positive feedback through the factors V, VIII, and IX.

In the case of blood coagulation in flow, especially in large arteries, the initiation of coagulation reactions can be downregulated by the flow removing blood factors from the injury site. Platelet aggregate forming at the injury site decreases the flow speed and allows the reactions to start [45,46].

After the initiation and propagation stages of coagulation reactions, the crucial question is about coagulation arrest. There are three main mechanisms inhibiting coagulation cascade: thrombin elimination by antithrombin, protein C pathway, and blood flow. We will return to them below.

4.1.4. Bleeding disorders and thrombosis

Insufficient blood coagulation leads to bleeding disorders, such as hemophilia, and excessive blood coagulation to thrombosis. In the first case, the clot may not form because of the deficiency at the stage of initiation, or it can form with a low rate. We will see below that blood coagulation represents a reaction-diffusion wave in the bistable case. Therefore, normal clotting implies that

the initial condition is sufficiently large, and the speed of the wave belongs to its normal range. If the initial condition is less than some critical value, which will be determined below, then the wave will not propagate. If the wave speed is too small, then the final clot size will be less than it is supposed to be. On the contrary, if the speed is too large, then the clot size will exceed the normal value possibly leading to vessel occlusion. Normal functioning of hemostasis depends on other physiological conditions. In particular, chronic inflammation and cancer represent pro-coagulatory factors often leading to thrombosis. In the next section we will determine critical conditions of blood coagulation, the speed of reaction-diffusion wave of blood coagulation, and the conditions of clot growth arrest.

4.2. Modeling of blood coagulation in quiescent plasma

modeling of blood coagulation implies consecutive stages beginning from thrombin generation curves described by kinetic equations [47–56]. Introduction of the spatial variable allows the description of spatiotemporal evolution of blood factors (see below). modeling of in vivo blood coagulation requires consideration of blood flow [57–64] and blood cells (see [44] and the references therein). We will consider in more detail spatial models of blood coagulation without and with flow but without blood cells.

Fig. 7. The main activation reactions of the intrinsic pathway of the coagulation cascade. Thrombin (IIa) catalyzes activation of factors V, VIII, XI; factors XIa and IXa catalyze activation of factors IX and X respectively; factors VIIIa and Va form active complexes with factors IXa and Xa respectively and further increase thrombin production.

Fig. 8. Schematic representation of the coagulation initiation (left) and termination with activated protein C (right). Factor VII in plasma forms a complex with tissue factor located at the damaged vessel wall. This complex activates factors IX and X that allow the conversion of prothrombin into thrombin. Thrombin produced near the injury site diffuses to the intact walls where it forms a complex with thrombomodulin. This complex activates protein C which downregulates thrombin production.

4.2.1. Reaction–diffusion model for the coagulation cascade

Self-sustained propagation of thrombin in plasma proceeds with a constant speed and it can be considered as a traveling wave solution of the system of partial differential equations describing the coagulation cascade [65–68]. Consider the simplified model:

$$\frac{\partial v_1}{\partial t} = D \frac{\partial^2 v_1}{\partial x^2} + k_1 T - h_1 v_1, \quad (4.1)$$

$$\frac{\partial v_2}{\partial t} = D \frac{\partial^2 v_2}{\partial x^2} + k_2 T - h_2 v_2, \quad (4.2)$$

$$\frac{\partial v_3}{\partial t} = D \frac{\partial^2 v_3}{\partial x^2} + k_3 T - h_3 v_3, \quad (4.3)$$

$$\frac{\partial v_4}{\partial t} = D \frac{\partial^2 v_4}{\partial x^2} + k_4 v_3 - h_4 v_4, \quad (4.4)$$

$$\frac{\partial v_5}{\partial t} = D \frac{\partial^2 v_5}{\partial x^2} + k_5 v_4 + k_1^c v_2 v_4 - h_5 v_5, \quad (4.5)$$

$$\frac{\partial T}{\partial t} = D \frac{\partial^2 T}{\partial x^2} + (k_6 v_5 + k_2^c v_1 v_5) \left(1 - \frac{T}{T_0}\right) - \sigma T \quad (4.6)$$

obtained as an approximation of the more complete model of blood coagulation in quiescent platelet free plasma [66,67]. Here T denotes thrombin concentration, T_0 is its maximal concentration equal to the initial concentration of prothrombin. The variables v_i , $i = 1, \dots, 5$ correspond to the concentrations of active forms of the factors V, VIII, XI, IX and X, respectively. The coefficients k_i denote activation reaction rates, and h_i inhibition rates. Factors VIIIa and IXa form the intrinsic kinase complexes [VIIIa, IXa], while factors Va and Xa form the prothrombinase complex [Va, Xa]. They further accelerate thrombin production. They enter Eqs. (4.5) and (4.6) in the form $k_1^c v_2 v_4$ and $k_2^c v_5 v_1$ obtained under the assumption of detailed equilibrium. This system of equations should be completed by the equations of fibrin production from fibrinogen (activated by thrombin) and fibrin polymer production from fibrin. These equations do not influence the equations of thrombin production, so that system (4.1)–(4.6) can be considered independently.

For further presentation, it is convenient to write system (4.1)–(4.6) in the vector form

$$\frac{\partial \mathbf{u}}{\partial t} = D \frac{\partial^2 \mathbf{u}}{\partial x^2} + \mathbf{F}(\mathbf{u}), \quad (4.7)$$

where $\mathbf{u} = (v_1, \dots, v_5, T)$ is the vector of concentrations and $\mathbf{F}(\mathbf{u})$ is the vector of reaction rates in the equations above. We can analyze this system directly and also use the method of reduction where some of the variables are expressed through the others under the assumption of detailed equilibrium. In this case we will obtain a scalar equation instead of the system of equations.

4.2.2. Existence and stability of waves

In the biologically relevant case, the vector-function $\mathbf{F}(\mathbf{u})$ in (4.7) has three zeros, $\mathbf{u}_+ = 0$, $\mathbf{u}_0$, and $\mathbf{u}_-$. The matrices $\mathbf{F}'(\mathbf{u}_\pm)$ have all eigenvalues in the left-half plane, and the matrix $\mathbf{F}'(\mathbf{u}_0)$ has a positive eigenvalue [69,70]. In this case the thrombin free equilibrium $\mathbf{u}_+$ is stable. Coagulation begins if the initial thrombin concentration exceeds some threshold level.

Let us note that the vector-function $\mathbf{F}(\mathbf{u})$ satisfies the following condition:

$$\frac{\partial F_i}{\partial u_j} \geq 0, \quad i, j = 1, \dots, 6, \quad i \neq j. \quad (4.8)$$

Therefore, if we consider system (4.7) on the whole axis, then it has a traveling wave solution $\mathbf{u}(x, t) = \mathbf{w}(x - ct)$ [3,69–71]. Here c is the wave speed, and the function $\mathbf{w}(x)$ satisfies the equation

$$D\mathbf{w}'' + c\mathbf{w}' + \mathbf{F}(\mathbf{w}) = 0, \quad \mathbf{w}(\pm\infty) = \mathbf{u}_\pm. \quad (4.9)$$

This wave is unique up to translation in space and it is globally asymptotically stable in the following sense. Consider system (4.7) with the initial condition $\mathbf{u}(x, 0) = \mathbf{u}_0(x)$ such that it is a monotonically decreasing (component-wise) function with the limits $\mathbf{u}_0(\pm\infty) = \mathbf{u}_\pm$. Then the solution converges to the wave, $\mathbf{u}(x, t) \rightarrow \mathbf{w}(x - ct)$ as $t \rightarrow \infty$.

Convergence to the wave may not hold if the initial condition $\mathbf{u}_0(x)$ is not a monotone function. As it is usually the case for

the bistable reaction–diffusion systems, it should exceed some threshold. System (4.7) has a stationary pulse solution, that is a positive stationary solution $\mathbf{w}_0(x)$ with the zero limits at infinity [10,11,70]. Moreover, such solution exists if and only if the wave speed is positive. If $\mathbf{u}_0(x) > \mathbf{w}_0(x)$ for all x , then the solution grows and converges to the wave. If $\mathbf{u}_0(x) < \mathbf{w}_0(x)$ for all x , then the solution decays and vanishes. Thus, the pulse solution determines the initial thrombin concentration that provides clot growth.

Thus, we obtain two conditions of clot growth. The first one is the condition on the parameters which provides the positiveness of the wave speed. This condition is equivalent to the existence of the pulse solution. The second condition concerns the initial thrombin concentration. It should exceed the pulse solution. Hence we need to determine the wave speed and the conditions of wave initiation.

4.2.3. Reduction to the scalar equation and the speed of wave propagation

Under the assumption of detailed equilibrium (large reaction constants) we express the variables $v_1, \dots, v_5$ from Eqs. (4.1)–(4.5):

$$\begin{aligned} v_1 &= \frac{k_1}{h_1} T, \quad v_2 = \frac{k_2}{h_2} T, \quad v_3 = \frac{k_3}{h_3} T, \quad v_4 = \frac{k_3 k_4}{h_3 h_4} T, \\ v_5 &= \frac{k_3 k_4}{h_3 h_4} T \left(\frac{k_5}{h_5} + \frac{k_1^c k_2}{h_2 h_5} T \right) \end{aligned} \quad (4.10)$$

and substitute them into the last equation. We get the single reaction–diffusion equation

$$\frac{\partial \theta}{\partial t} = d \frac{\partial^2 \theta}{\partial x^2} + b \theta^3 (1 - \theta) - \sigma \theta \quad (4.11)$$

for the dimensionless thrombin concentration $\theta = T/T_0$, where

$$b = \frac{k_1 k_2 k_3 k_4 k_1^c k_2^c}{h_1 h_2 h_3 h_4 h_5} T_0^2$$

and some small terms are neglected. From the mathematical point of view this means that we consider a singular perturbation problem with a small parameter ϵ in all equations except for the last one: $F_i(u) = F_i^0(u)/\epsilon$, $i = 1, \dots, 5$. Since we are interested in the speed of wave propagation, we need to show that the speed c_ϵ for system (4.7) converges to the speed c_0 for Eq. (4.11). From the minimax representation (2.8) of the wave speed we get the estimate

$$\inf_{x,i} \frac{d_i \rho_i'' + F_i(\rho)}{-\rho_i'} \leq c_\epsilon \leq \sup_{x,i} \frac{d_i \rho_i'' + F_i(\rho)}{-\rho_i'}$$

for any test function ρ . Choosing an appropriate test function, we can show that

$$c_0 + \epsilon \inf_x H(x) \leq c_\epsilon \leq c_0 + \epsilon \sup_x H(x),$$

where $H(x)$ is a bounded function independent of ϵ [69,70]. This estimate provides the desired convergence.

The wave speed c for system (4.1)–(4.6) converges to the speed c_0 for the single equation (4.11) in the limit of large reaction rate constants (coefficients in the first five equations) [69,70]. The latter can be estimated by the dimensionality considerations:

$$c_0 \sim \sqrt{bD} \left(\alpha - \frac{\sigma}{b} \right),$$

where $\alpha \approx 0.1$ is a number chosen from the condition that the speed vanishes if the integral of the nonlinearity equals 0. More precise but qualitatively similar formula can be obtained by a narrow reaction zone method and by a piecewise linear approximation [70]. They give a good approximation for the speed of the complete system and of the experimental results (Fig. 9).

4.2.4. 2D models and clot growth arrest

Blood coagulation is inhibited via TFPI (tissue factor pathway inhibitor) present at the surface of endothelial cells, antithrombin, which is present in blood plasma, and protein C, which is activated at the intact surface by a complex of thrombin and thrombomodulin and then diffuses in the vessel lumen. The concentration of antithrombin should be sufficiently low in order to allow the initiation and propagation of the thrombin wave. TFPI regulates the activation of the coagulation cascade at the vessel wall. Once propagation commences, neither of them (antithrombin, TFPI) cannot stop it. It can be verified that in the 1D model activated protein C does not stop the wave propagation either. Therefore, in order to describe clot growth arrest via activated protein C we need to consider 2D or 3D formulations, with an injured part of the wall where thrombin production is initiated and an intact part of the wall where protein C is activated. Protein C can then diffuse to the injured site and decelerate thrombin production. There is competition between self-sustained thrombin production and thrombin induced protein C activation, which downregulates thrombin production. For fixed values of other parameters, the result of this competition is determined by the injury size. If the injury size is sufficiently small, then the diffusion of activated protein C to the damaged site will be large enough to arrest thrombin production. If the injury is sufficiently large, then protein C does not stop wave propagation.

Thus, there are three regimes of clot growth in quiescent plasma: initiation without propagation, initiation with propagation and with growth arrest and initiation with propagation without growth arrest [73].

4.3. The influence of blood flow

There is a vast literature devoted to the interaction of blood coagulation with hemodynamics (see, e.g., [44,57–63]). A conventional model describing this process consists of a reaction–diffusion system of equations coupled with the Navier–Stokes equations. The latter can contain some additional terms describing the deceleration of fluid inside the clot [74–76]. An example of numerical simulations is shown in Fig. 10. Detailed analysis of this question is beyond the scope of this paper, and we will limit ourselves to some aspects related to the propagation of reaction waves.

The equation for the thrombin concentration with convection becomes as follows:

$$\frac{\partial \theta}{\partial t} + \mathbf{v} \cdot \nabla \theta = D \Delta \theta + f(\theta), \quad (4.12)$$

where $\mathbf{v} = (v_x, v_y)$ is the velocity vector, $f(\theta) = b\theta^3(1 - \theta) - \sigma\theta$, x is the variable along the axis of the cylinder, y is the orthogonal variable. If we neglect the influence of clot on blood flow and suppose that v_x is a given (e.g., parabolic) function of y , and $v_y = 0$, then existence and stability of waves described by Eq. (4.12) can be studied [3,77]. However, these qualitative results do not allow us to get analytical estimates of the speed of clot growth and of its size.

In order to get analytical estimates of clot growth, we consider the simplified equation for thrombin concentration

$$\frac{\partial \theta}{\partial t} = D \frac{\partial^2 \theta}{\partial y^2} + \Phi(\theta, y) \quad (4.13)$$

obtained as an approximation of Eq. (4.12) after its integration along the axis of the flow. Here,

$$\Phi(\theta, y) = b\theta^3(1 - \theta) - \sigma(y)\theta, \quad (4.14)$$

and $\sigma(y) = \sigma_0 + \sigma_1 v_x(y)$. It is similar to Eq. (4.11) with an additional term in $\sigma(x)$. The first term in the expression for σ

Fig. 9. Left: speed of the thrombin wave propagation (mm/min) as function of percentage of factor IX activity. Dots: experimental data [72]; dashed line: narrow reaction zone approximation; dash-dot line: piecewise linear approximation. Right: thrombin distribution at the consecutive moments of time for the complete model (a) and for the single equation (4.11) (b). Thrombin profile becomes monotone for the one-equation model because the equation for prothrombin is eliminated. Despite this difference, the speeds of propagation remain close. Reprinted with permission from [70].

Fig. 10. Snapshots of thrombin distribution (left) and fibrin distribution (right) in the beginning of clot growth (upper row) and before complete occlusion (lower row). Reprinted from [75] with permission.

takes into account the inhibition by antithrombin, the second term describes the loss of thrombin due to blood flow. Velocity profile $v_x(y)$ can be parabolic or it can take into account the influence of clot on flow velocity. For simplicity of notation we denote here by y the variable perpendicular to the injured vessel wall.

Set $I(y) = \int_0^{\theta_*(y)} \Phi(\theta, y) d\theta$, where $\theta_*(y)$ is the largest zero of the function $\Phi(\theta, y)$ for fixed y . In the case without flow, where the function and the integral do not depend on y , condition $I > 0$ provides the positiveness of the wave speed, which is

the condition of clot growth. Since the thrombin wave front is sufficiently narrow, then a similar condition can be applied for every fixed x . Therefore, thrombin wave propagates for the values of y where $I(y) > 0$. Condition $I(y) = 0$ determines the maximal clot size if it is less than H . Fig. 11 shows the maximal clot size as a function of the maximal flow velocity obtained in direct numerical simulations of the 2D problem and in the approximate 1D equation (4.13). The analytical condition through the sign of the integral and numerical solution of the simplified

Fig. 11. The final height of the clot in case of different values of the initial blood flow velocity. Left: numerical simulations for different values of antithrombin concentration A in plasma. Right: numerical (solid line) and analytical (dashed line) solutions of the simplified model for $A = 0.4$. For both models, complete occlusion occurs for low flow velocities and partial occlusion for high velocities with a sharp transition between these two regimes. Reprinted from [76] with permission.

equation give a good approximation of the final clot size and of the conditions of complete occlusion.

4.4. Regimes of blood coagulation

We can now summarize the results presented above and determine the regimes of blood coagulation in relation with different physiological conditions. We will explain them with the help of the simplified model (4.13) where the function

$$\Phi(\theta, y) = (b\theta^3 + \tau e^{-\mu y})(1 - \theta) - \sigma(y)\theta$$

contains additional exponential term (see (4.14)) that describes the initiation of coagulation at the vessel wall. The function $\tau \exp(-\mu y)$ gives the stationary distribution of activated factor IX which initiates production of thrombin. The constant τ depends on the expression of the tissue factor at the endothelial cells. Eq. (4.13) is considered on the half-axis $y \geq 0$ with the no-flux boundary condition $\theta'(0) = 0$.

In order to determine the regimes of blood coagulation, let us briefly analyze behavior of solutions of Eq. (4.13). During the initiation stage, Eq. (4.13) in its stationary form can be approximated by the equation

$$d\theta'' - \sigma_0\theta + \tau e^{-\mu y} = 0, \quad \theta'(0) = 0. \quad (4.15)$$

We neglect θ^3 since the concentration of thrombin is small, $\theta \ll 1$, and we neglect the flow velocity near the boundary. We get the solution

$$\theta_0(y) = \frac{\tau}{\lambda(\sigma_0 - d\mu^2)} (\lambda e^{-\mu y} - \mu e^{-\lambda y}),$$

where $\lambda = \sqrt{\sigma_0/d}$.

On the other hand, neglecting the initiation term, we obtain the equation

$$d\theta'' + b\theta^3(1 - \theta) - \sigma_0\theta = 0, \quad \theta'(0) = 0. \quad (4.16)$$

It can be explicitly verified that if the inequality

$$\int_0^{\theta_*} (b\theta^3(1 - \theta) - \sigma_0\theta) d\theta > 0 \quad (4.17)$$

holds, then problem (4.16) has a positive decaying solution $\theta_p(y)$. Here θ_* is the maximal solution of the equation $\theta^2(1 - \theta) = \sigma_0/b$ assuming that it exists.

The behavior of solution of Eq. (4.13) ($\tau = 0, \sigma_1 = 0$) depends on the initial condition $u(y, 0)$. If $u(y, 0) \geq \theta_p(y)$ for all $y \geq 0$, then the solution will converge to the traveling wave solution whose speed is positive due to condition (4.17). If $u(y, 0) \leq \theta_p(y)$, then the solution converges to zero. If none of the inequalities hold, then we cannot make a precise conclusion about the behavior of solution. However, the qualitative conclusion is that for $u(y, 0)$ sufficiently large, the solution grows, if the initial condition is sufficiently small, then the solution decays.

4.4.1. Initiation without propagation

The initial quantity of thrombin $\theta_0(y)$ produced at the initiation stage should be compared with the critical solution $\theta_p(y)$. If $\theta_0(y)$ is sufficiently small, then solution of Eq. (4.13) also remains small, and clot does not grow. In particular, this is the case for small values of τ (tissue factor expression).

There are two main physiological situations where initiation without propagation can be realized. The first one is venous thrombosis where a large part of a vein can contain chronic thrombus. The initiation of coagulation can occur due to the inflammation of vessel wall leading to the expression of tissue factor by endothelial cells. Let us stress that this regime of blood coagulation does not depend on the size of the damaged area.

The second situation concerns severe hemophilia where condition (4.17) does not hold since parameter b is too small. In this case, the wave speed is negative, and clot will not grow for any initial condition.

4.4.2. Initiation with propagation and with growth arrest

If $\theta_0(y)$ is sufficiently large, then the initiation stage is followed by the propagation stage. When the thrombin wave goes from the boundary inside the vessel, blood flow velocity increases, and it should be taken into account. Instead of condition (4.17) we have now condition

$$\int_0^{\theta_*} (b\theta^3(1 - \theta) - \sigma(y)\theta) d\theta > 0$$

with a space dependent function $\sigma(y)$. We assume here that the width of the thrombin gradient is sufficiently narrow with respect to the vessel radius R . This condition should be verified for each y fixed. The maximal value of y_m for which it holds determines the final clot size (Fig. 11, right). If $y_m < R$, then blood flow stops clot growth. Otherwise it fills completely the vessel cross section leading to the complete occlusion. Thus, condition

$$\int_0^{\theta_*} (b\theta^3(1 - \theta) - \sigma(y_m)\theta) d\theta \leq 0 \quad (4.18)$$

determines the regime of normal functioning or partial occlusion. This can be also the case of weak hemophilia. The difference between normal case and pathologies is determined by the value of parameter b which includes the reaction rate constants.

Let us note that activated protein C represents another mechanism of clot growth arrest not taken into account by condition (4.18). As discussed above, we need to consider a 2D formulation in order to describe it.

4.4.3. Initiation with propagation and without growth arrest

If condition (4.18) is not satisfied, then clot growth is not limited by flow. It fills the whole vessel cross section leading to

Fig. 12. Schematic representation of the conduction pathway for the heart.

complete vessel occlusion. This situation can occur in the case of local occlusion and for disseminated coagulation specific for sepsis where large areas of the endothelial layer are damaged. Similar to the initiation without propagation, this limiting case is independent of the injury size.

We remark that condition (4.18) does not take into account the variation of the flow velocity during clot growth. Flow velocity can increase or decrease during clot growth depending on the clot size. Some analytical estimates can be obtained in the approximation of Poiseuille flow [44].

5. Basic principles of cardiac arrhythmia modeling

5.1. Normal function of the heart

5.1.1. Physiology of the heart

The pumping of the heart is controlled by electrical waves of excitation, which propagate through the heart and initiate cardiac contraction. As in all mammals, the human heart contains four chambers. The lower two chambers, called ventricles, pump blood to the lungs and body. Therefore the cardiac ventricles have thick walls: the right ventricle of about 0.3–0.5 cm, the left ventricle of about 0.7–1.1 cm. The upper two chambers of the heart are called the atria. The left and right atria respectively receive blood from the lungs and the venous circulation and contraction of the atria helps to deliver it to the ventricles. Compared to the ventricles, less mechanical power is required, which is why the atria have thinner walls, of about 0.3 cm.

5.1.2. Excitation sequence in sinus rhythm

The normal excitation of the heart is initiated at its natural pacemaker, the sinoatrial (SA) node, which is located in the right atrium (Fig. 12). The cells of the SA node can spontaneously depolarize and produce an electrical excitation wave. From the SA node, the wave propagates through the atria by depolarization of the atrial muscle cells (myocytes) and initiates their contraction. The atria and ventricles are not electrically coupled to each other, except for one region called the atrioventricular (AV) node. However, wave propagation through the AV node is very slow, which causes some delay between excitation of atria and ventricles. As a result, the blood is delivered from atria before ventricular contraction. Excitation of the ventricles progresses from the AV node through the ventricular conduction system (Purkinje network) and exits into ventricle at multiple locations to ensure a fast excitation, which in normal conditions should yield nearly synchronous mechanical contraction of the cardiac ventricles.

5.2. Brief overview of cardiac arrhythmias

The abnormal excitation of the heart results in cardiac arrhythmias. The most important arrhythmias are associated with an abnormally fast heart rate and are called tachycardia. Because the atria and ventricles are coupled only via the AV node, most cardiac arrhythmias occur either in the atria or in the ventricles of the heart.

The excitation of the heart can also become spatially disorganized. Since mechanical contraction of heart cells is triggered by the electrical depolarization waves, a spatially disorganized electrical activation pattern will cause contraction failure. If this occurs in the ventricles, it will result in cardiac arrest and sudden cardiac death. Sudden cardiac death due to ventricular fibrillation (VF) is one of the largest causes of death in the industrialized world and accounts for approximately 1 death in 10 [78]. In the first minutes of VF, it is possible to apply a strong electric shock to the heart, to defibrillate it and restore spatial synchronization. Current methods for defibrillation use electrodes on the heart or on the body surface and release a relatively large amount of energy, which can be painful and damage the heart [79,80]. VF also has another important consequence: more than half of all drug withdrawals in recent years were due to cardiac side effects contributing to the onset of cardiac arrhythmias and sudden cardiac death [81].

Chaotic excitation patterns that occur in the cardiac atria give rise to so-called atrial fibrillation (AF). Although the atria of the heart do not contract under this condition, AF does not lead to sudden cardiac death, because blood can still enter the ventricle during its relaxation. In normal conditions, around 70%–80% of the volume of the left ventricle is filled passively, and the ‘atrial kick’ accounts for only 20%–30%. Unfortunately, in the absence of active contraction, the blood flow in the atria is turbulent and much slower than in normal conditions. Therefore, AF induces stagnation of blood and creates conditions for thrombus formation. If a thrombus formed in the atria enters the circulatory system, this may also be life-threatening, e.g. via occlusion of coronary arteries, pulmonary embolism or stroke. In relation to the latter, AF causes a five fold increase in stroke risk [2]. Atrial fibrillation is probably one of the most important and challenging problems in modern cardiac electrophysiology, as it is the most common sustained cardiac arrhythmia encountered in clinical practice. It is estimated that AF occurs in approximately 3% of adults aged 20 years or older. Its incidence rapidly increases with age and it is estimated that one in four middle-aged adults in Europe and the US will develop AF during their life [82].

5.3. Rotor formation

The mechanisms of cardiac arrhythmias are directly related to abnormal wave propagation in the heart. Although electrical waves in the heart are also described by the reaction–diffusion equations, it will be easier to introduce the main abnormal regimes using simpler, rule-based models. These models are based on very well established properties of cardiac cells, but represent them in a simplified way. We assume the following: (1) the cells can be excited by their neighbors forming a propagating excitation wave with velocity v , (2) some time after excitation, the cell cannot be excited again. This time period is called refractory period and will be denoted R .

In such systems, we can observe the formation of sustained rotational activity. First, consider an inexcitable portion of the cardiac wall, e.g. scar tissue or a vein or artery leaving the heart. If the perimeter of this obstacle is L then the period of rotation will be $T = L/v$. If this period is longer than the refractory period R the wave can rotate around this obstacle and will excite the

heart with period T (Fig. 13a). The minimally possible value of T is R , i.e. about 250 – 300 ms. This value is 3–5 times lower than normal heart rate, leading to a heart beat that is 3–5 times faster than the normal heart rhythm. This ‘re-entrant’ activity around an obstacle thus results in a tachyarrhythmia. The ideas of such sources of arrhythmia are known for more than a century [83].

It was also found that such reentrant activity can occur in cardiac tissue without any anatomical obstacle. To this aim, consider initial conditions in the form of a wave break (Fig. 13b). From the rules stated above, it follows that the wave will propagate along the boundary of refractory tissue until the refractory period ends. Then it will make a U-turn and again follow the refractory tail (Fig. 13c), eventually creating a self-sustained source of electrical activation in two spatial dimensions (Fig. 13d). It is easy to see that if one assumes constant propagation velocity of the wave front, the pattern will eventually take the shape of a spiral to keep the rotation period and propagation velocity the same at all distances from the center.

These waves are in different communities known as ‘spiral waves’, ‘electrical vortices’ or ‘cardiac rotors’.

We will discuss the dynamics of spiral waves in more detail in the next sections. Here we note, that because in our example the wave front closely follows the refractory tail of the wave, the rotation period will be close to the minimal possible period R , and thus such sources will produce the fastest heart rhythms and thus the most dangerous cardiac arrhythmia.

5.4. Electrical activity of the heart as a reaction–diffusion system

5.4.1. Cell model

Cardiac cells maintain a difference in the electrical potential between the inside and outside of their cell membrane. This voltage difference V_m is a result of physico-chemical equilibrium between different concentrations of ions inside and outside the cell. The ion that is most important in maintaining the resting state and determining the resting potential is potassium (K^+).

The electrical signaling in the heart follows from a rapid change of the transmembrane voltage which is caused by the flow of other ions through the membrane, mainly Na^+ and Ca^{2+} . These currents are transient and after a while the resting potential is restored by the flux of K^+ ions. The process of excitation and relaxation of a portion of the cell membrane of a single cell can be described by the following ordinary differential equation (ODE):

$$C_m \frac{dV_m}{dt} = -I_{ion}, \quad I_{ion} = I_{Na} + I_K + I_{Ca} + \dots, \quad (5.1)$$

where C_m is the electrical capacity of the cell membrane, V_m the transmembrane voltage over the cell membrane and I_{ion} represent the sum of ion currents which flow through the membrane and change its potential. The description of dynamics of these current requires additional equations, as each of the currents typically depends on V_m , local ion concentrations and time. Usually the time dependency is given by a description of the opening and closing of channels in the cell membrane that allow specific ion species to pass. The state of a local ensemble of such channels is summarized in so-called gating variables, often denoted $g, m, h, \dots$. For example, the potassium current I_K in one of the most widely known models for excitable cells, i.e. the Hodgkin–Huxley model [84], is represented as follows:

$$I_K = G_K n^4 (V_m - V_K), \quad (5.2)$$

$$\frac{dn}{dt} = \frac{n_\infty(V_m) - n}{\tau_n(V_m)}, \quad (5.3)$$

where the gating variable n (scaled between 0 and 1) approaches its voltage-dependent steady state $n_\infty(V_m)$ with characteristic time $\tau_n(V_m)$ (5.3), G_K is a scaling constant, and V_K is a so-called

Nernst potential, i.e. a voltage for which this current is zero. Mathematically V_K gives the equilibrium of (5.1) if only the current I_K would be present. The functions $n_\infty(V_m)$ and $\tau_n(V_m)$ are fitted to experimental measurements and the power 4 in Eq. (5.3) was chosen to give the best fit to the data. The Hodgkin–Huxley model is probably one of the most cited papers in computational electro- or neurophysiology and since it was published in 1952 it has been cited nearly once per-day on average. This paper provided a framework for developing most of the other currently used models for cardiac and nerve cells. In such models, the ion currents are represented in a similar fashion, but of course fitted to other experimental data. Most ionic currents that have been identified in cardiac cells have one or two gating variables and may include additional functions in the right-hand side, e.g. $I_X = f(n_X, V_m, X)$. For example the scaling factor G_X may have dependency on voltage, etc. Some currents can also depend on concentrations. For some channels, especially when there are few channels per surface area on the cell membrane, a more complex Markov state representation may be appropriate [85].

Currently, more than 280 models for various aspects of electrical activity of excitable cells are listed in the online repository CellML [86]. Such models describe the detailed underlying biophysical mechanism of cardiac excitation and are called ionic models. They have different complexity and can include from 3 to more than 100 equations and hundreds of parameters.

Another class of cardiac models are phenomenological models, which can have as few as two ODEs. The first model of that kind was developed by R. FitzHugh in 1961 [87] as a generic two-variable system to describe dynamics of excitation observed in the Hodgkin–Huxley model. Later, many similar models were proposed. A typical form for such models is:

$$\frac{dV}{dt} = f(V, g), \quad (5.4)$$

$$\frac{dg}{dt} = \varepsilon \phi(V, g), \quad (5.5)$$

where in most of the cases $f(V, g)$ can be represented as $f(V, g) = f(V) - g$, where function $f(V)$ has cubic non-linearity, and the function $\phi(V, g)$ is a monotonic function, for example $\phi(V, g) = kV - g$ and $\varepsilon \ll 1$. The most widely used models of this type are those of FitzHugh–Nagumo [87], Rinzel & Keller [88], Barkley [89], Aliev–Panfilov [90], Mitchell–Schaeffer [91], Karma [92], Fenton–Karma [93] and Bueno-Orovio et al. [94]. The non-linear functions used in these models can be found in those references or works that use of these models to study or explain emergent dynamics.

In summary, simplified or detailed ionic models of cardiac cell can be represented as a set of autonomous coupled ODEs:

$$\frac{d\mathbf{u}}{dt} = \mathbf{F}(\mathbf{u}), \quad (5.6)$$

where the state vector $\mathbf{u}$ contains the (possibly normalized) transmembrane voltage V_m as its first element. Other state variables may include ionic concentrations or gating variables or phenomenological descriptions thereof. For non-self-pacemaking cells, the resting state will be a stable equilibrium to Eq. (5.6). For self-pacemaking cells (such as those in the SA node), the non-linear functions in the right-hand side of Eq. (5.6) will be tuned such that it possesses a stable limit cycle.

5.4.2. Anisotropic coupling of cells

The cells in cardiac tissue are electrically connected to each other and their coupling is commonly described by representing the tissue as a resistive network. This approach, however, has to take into account the structural anisotropy of cardiac tissue, which will impact on wave propagation and excitation pattern dynamics.

Fig. 13. Schematic representation of a wave propagating around an obstacle (a) and of spiral wave formation (b–d). White denotes resting state, red is the wave front and green refractory (recovering) tissue that cannot be re-excited yet. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

In a structurally normal heart, cardiac myocytes are organized in myofibers where there is more end-to-end coupling between cells than lateral coupling, resulting in a propagation velocity for electrical signaling that is 2–3 times larger in the myofiber direction than in transverse directions.

The fibers in the heart have a complex spatial organization. In the ventricles they exhibit a pattern, called rotational anisotropy: if we consider a small cube made from cardiac ventricle then at each layer orthogonal to the transmural axis we will have parallel fibers, but the angle of the fibers will rotate when we move along the vertical (transmural) direction. Depending on the region of the ventricle this rotation can be from 90 to 150 degrees. Such anisotropy can be represented by various analytical geometrical models [95–97]. In the mid-ventricular walls, there is additional laminar structure which reduces shear stresses during contraction and allows a significantly higher volume reduction of the cardiac cavities during the pumping of blood [98,99]. In atrial tissue the organization of the anisotropy is much more complex [100].

From homogenization theory applied to an ensemble of cardiac cells, one can obtain the so-called bidomain equations [101–105]:

$$\begin{aligned} \partial_t V_m &= \bar{\nabla} \cdot (\bar{D}_i \cdot \bar{\nabla} V_i) + F_1(V_m, \tilde{\mathbf{u}}), \\ \bar{\nabla} \cdot (\bar{D}_i \cdot \bar{\nabla} V_i) + \bar{\nabla} \cdot (\bar{D}_e \cdot \bar{\nabla} V_e) &= 0, \\ \partial_t \tilde{\mathbf{u}} &= \tilde{\mathbf{F}}(V_m, \tilde{\mathbf{u}}). \end{aligned} \quad (5.7)$$

Here, the original state vector from the cell model (5.6) is written as $\mathbf{u} = [V_m \tilde{\mathbf{u}}]^T$, and $\mathbf{F} = [F_1, \tilde{\mathbf{F}}]^T$, where $\tilde{\mathbf{u}}$ contains ionic concentrations and gating variables, and $\tilde{\mathbf{F}}$ their reaction kinetics. F_1 is the essentially the prescription for transmembrane current.

For the spatial coupling, the transmembrane potential is stated explicitly to the difference between intra- and extracellular potentials: $V_m = V_i - V_e$. All effects of the microstructural anisotropy (e.g. partial volume and shape of fluid compartments, density and direction of gap junctions) are lumped into two effective diffusion tensors $\mathbf{D}_i$ and $\mathbf{D}_e$ for V_i and V_e , respectively. Each of them can be written in a basis of eigenvectors, corresponding with the local principal axes of the tissue:

$$D_c^{jk} = \sum_{a=1}^n D_{a,c} e_a^j e_a^k, \quad c \in \{i, e\}. \quad (5.8)$$

5.4.3. Monodomain model

The need of taking into account separate extra- and intracellular potentials is often relieved by assuming equal anisotropy ratios for the intra- and extracellular domains, i.e. $D_{a,i} = \lambda D_{a,e}$, in which case Eqs. (5.8) reduce to the monodomain equations:

$$\begin{aligned} \partial_t V_m &= \sum_{i,j=1}^n \partial_i (D^{ij} \partial_j V_m) + F_1(V_m, \tilde{\mathbf{u}}), \\ \partial_t \tilde{\mathbf{u}} &= \tilde{\mathbf{F}}(V_m, \tilde{\mathbf{u}}). \end{aligned} \quad (5.9)$$

where $\bar{D}^{-1} = \bar{D}_i^{-1} + \bar{D}_e^{-1}$. Eqs. (5.9) can be concisely written as a multi-component reaction–diffusion system

$$\partial_t \mathbf{u} = \mathbf{P} \sum_{i,j=1}^n \partial_i (D^{ij} \partial_j \mathbf{u}) + \mathbf{F}(\mathbf{u}) \quad (5.10)$$

with the matrix $\mathbf{P} = \text{diag}(1, 0, \dots, 0)$. For $\mathbf{F}(\mathbf{u})$ one can use any of the ionic or simplified cell models discussed above. Numerical studies have shown that, in the absence of external electrical stimulation, numerical solutions to the monodomain equations are good approximations to bidomain solutions [106,107]. Deviations between mono- and bidomain are found under external electrical stimulation (e.g. during defibrillation), high myofiber curvature and as boundary effects [108].

The main idea behind Eq. (5.10) can be easily understood from a 1D system. In that case the spatial current in the system is given by $D \frac{\partial V_m}{\partial x}$. The divergence of this current, namely $\frac{\partial}{\partial x} (D \frac{\partial V_m}{\partial x})$ flows through the membrane and is added to ion currents of the cell. The generalization of this approach gives Eq. (5.10).

The diffusion tensor $\mathbf{D}$ accounts for tissue anisotropy, as the resistivity along cardiac fibers and in the orthogonal directions is different. Often, one only takes into account the effect of myofiber direction $\bar{e}_1$ and thus assumes that $D_2 = D_3$ whence

$$D^{ij} = D_2 \delta^{ij} + (D_1 - D_2) e_1^i e_1^j. \quad (5.11)$$

Although diffusion properties can be expected to vary during cardiac contraction, it is common to take D^{ij} time-independent. For anatomically accurate modeling, the heart's shape can be segmented from medical images, e.g., computed tomography or magnetic resonance imaging (MRI). The myofiber direction can either be added rule-based [95], or interpolated from diffusion-weighted MRI, which is now feasible for both ventricles [109] and atria [110].

If one wishes to model the heart as uniaxial rather than laminar, the laminar structure of the cardiac ventricles can be assessed by MRI imaging [109,111] and structure tensor imaging [112].

5.5. Approaches beyond the mono- and bidomain models

Other approaches than the mono- and bidomain equations have also been proposed and used for cardiac modeling. The effect of resistive coupling with fibroblasts in cardiac tissue has been incorporated in the tridomain model [113]. In recent years, cardiac tissue has also been regarded as a porous medium [114] or as a medium undergoing fractional diffusion [115]. Recently, hyperbolic models were introduced to account for induction effects [116].

The forementioned approaches describe models for the electrophysiology. To study their effect on mechanical contraction of the heart, mono- and bidomain models have been coupled to mechanical equations, sometimes with an electromechanical feedback term that represents stretch-activated channels in the myocyte membranes [117]. Different approaches have been taken for the mechanical equations, including mass–spring models [118]

and finite deformation [119–122]. Since both the bidomain model and finite deformation mechanics are ruled by elliptic rather than parabolic partial differential equations, they can be treated together if they need to be both included in the description; see e.g. [123].

6. Dynamics of wave patterns underlying cardiac arrhythmias

6.1. Numerical approach to cardiac arrhythmia management

Many results on spiral wave dynamic in cardiac models have been obtained using numerical, or combined numerical–analytical approaches, as unfortunately the equations, even for FitzHugh–Nagumo models, resist to most theoretical analysis (although we provide some exceptions in Sections 6.2–6.4). Currently we can distinguish two general directions of research in cardiac modeling: (1) numerical studies for understanding the fundamental processes underlying the dynamics of arrhythmia sources in the heart. Part of this field included the numerical study of the dynamics of spiral waves and processes by which they can be initiated, sustained or terminated; (2) numerical studies with direct clinical or experimental applications.

6.1.1. Numerical studies of fundamental dynamics

The studies of fundamental dynamics have revealed most of the known properties of spiral waves. It was shown that in addition to stationary circular rotation there are more complex regimes of rotation of the tip of the spiral wave, which is called meandering. The meandering of spiral waves was first observed by Zykov [124] in low-dimensional models of excitable media, where he showed that trajectories of the core can have epicycloidal rotation (inward petals) or hypocycloidal rotation (outward petals) with intermediate regimes of drift of the spiral. These types of dynamics turned out to be very general and were later found in many detailed models of cardiac tissue [125,126]. Fig. 14 shows qualitatively different meandering patterns obtained in the three-variable Fenton–Karma model [93]. Typically, when one increases the excitability of the tissue, the meandering pattern changes first from a circular core to inward petals, then to outward petals.

Another important type of dynamics is drift of spiral waves. This drift occurs if the properties of cardiac tissue are not uniform in space. The most simple situation is heterogeneity in the “reaction” terms in Eq. (5.10). It accounts for spatial heterogeneity in properties of cardiac cells which is very well established in the heart [127]. Numerical studies using FitzHugh–Nagumo models show that in case of heterogeneity the spiral wave drifts and the drift has two components, one in the direction of the gradient of the heterogeneity and one in the orthogonal direction [128,129]. The orthogonal component depends on the direction of rotation of spiral waves (i.e. chirality), and the longitudinal component was always directed towards the region where the spiral wave has a longer period. This result was later confirmed in an ionic model of cardiac tissue [130], where the drift was also always directed towards the region of a longer period. Although that effect is very robust and reproducible, we still do not know its mechanism. In addition, drift of spiral waves can also be induced by spatial changes in the anisotropy of cardiac tissue [131,132] and by change in the Gaussian curvature of the surface [133]. Such dynamics of spiral waves found through the numerical experiments turned out to be important. In particular, stable rotation along a circular trajectory or rotation along an obstacle will produce a periodic electrical activity typical for arrhythmias called monomorphic tachycardia, which is characterized by a periodic electrocardiogram (ECG). Meandering or drift has more complex dynamics and results in quasi-periodic or even more complex

activity. Therefore, it has been related to another arrhythmia, called polymorphic ventricular tachycardia characterized by aperiodic ECGs [134–136]. Numerical studies also proposed many mechanisms of initiation of spiral waves and proposed novel methods of spiral wave removal from the heart (see, e.g., the review in [137]).

Another class of important results which were obtained using generic computer modeling was the study of properties of three-dimensional spiral waves, called scroll waves. A scroll wave is just a three-dimensional generalization of the spiral wave and it can be obtained by shifting a 2D spiral wave in the vertical direction (Fig. 15a). The centers of rotation of spiral waves will form a line called filament (Fig. 15b, green).

The dynamics of scroll waves in 3D are often represented by dynamics of their filaments. First numerical studies of scroll waves revealed that their dynamics is affected by the curvature of filament and in case of a circular filament (Fig. 16a) it can either shrink, which leads to annihilation of the vortex (Fig. 16b) or it can expand in space (Fig. 16c) [138]. Such property of the filaments was interpreted as filament tension [139], and the shrinking regime was associated with the positive filament tension, i.e. the situation when the filament tends to decrease its length, or negative filament tension when the filament tends to increase its length.

Filament tension substantially affects the overall dynamics of the scroll waves. If one considers an oblique filament in a slab of cardiac tissue, then in the case of positive tension the filament will try to reduce its length and thus will evolve and stabilizes orthogonal to the boundaries (Fig. 17 the upper panel). In the case of negative filament tension, the length of the filament will continue to grow, it will collide with the boundaries and form new vortices (Fig. 17 the lower panel). Such dynamics was considered as one of possible mechanisms of ventricular fibrillation in a famous paper by A. Winfree [78] who claimed that in most of the cases ventricular fibrillation is a 3D phenomenon (see also [140]). However, it was also shown that if tension is not sufficiently negative compared to the filament rigidity and wall thickness, there is insufficient space for breakup and the filament will undergo Euler buckling instead [141].

Currently, the further study of filament dynamics is an important direction in cardiac modeling. Many other types of filament dynamics were also found and studied using computational and analytical approaches. We will provide examples of that below.

Another important contribution of generic computer modeling studies was finding of the process of spiral breakup and bringing up ideas of dynamical instabilities. It will be considered in details in Section 6.2.

6.1.2. Numerical modeling with direct clinical or experimental applications

In addition to obtaining novel biological insights in many disease conditions, a subdiscipline within cardiac modeling concerns numerical studies in view of direct clinical or experimental applications. Here, the modellers aim to reproduce quantitatively the electrical activity in the heart of an individual patient and using it propose or assist clinical interventions. This is at present a fairly large area of research and it includes a lot of engineering methodology, such as image segmentation (in order to obtain a patient-specific heart model), mesh generation, tuning the model to electrical measurement of the heart and adjustment of the model equations based on these measurements.

After formulation of a patient-specific model, clinical interventions can be planned. With regard to arrhythmia management, cardiac surgery typically consist of implanting pacemakers, cardioversion devices (i.e. implanted defibrillators) or ablation of the arrhythmogenic substrate.

Fig. 14. Spiral tip trajectories in the Fenton–Karma model for two different parameter settings. Low excitability yields a cycloidal tip trajectory (a), while high excitability a star-like or “linear” core (b). Color denotes transmembrane voltage V_m ; the medium is assumed isotropic. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

Fig. 15. (a) In three dimensions, spiral waves are called scroll waves, for their similarity with ancient book scrolls. (b) The rotation axis in every cross-section of the scroll wave is usually called the filament (green). From simulated or numerical data, one often determines the line of spiral wave tips (blue) instead, which will for a circular-core scroll wave revolve around the filament and trace out the spiral wave core. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

Fig. 16. A closed filament in 3D corresponds to a scroll ring. Historically, 2 regimes were found: contracting rings and expanding rings, which correspond to the regimes of positive and negative filament tension. Picture reproduced with permission from Panfilov & Rudenko, *Physica D* 28, p. 215–218, 1987.

During an ablation procedure, the cardiologist inserts one or several catheters through the patient's blood vessels into the cavities of the heart. The catheters can record electrical activity and can also destroy tissue at desired location. The main idea behind this procedure is to destroy part of the tissue which are essential for rotational activity or focal arrhythmia sources, such that after the arrhythmia will not be initiated or sustained. In recent studies various modes of imaging and patient-specific modeling were combined to predict the best places for ablation for the individual patients [142,143].

To date, an increased performance compared to existing clinical methods has been suggested for the prediction of ventricular tachyarrhythmias and in post-myocardial infarction patients, where the anatomical substrate is modulating the arrhythmic risk.

6.2. Spiral breakup during fibrillation explained as a 1D map

One interesting novel phenomenon which has been established through modeling studies and had important consequence

Fig. 17. Schematic overview of filament evolution in the regimes of positive and negative tension.

for experimental and clinical research was the relation between dynamical instability and cardiac arrhythmias. In modern view, this phenomenon was found when researchers started to make simulations using ionic models of cardiac tissue. Before that simulations of spiral wave dynamics were performed in low dimensional models of the FitzHugh–Nagumo type. The first full size paper describing rotation of spiral wave and spiral breakup in cardiac model was published in 1990 [144]. In this paper the authors used the first ionic model of cardiac cells, Noble 1962 model [145], initiated a spiral wave and studied its dynamics. Unexpectedly they found that instead of stationary rotation or meandering the spiral wave broke down to multiple vortices, also called wavelets, see Fig. 18.

The excitation pattern was similar to the excitation pattern that can be expected from the so-called multiple wavelet hypothesis, which was the main prevailing idea in clinical and experimental electrophysiology. It was explaining fibrillation by the existence of multiple small vortices (wavelets), with the wavelets being generated by the heterogeneity of cardiac tissue. The numerical study in [144] showed that such heterogeneities are not necessary and complex excitation patterns can also occur in fully homogeneous tissue.

The mechanism of the breakup, however, was unknown at that time. Few years later, a similar phenomenon was almost simultaneously reproduced by several groups in low-dimensional models of cardiac tissue [146–148] and at that time the link with dynamical instabilities was made. In [146], the authors showed that the forcing of one-dimensional tissue at a period of a spiral wave causes an oscillatory instability. In [148] a model that captures the main dynamics of the Noble 1962 model was proposed and it was shown that the mechanism of that instability was a period-doubling instability, which in physiology was known as alternans. The phenomenon of alternans has been known since the beginning of the twentieth century [149]. In an informal way, it means that under periodic stimulations the response of cardiac cell/tissue becomes non-constant in a specific way. That is, one can see alterations in the durations of the action potential duration: after an initial transient, all odd beats may have longer action potential duration (APD) and even beats may have shorter APD.

The phenomenon of alternans can be understood from studying the periodic stimulation of a single cardiac cell, say with period T , as shown in Fig. 19.

For large enough T each stimulus generates an action potential and as a result we will get a periodic sequence of action potentials with a period T . If we denote the duration of the action potential by APD, and the resting time between the previous and the next action potential as a diastolic interval (DI) we obviously have the following relation (Fig. 19):

$$APD_n + DI_n = T. \quad (6.1)$$

What determines the APD? It is widely accepted the main factor influencing APD for a given cell is the resting time after the previous excitation. This assumption implies that each new action potential resets the cell to some standard condition (no memory hypothesis). Mathematically this assumption tells us that the $(n + 1)$ th APD is a function of only the n th DI, or

$$APD_{n+1} = f(DI_n). \quad (6.2)$$

The graph of Eq. (6.2) is commonly called the APD restitution curve in electrophysiology. It normally has the following properties. If $DI \rightarrow \infty$, tissue is fully recovered and we obtain a ‘standard’ action potential, i.e. the curve $APD = f(DI)$ will approach some asymptotic value APD_∞ for large DI . For shorter DI one will have incomplete recovery, which results in shorter action potential, such that f is for large DI a monotonously increasing function. Remark also that for some small recovery time the cell will be in the refractory state when the next stimulus comes and no new AP will be initiated. Hence, the curve will end for some DI_{min} . Thus, we expect a typical restitution curve as shown in Fig. 20.

Let us rewrite Eq. (6.1) using (6.2) as a dynamical system:

$$APD_{n+1} = f(DI_n) = f(T - APD_n) = F(APD_n, T). \quad (6.3)$$

Thus, the process shown in Fig. 19 can be viewed as discrete map of a single variable APD with a parameter T .

For each T , we can look for an equilibrium to this map, that corresponds to a periodic regime shown in Fig. 19. The equilibrium will be denoted as APD^* and $DI^* = T - APD^*$ and is given

Fig. 18. Breakup of spiral wave in an ionic model [REF: Noble 62] of cardiac tissue. Reproduced from Panfilov and Holden (1990) [144] with permission. Excitation patterns a–g, h–j shown 1560 ms after initiation of the spiral and are 80 ms apart. There is a gap of 880 ms between frames g and h. Color represents the transmembrane voltage. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

Fig. 19. Three subsequent action potentials, with corresponding action potential duration (APD) and diastolic interval (DI). Horizontal axis denotes time and vertical axis voltage.

Fig. 20. Schematic representation of the APD restitution curve.

by:

$$f(DI^*) + DI^* = T \Leftrightarrow APD^* = F(APD^*, T). \quad (6.4)$$

Here we can apply a standard stability criterion from dynamical systems theory: if $x_{n+1} = f(x_n)$ exhibits a fixpoint x^* , then the linear stability of x^* for small perturbations follows from putting $x_n = x^* + \tilde{x}_n$. This delivers $\tilde{x}_{n+1} = f'(x^*)\tilde{x}_n$, such that the fixpoint is linearly stable if and only if $|f'(x^*)| < 1$.

In relation to the restitution curve, it follows that a stable APD can exist under periodic stimulation only if

$$\left| \frac{\partial F}{\partial APD}(APD^*, T) \right| = \left| -\frac{\partial f}{\partial DI}(DI^*) \right| < 1. \quad (6.5)$$

Since generally f is a monotonically increasing function, one finds $-f'$ is negative and instability can occur only if $-f' < -1$, i.e. if:

$$df(DI^*)/dDI^* > 1 \quad (6.6)$$

at the equilibrium point (APD^*, DI^*) for given period T . For discrete map (6.3) this is a condition for period doubling bifurcation [150]. It is well known that such bifurcation manifests itself as the onset of a period-two orbit, i.e. sequence of points oscillating around the equilibrium with period 2, what in our case means sequence short–long–short–etc action potentials.

In accordance with this analysis, the bifurcation was found to occur if the slope of the APD restitution curve for the given period of stimulation T will be steeper than 1. Because for lower periods of stimulation DI will be smaller and normally the slope of the restitution curve increases with a decrease in DI , the alternans instability normally occurs for short stimulation periods. This simple criterion for the onset of the alternans instability was proposed in [151,152]. In [148] it was shown that this instability causes the breakup of spiral waves in 2D. More recently, extensive studies of this instability in relation to spiral breakup and mechanisms of fibrillation were performed at UCLA and Cornell University [153,154] and the so-called restitution hypothesis of the cardiac fibrillation was formulated. According to this hypothesis, onset of fibrillation is a result of two factors: (1) the restitution curve of cardiac tissue must have a maximal slope steeper than one and (2) the arrhythmia must have a period for which the slope to the restitution curve is steeper than one. If those conditions are met, a complex spatiotemporal pattern of excitation associated with fibrillation will occur.

It was further suggested that to avoid the onset of fibrillation one should make the restitution curve less steep. The idea to flatten restitution curves has been tested in several experimental studies. In [153], the onset of fibrillation in a dog heart was studied. It was found that while in a control group, VF could be induced in all dogs, and if drugs were given which flatten the restitution curve, it became impossible to induce VF, thus conforming the hypothesis. Another research with similar objectives was performed in [154], using a drug called bretylium. This paper showed that this drug decreases the slope of the restitution curve and at the same time prevents onset of ventricular fibrillation. By mapping of electrical activity of the heart during drug application it was shown that bretylium converts a complex spatiotemporal pattern of ventricular fibrillation to a regular tachycardia pattern organized by a single spiral.

Of course the above analytical representation of cardiac dynamics is simplified and a lot of subsequent research has been performed to improve the analysis. In [155], the process of rotation of wave in a ring of excitable tissue was studied. It was shown that such dynamics can be described by a delayed differential equation and in that case the alternans instability manifests

itself as an infinite-dimensional Hopf bifurcation. For this case the condition (6.3) still holds. Later, in [156], it was shown that such degeneracy can be unfolded if one takes into account dependency of the conduction velocity on DI . In that case alternans instability can occur for different values of the slope which also depends on the conduction velocity restitution and may substantially differ from condition (6.3). Recent work has furthermore shown that anti-arrhythmic drug action is achieved by modulation of the steady-state action potential duration, rather than flattening of the restitution slope [157]. Another limitation of the restitution hypothesis (6.6) is that it assumes a periodic pacing. Experimental and numerical studies have shown that the condition is violated under non-periodic stimulation [158,159].

The forementioned studies initiated an extensive research on identification of other types of dynamical instabilities in cardiac tissue. Currently, more than eight types of the alternans instability have been identified [160]. For example, alternans at the tissue level can be spatially concordant or discordant. In mathematical terms, the modes in discordant alternans vary in space, while concordant alternans, the entire tissue alternates together (in phase) between short and long action potential duration. The onset of spatial dependency can be created by conduction velocity restitution [161]. Discordant alternans is considered to be more arrhythmogenic as it induces spatial differences in refractoriness, which can result in the onset of wave blocks and formation of arrhythmia sources. Several types of alternans can occur as a result of dynamical instabilities in intracellular calcium dynamics [162]. For a detailed description of these types of alternans instability and its relation to such factors as action potential duration, conduction velocity, and calcium concentration, see e.g. [163] or the review [160]. In addition, these studies initiated many experimental investigations on the quantification of the restitution properties in cardiac tissue. It became widely accepted that the refractory period in the heart is not constant, but it strongly depends on the history of excitation and it can create a huge dynamical heterogeneity in the heart even in the absence of any anatomical heterogeneity [164,165].

Not all instabilities that have been identified were linked to alternans. For example, short-term variability may exist, i.e. a situation when under periodic stimulation a beat-to-beat variability is observed which is not a period-two process. This type of instability is currently extensively studied in experimental and clinical settings. Some indications show that it may be a predictor of cardiac arrhythmias, sudden cardiac death and can also be used to rank efficacy of anti-arrhythmic drugs [166].

6.3. Dynamics of complex structures unraveled by properties of smaller building blocks

6.3.1. Motivation

Many patterns observed in reaction–diffusion systems are built up from smaller dynamical structures. If we consider any wave pattern in 2D or 3D myocardium then in any small section of the wave orthogonal to the wave front we will get a pattern resembling a plane wave (Fig. 21a), which is typical for 1D propagation. We can thus take this solution of the 1D problem and using it build the whole activation pattern by translating it along the level lines (in 2D) or level surfaces (in 3D) of the pattern. At another level we clearly see that in each section orthogonal to the filament of a 3D rotor we have a 2D spiral wave and, thus, we can build up such 3D solution from 2D spiral waves (Fig. 21b). The formal mathematical method to assemble these solutions along another spatial dimension is ‘gradient expansion’ and it allows us to deduce the behavior of the full pattern from its constituents, see e.g. [139,167–170]. For slightly or moderately curved wave

fronts or scroll wave filaments, this theory is well understood and outlined below.

The motivation of these works is the following: instead of using a computational approach to bridge across spatial scales, semi-analytical results form another, powerful means to demonstrate properties and stability of wave patterns. In contrast to direct numerical integration, analytical results allow us to find generic laws that apply to a broad class of models, e.g. all cardiac monodomain models or all reaction–diffusion systems supporting traveling waves. In such case, it becomes possible to identify emergent parameters that will govern the overall dynamics of the system, irrespective of the detailed ionic model used. In mathematical terms, this comes down to finding bifurcation parameters that control the system. Two examples of these are the slope of the restitution curve (see Eq. (6.6)) and the sign of the filament tension γ_1 (see Eq. (6.32)). Also, a multi-spiral state such as fibrillation can be seen as different spiral waves interacting with each other, and how these building blocks interact with each other or medium boundaries is still a topic of active investigation [171–173].

6.3.2. Response function theory of spiral waves

The response function theory for traveling waves has been explained above in Section 2.3.1, since it applies to both atherosclerosis waves and electrical disturbances.

Response function theory can also be applied to spiral waves, i.e. rotors in the context of cardiac electrophysiology. The essence of this approach is the following: if the spiral wave (rotor) is stable, external stimuli below a critical strength will not cause wave breaks, such that the spiral remains to be present. After some transient time, the spiral will be the same as before, but it could have shifted relative to the case where the same spiral was not subjected to the perturbation. In general, such shift can be expected for every (continuous, Galilean) symmetry of the reaction–diffusion equations that is not present in the spiral wave. That is, circular-core spirals in 2D may shift the position of their tip in the X and Y directions, as well as their orientation, i.e. rotation phase Φ . Meandering spiral waves (see Section 6.1.1) may by external factors be pushed forward and backward in their meander cycle of periodic deformation, such that the meander phase Ψ is the fourth degree of freedom that can be affected [174].

Now, if one hypothetically records the shifts in X, Y, Φ, Ψ induced by a localized stimulus delivered at position $\vec{x}_0$ and during meander phase Ψ_0 to the j th state variable, these data can be gathered into a set of functions $W_j^\mu(\vec{x}_0, \Psi_0)$, where $\mu \in \{X, Y, \Phi, \Psi\}$. In general, the effect of an arbitrary spatiotemporal perturbation $-\mathbf{h}$ then follows from the linear superposition of the elementary responses to localized sources:

$$X^\mu(t) - X_{\text{ref}}^\mu(t) = \int_{-\infty}^t d\tau \int_{\mathbb{R}^2} d^2x_0 \sum_j W_j^\mu(\vec{x}_0, \Psi(\tau)) h_j(\vec{x}_0, \tau),$$

$$\mu \in \{X, Y, \Phi, \Psi\}.$$
(6.7)

In paragraph 6.3.4 we will show in mathematical detail that the $\mathbf{W}^\mu$ are the critical eigenmodes of the adjoint linearized operator associated with a spiral wave solution; see also [175] for the circular-core case and [174] for the meander case. The adjoint property (Eq. (6.19) below) [173,176–178] allows us to accurately evaluate spiral wave response functions; the operational definition (6.7) has also been used to estimate a response function for meander [179] in view of a possible future real-life measurement of response functions in experiments. It turns out that for all “classical” spiral waves, numerically computed

Fig. 21. (a) Reduction of a complex pattern to smaller entities: wave trains, rotors, wave fronts and action potentials. (b) In every section of a scroll wave across its rotation axis or filament (green), one finds a spiral wave. This illustrates that a 3D pattern can be broken down to a stack of 2D solutions. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

response functions are strongly localized in space around the spiral wave tip; see the example in Fig. 22. The only exception found was a counter-rotating spiral [180,181], i.e. with concave instead of convex wave fronts along the spiral arms. In [179], it was furthermore remarked that for a spiral wave with “linear” core, the response function was localized near the turning points of the tip trajectory.

The observation that spiral wave response functions are localized is a key to understanding arrhythmias: since the rotor tip is insensitive to stimuli applied to its periphery, one can only control spiral waves (and, therefore, many types of cardiac arrhythmia) if one manages to act directly on the spiral wave tip. Thus, localization of response functions suggests that control of electrical rotors in the heart using a low-amplitude stimulus should be preferentially targeted to the spiral wave cores.

The spatial localization of spiral wave response functions demonstrates that they behave as localized “particles”, a property sometimes referred to as “particle-wave dualism” of spiral waves [178]. This dualism further supports the “building block” approach to understand pattern formation in the heart that we review in this section.

6.3.3. Spiral wave response functions

Suppose that one has a multi-component reaction–diffusion system

$$\partial_t \mathbf{u} = \hat{\mathbf{P}}(\partial_x^2 + \partial_y^2) \mathbf{u} + \mathbf{F}(\mathbf{u}) - \mathbf{h} \quad (6.8)$$

that exhibits a stable spiral wave solution $\mathbf{u}_0(x, y, t)$ for $\mathbf{h} = \mathbf{0}$:

$$\partial_t \mathbf{u}_0 = \hat{\mathbf{P}}(\partial_x^2 + \partial_y^2) \mathbf{u}_0 + \mathbf{F}(\mathbf{u}_0). \quad (6.9)$$

We can linearize the problem around the solution $\mathbf{u}_0$ by considering a reference frame in which the spiral wave solution becomes stationary (for circular-core spirals) or periodic (for meandering spirals). We restrict our detailed discussion to circular-core spiral waves here. Consider a frame whose origin coincides with the spiral wave tip (having Cartesian coordinates $X^a(t)$), and whose first coordinate axis encloses an angle $\Phi(t)$ with the positive X -axis :

$$x^a = X^a(t) + \sum_{A=1}^2 R_A^a(\Phi(t)) \rho^A, \quad (6.10)$$

where $R_A^a(\Phi)$ is the rotation matrix $\begin{pmatrix} \cos \Phi & -\sin \Phi \\ \sin \Phi & \cos \Phi \end{pmatrix}$ with matrix inverse $R_A^a(\Phi)$. Below, we will use indices $a, b, \dots$ to indicate the lab frame of reference, and $A, B, \dots$ to denote the co-rotating frame. Implicit summation over repeated indices is assumed henceforth. In the lab frame, the state-variable field is denoted by $\mathbf{u}$ and time by t , while in the moving frame, it will be $\mathbf{U}$ and τ , respectively: $\mathbf{u}(x^a, t) = \mathbf{U}(\rho^A, \tau)$.

We choose for the rotating frame $\partial_t \Phi = \omega$, $\partial_t X^a = R_A^a(\Phi(t)) v^A$ with v^A constant, such that the origin of the rotating frame rotates together with the spiral wave on a circle at constant speed $v = \sqrt{(v^1)^2 + (v^2)^2}$. In this frame of reference, it follows from the chain rule that

$$\begin{aligned} \partial_t \mathbf{u} &= \partial_A \mathbf{U} \partial_t \rho^A + \partial_\tau \mathbf{U} \partial_t \tau \\ &= [\partial_\tau R_A^a(x^a - X^a) - R_A^a \partial_\tau X^a] \partial_A \mathbf{U} + \partial_\tau \mathbf{U} \\ &= (\partial_\tau - v^A \partial_A - \omega \partial_\theta) \mathbf{U}, \end{aligned} \quad (6.11)$$

where $\partial_\theta = \epsilon_A^B \rho^A \partial_B$ is the derivative with respect to the polar angle θ in the rotating frame. Here and below, we use the Levi-Civita symbol ϵ_A^B , given by the matrix $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$.

Eqs. (6.8) and (6.9) become in the rotating frame as follows:

$$\partial_\tau \mathbf{U} = \hat{\mathbf{P}} \Delta \mathbf{U} + \omega \partial_\theta \mathbf{U} + v^A \partial_A \mathbf{U} + \mathbf{F}(\mathbf{U}) - \mathbf{h}, \quad (6.12)$$

$$\mathbf{0} = \hat{\mathbf{P}} \Delta \mathbf{U}_0 + \omega \partial_\theta \mathbf{U}_0 + v^A \partial_A \mathbf{U}_0 + \mathbf{F}(\mathbf{U}_0) \quad (6.13)$$

since $\mathbf{U}_0$ is constant in time.

If at $\tau = 0$ the state is $\mathbf{U}_0 + \tilde{\mathbf{U}}$ with $\tilde{\mathbf{U}}$ small, it will evolve according to $\partial_\tau \tilde{\mathbf{U}} = \hat{\mathbf{L}} \tilde{\mathbf{U}}$ (i.e. Eq. (2.12a)), where

$$\hat{\mathbf{L}} = \hat{\mathbf{P}} \Delta + v^A \partial_A + \omega \partial_\theta + \mathbf{F}'(\mathbf{U}_0). \quad (6.14)$$

Our assumption that the spiral wave is stable thus implies that $\hat{\mathbf{L}}$ has no eigenvalues with positive real part.

Next, let us investigate the effect of symmetries. To this aim, differentiate Eq. (6.13) with respect to ρ^1, ρ^2 or θ to find:

$$\hat{\mathbf{L}} \partial_A \mathbf{U}_0 = -\omega \epsilon_A^B \partial_B \mathbf{U}_0, \quad \hat{\mathbf{L}} \partial_\theta \mathbf{U}_0 = v^A \epsilon_A^B \partial_B \mathbf{U}_0. \quad (6.15)$$

By taking appropriate linear combinations in this subspace, one finds that underlying Euclidean symmetry gives the operator $\hat{\mathbf{L}}$ three eigenvalues with vanishing real part:

$$\hat{\mathbf{L}} \mathbf{V}_n = in \omega \mathbf{V}_n, \quad n \in \{-1, 0, 1\} \quad (6.16)$$

$$\mathbf{V}_{\pm 1} = -\frac{1}{2}(\partial_1 \mathbf{U}_0 \pm i \partial_2 \mathbf{U}_0), \quad \mathbf{V}_0 = -\partial_\theta \mathbf{U}_0 + \frac{v^A}{\omega} \epsilon_A^B \partial_B \mathbf{U}_0. \quad (6.17)$$

On the function space of right-hand eigenfunctions $\mathbb{R}^2 \times I \rightarrow \mathbb{C}^m$, it is natural to use the inner product (2.14), for $n = 2$:

$$\langle \mathbf{f} | \mathbf{g} \rangle = \iint_{\mathbb{R}^2} \mathbf{f}^H \mathbf{g} \, d^2 x. \quad (6.18)$$

If the products $\mathbf{f}^H \mathbf{g}$ are tempered at large distance from the origin, the adjoint operator to $\hat{\mathbf{L}}$ can be defined as $\hat{\mathbf{L}}^\dagger = \hat{\mathbf{P}}^H \Delta - v^A \partial_A - \omega \partial_\theta + \mathbf{F}^H(\mathbf{U}_0)$. It is now assumed that $\hat{\mathbf{L}}^\dagger$ has three critical modes $\mathbf{W}^n$ (“response functions”) that satisfy

$$\hat{\mathbf{L}} \mathbf{W}^n = -in \omega \mathbf{W}^n, \quad \langle \mathbf{W}^m | \mathbf{V}_n \rangle = \delta_n^m, \quad m, n \in \{-1, 0, 1\}. \quad (6.19)$$

These response functions in a complex basis can be converted to a real basis again by using the Hermitian transpose (i.e. adjoint)

Fig. 22. Numerical evaluation of critical eigenmodes $\mathbf{V}_\mu$ and adjoint critical eigenmodes (response functions) $\mathbf{W}^\mu$ for a circular-core spiral wave in a reaction–diffusion system with Aliev–Panfilov kinetics [182]. To this aim, linear and adjoint linear problems were solved using DXSpiral software [183,184]. First and second rows show the first (u) and second (v) variable of the system, where u is normalized transmembrane voltage.

matrix:

$$\begin{pmatrix} \mathbf{V}_1 \\ \mathbf{V}_{-1} \\ \mathbf{V}_0 \end{pmatrix} = \begin{pmatrix} -\frac{1}{2} & \frac{i}{2} & 0 \\ -\frac{1}{2} & -\frac{i}{2} & 0 \\ 0 & 0 & -1 \end{pmatrix} \begin{pmatrix} \partial_x \mathbf{U}_0 \\ \partial_y \mathbf{U}_0 \\ \partial_\theta \mathbf{U}_0 \end{pmatrix} \Rightarrow \begin{pmatrix} \mathbf{W}^x \\ \mathbf{W}^y \\ \mathbf{W}^\theta \end{pmatrix} \\ = \begin{pmatrix} -\frac{1}{2} & -\frac{1}{2} & \frac{v^x}{\omega} \\ -\frac{i}{2} & \frac{i}{2} & \frac{v^x}{\omega} \\ 0 & 0 & -1 \end{pmatrix} \begin{pmatrix} \mathbf{W}^1 \\ \mathbf{W}^{-1} \\ \mathbf{W}^0 \end{pmatrix}. \quad (6.20)$$

6.3.4. Response of a spiral wave to small perturbations

In analogy to Section 2.3.1, let us approximate the solution to the perturbed reaction–diffusion problem (6.8) with $\mathbf{h} = \mathcal{O}(\eta)$ as

$$\begin{aligned} \mathbf{u}(x, y, t) &= \mathbf{U}_0(\rho^A) + \tilde{\mathbf{U}}(\rho^A, t) + \mathcal{O}(\eta^2), \\ \partial_t X^a(t) &= R_A^a(\Phi) v^A + \mathcal{O}(\eta), \\ \partial_t \Phi(t) &= \omega + \mathcal{O}(\eta). \end{aligned} \quad (6.21)$$

The definition of $\tilde{\mathbf{U}}$ can be made unambiguous by the gauge condition

$$\langle \mathbf{W}^\mu | \tilde{\mathbf{U}} \rangle = 0, \quad \mu \in \{x, y, \theta\} \quad (6.22)$$

because, if $\tilde{\mathbf{U}}$ possesses a small component along $\partial_\mu \mathbf{U}_0$, the spiral wave's position and orientation can be changed to satisfy (6.22). As Eq. (2.20) is valid for all times, it follows that $\langle \mathbf{W}^\mu | \partial_\tau \tilde{\mathbf{U}} \rangle = 0$. Then, inserting the ansatz (6.21) into (6.8) and retaining only terms up to order η delivers

$$\partial_\tau \tilde{\mathbf{U}} - \hat{\mathbf{L}} \tilde{\mathbf{U}} = (\partial_t \Phi - \omega) \partial_\theta \mathbf{U}_0 + (\partial_\tau X^A - v^A) \partial_A \mathbf{U}_0 - \mathbf{h} \mathcal{O}(\eta). \quad (6.23)$$

Taking the inner product with any of the response function $\mathbf{W}^x, \mathbf{W}^y, \mathbf{W}^\theta$ then yields

$$\begin{aligned} \partial_t X^a(t) &= R_A^a(\Phi(t)) [v^A + \langle \mathbf{W}^A | \mathbf{h} \rangle] + \mathcal{O}(\eta^2), \\ \partial_t \Phi &= \omega + \langle \mathbf{W}^\theta | \mathbf{h} \rangle + \mathcal{O}(\eta^2). \end{aligned} \quad (6.24)$$

The above relations have hitherto mostly been formulated for $v^A = 0$ [175,183,185], in which case the dynamical law (6.24) applies to the rotation center of the spiral instead of its tip. As in [168], we allow zeroth order motion of the tip and use the tip reference frame. [179].

In the cardiac tissue, spiral waves are often seen to meander, i.e. in a frame that moves and co-rotates with the spiral wave tip, the spiral wave solution is not constant but periodic in time with meander period T . In this case, the progression along the meander cycle can be tracked by the temporal or meander phase

Ψ . Defining $\Omega = 2\pi/T$, one has that $d\Psi/dt = \Omega$ and the solution $\mathbf{u}_0(\rho^1, \rho^2, \Psi)$ will depend on Ψ . Besides the temporal or meander phase Ψ , the solution still depends on the spatial or rotation phase Φ . Here, the rotation phase determines the rotation of the frame in which the unperturbed solution is periodic. If β is the angle between two subsequently visited ‘petals’ of the meander flower, then $d\Phi/dt = \omega$ can be taken constant equal to $(\beta + 2n\pi)/T$, ($n \in \mathbb{Z}$). Another choice is to let the instantaneous rotation of the frame coincide with the rotation at the spiral wave tip, leading to $\omega(\Psi)$ that is 2π -periodic which still satisfies $\frac{1}{2\pi} \int_0^{2\pi} \omega_0 d\psi = \beta/T$. Generally speaking, in the case of meander, the solution, its unperturbed motion and response functions will depend periodically on the meander phase Ψ , and a derivation similar to the circular-core case leads to [174]:

$$\begin{aligned} \partial_t X^a(t) &= R_A^a(\Phi(t)) [v^A(\Psi(t)) + \langle \mathbf{W}^A(\Psi) | \mathbf{h} \rangle] + \mathcal{O}(\eta^2), \\ \partial_t \Phi &= \omega(\Psi(t)) + \langle \mathbf{W}^\theta(\Psi(t)) | \mathbf{h} \rangle + \mathcal{O}(\eta^2), \\ \partial_t \Psi &= \Omega + \langle \mathbf{W}^\Psi(\Psi(t)) | \mathbf{h} \rangle + \mathcal{O}(\eta^2). \end{aligned} \quad (6.25)$$

Eqs. (6.24)–(6.25) are the spiral wave counterpart of Eq. (2.24). They are more complex since spiral waves have more degrees of freedom (broken symmetries) as traveling waves, but all of them can be written as

$$\partial_t X^\mu(t) = v^\mu(X^\mu) + \langle \mathbf{W}^\mu | \mathbf{h} \rangle + \mathcal{O}(\eta^2), \quad (6.26)$$

which is essentially a reduction of the infinite-dimensional phase space to the low-dimensional slow manifold of the system. Note that by taking $h_k = \eta \delta_{jk} \delta(\vec{x} - \vec{x}_0) \delta(\psi - \psi_0)$, and integrating over the full history of the system, one recovers the ‘operational’ response function definition, Eq. (6.7).

Eq. (6.24) has been used to analyze spiral wave drift under different perturbations, including parameter gradients [186], surface curvature [132] and 3D effects [168,169,187,188]. The effect of externally imposed electrical fields cannot be directly inferred, as that would require the bidomain model (Eqs. (5.8)). While response function theory concerns only the eigenmodes with vanishing real part, it is also possible to compute unstable eigenvalues and eigenmodes [189], which then offers insights in the mechanism of instability, e.g. for the transition between circular-core and meandering spiral waves [190] or the development of three-dimensional instability [141].

We also want to mention two alternative approaches to describe spiral wave tip dynamics. One is the ‘kinematic method’ [191], which relates the spiral wave tip velocity to the curvature of the wave front at the tip. Under certain conditions, this behavior can be derived from response function theory [192]. A second method is asymptotic matching of excited and quiescent tissue regions; see e.g. [167,193].

6.3.5. Dynamics of filaments of scroll waves

For three-dimensional rotors which rotate around a filament curve, the filament curvature will also affect its motion [168]. Approximating a portion of the filament as a circular arc with radius of curvature r and curvature $k = 1/r$, one finds from Eqs. (2.25) and (6.24) that

$$\partial_t X^a(t) = R_A^a(\Phi, \Psi)[v^A(\Psi) + \langle \mathbf{W}^A(\Psi) | \hat{\mathbf{P}}_{\partial_B \mathbf{U}_0} \rangle \partial_s^2 X^B], \quad (6.27)$$

which is the leading-order law of motion for the spiral wave that is found in every section transverse to the scroll wave filament. In the original result by Keener [168], both the translational and rotational degrees of freedom were derived. If we put $X^m = (X^1, X^2, \Phi)$ and X^M to represent the same quantities in the co-rotating reference frame, and $v^\phi = \omega$, and

$$R_M^m(\Phi) = \begin{pmatrix} R_A^a(\Phi) & 0 \\ 0 & 1 \end{pmatrix}, \quad (6.28)$$

the result (6.27) can be written as

$$\partial_t X^m = R_M^m[v^M + \langle \mathbf{W}^M(\Psi) | \hat{\mathbf{P}}_{\partial_N \mathbf{U}_0} \rangle \partial_s^2 X^N]. \quad (6.29)$$

Thus, we recover Keener's result [168], showing coupling between rotational and translational degrees of freedom in non-equal diffusion systems ($\hat{\mathbf{P}} \neq \mathbf{DI}$).

In the regime of small curvature, one may average over one spiral wave rotation, or even a full rotation meander cycle (if applicable), to find [139,174,179]

$$\partial_t X(\sigma, t) = \gamma_1 k \vec{N} + \gamma_2 k \vec{B} + \mathcal{O}(k^3), \quad (6.30)$$

where σ is a parameter along the filament (e.g. arc length) and $\vec{N}$ and $\vec{B}$ the unit normal and binormal vectors to it. The values

$$\gamma_1 = \frac{1}{2} \sum_{A=1}^2 \langle \mathbf{W}^A | \hat{\mathbf{P}}_{\partial_B \mathbf{U}_0} \rangle, \quad \gamma_2 = \frac{1}{2} \sum_{A,B=1}^2 \epsilon_{AB}^B \langle \mathbf{W}^A | \hat{\mathbf{P}}_{\partial_B \mathbf{U}_0} \rangle \quad (6.31)$$

are known as the scalar and pseudoscalar filament tension coefficients. In case of meander, an additional averaging over Ψ needs to be included. When a spiral's chirality is reversed, γ_2 switches sign but γ_1 remains the same.

The sign of the filament tension determines much of the linear stability of scroll waves, since Biktashev et al. showed that total filament length L evolves as

$$\partial_t L = -\gamma_1 \int k^2 ds. \quad (6.32)$$

Negative filament tension is therefore expected to lead to filament buckling and multiplication and was numerically seen to convert a single rotor state 'tachycardia' to a fibrillation-like state. However, it was later demonstrated that sufficient wall thickness is needed to give room for this instability to develop [141].

6.4. Curved-space viewpoint on anisotropic wave propagation in the heart

6.4.1. Motivation

The electrical excitation pattern and therefore the efficiency of cardiac contraction depends on the local propagation speed of the signaling waves. Therefore, the cardiac modeling in a realistic geometry should take into account the local fiber direction (either rule-based or image-based) and an estimate of the principal conduction velocities along the principal material axes, i.e. fiber direction, sheet direction and sheet normal direction [111].

On the other hand, several (semi)-analytical results have been derived for isotropic reaction-diffusion systems in the context of pattern formation in oscillating chemical reactions. These results

for an isotropic medium include the velocity-curvature relation (2.26) and the dynamical law of motion for scroll wave filaments by Keener [168] and period-averaged version [139] quoted as Eq. (6.30) above. The concept of filament tension as well as the $c(K)$ relation are fundamental and powerful properties, and it would be convenient to generalize them to an anisotropic context. For the $c(K)$ relation, an expression is given in [194], which however lacks the clear physical meaning given by the curved-space viewpoint below.

6.4.2. Operational definition of distance

Let us start with a simple observation. If a portion of the cardiac wall would have parallel fibers along the X-direction with longitudinal and transverse wave propagation velocities c_1 and c_2 , then isotropy could be restored by scaling the variables as

$$\xi = x/c_1, \quad \eta = y/c_2. \quad (6.33)$$

In the new coordinates (ξ, η) , the propagation velocity becomes $c = 1$ and isotropy is restored. In the new coordinates, the Euclidean distance between two points P_1, P_2 with respective coordinates $(\xi_1, \eta_1), (\xi_2, \eta_2)$ will be

$$s_{12} = \sqrt{(\xi_1 - \xi_2)^2 + (\eta_1 - \eta_2)^2} \quad (6.34)$$

and since $c = 1$, s_{12} equals the travel time for a signal between P_1 and P_2 .

A small remark is appropriate here: due to the dependency of wave velocity on wave front curvature, the original velocities c_1 and c_2 are only reached for plane waves. The time difference for a plane wave to reach P_1 and P_2 depends on the direction of propagation: it could happen that P_1 and P_2 lie in the same plane wave front, which makes the time difference between reaching P_1 and P_2 zero. This is hardly a travel time, since no information is conveyed in the process of reaching P_1 and P_2 . To relieve this ambiguity in the orientation of the plane wave, the travel time is defined as the *maximal* time difference for a plane wave reaching P_1 and P_2 , over all orientations.

Can travel time as a measure of distance be used with locally varying anisotropy? The answer is definitely yes, and will be guided by Einstein's famous theory of general relativity. In it, he states the light speed (not incidentally also annotated as c) should be constant, and that the distance between a pair of points needs to be measured by the time of travel of a light pulse between them.

There are also more mundane examples on using travel time as the operational definition of distance. When planning or navigating a route by car, the cost function one usually minimizes is the travel time such that e.g. road works, traffic jams and speed limits are accounted for. (In modern days, a penalty term for carbon emission should be also added). Fig. 23 presents an example. If one navigates from city A to B or C, which one is closer? Although B is closer to A than C, they are effectively at the same travel time of 30 min from A. The map in Fig. 23b has been modified to visually account for the faster time of travel along the highway running north-south. In general, one cannot draw the result on a plane anymore and ends up with an intrinsically curved surface. A similar phenomenon occurs when dealing with local anisotropy of cardiac tissue: with locally rescaled distance, one ends up with an intrinsically curved volume, that can only be unfolded and made isotropic again when going to more than 3 dimensions. Fortunately, there is no need to do this embedding explicitly; rather one can introduce a new notion of distance. For every nearby (i.e. infinitesimally spaced) pair of points, one defines the effective electrophysiological distance as [169,195–197]

$$ds^2 = \sum_{i,j=1}^3 g_{ij}(\vec{x}) dx^i dx^j, \quad (6.35)$$

Fig. 23. Illustration of how rescaling the unit of distance gives rise to an intrinsically curved space. (a) Map showing Euclidean distance. (b) Attempt to redraw the map such that the local distances are scaled according to velocity of travel. In general, the resulting map can no longer be drawn in the plane.

where $\vec{dx}$ is the difference between the Cartesian coordinates of the pair of points and g_{ij} is the metric tensor, which relates local coordinates to distance via Eq. (6.35). The expression (6.35) is well-known in physics and mathematics, as it also occurs, e.g., with polar coordinates (r, ϕ) defined by $x = r \cos \phi$, $y = r \sin \phi$, in which case $ds^2 = dr^2 + r^2 d\phi^2$. For the cardiac context, Wellner et al. proposed that for the metric tensor $\mathbf{g}$ having elements g_{ij} ,

$$\mathbf{g} = D_0 \mathbf{D}^{-1}. \quad (6.36)$$

Essentially, the metric tensor is proportional to the inverse diffusion tensor $\mathbf{S}$ from the monodomain cardiac model, since larger diffusion of transmembrane potential along the line connecting two nearby points brings them effectively closer to each other. We include a constant factor D_0 here, which is the diffusion coefficient for an isotropic medium which makes $c = 1$. Then, the distance ds given by Eq. (6.35) will be equal to the travel time between two points, which can be easily checked along the coordinate axes, given relation (2.11).

The use of the metric tensor (6.36) allows a direct generalization of many concepts from isotropic to anisotropic space by local rescaling; see Section 6.4.4. However, the effect of the space curvature will become noticeable if one considers objects of size that is no longer small relative to the intrinsic curvature of space. For example, one may draw a map of a small country in a plane without much distortion, but for a region of diameter 1000 km, the earth's radius of curvature (6.3710 1/km) can no longer be neglected.

The consequence of working in a curved space is that one needs to adopt mathematical concepts from differential geometry, e.g., parallel transport, geodesics, covariant differentiation. These concepts are well established and can be found in textbooks; a brief overview is contained in Appendix.

6.4.3. Role of extrinsic and intrinsic curvature in the dynamics of excitation patterns

From the basic principles it is possible to distinguish between the roles of extrinsic curvature (embedding in ambient space) versus intrinsic curvature, i.e., deviations of the geometry from $\mathbb{R}^N$ in view of wave propagation.

First, note that since wave fronts are surfaces with constant value of state variables, there is no effective diffusion in them, and

the intrinsic curvature of a wave front does not affect its propagation speed. The quantity which matters there is the extrinsic curvature, since this will determine how many cells in ambient space need to be excited by the wave front.

A similar argument holds for the filament of a scroll wave: there is no diffusion along the filament of a scroll wave in 3D and therefore, the extrinsic curvature of the filament (called k above) will matter.

Conversely, spiral waves rotate within a surface, and if any of the principal radii of curvature is not much larger than the spiral wave's core size, intrinsic curvature effects will come into play. In lowest order, the Gaussian curvature of the surface will affect the rotation frequency of the spiral.

The orientation-averaged three-dimensional intrinsic curvature of the cardiac wall can be calculated from the diffusion tensor using (6.36). In the heart, assuming a constant myofiber rotation of Θ radians over a wall thickness L , one finds that [170, 197]

$$\mathcal{R} = - \left(\frac{\Theta}{L} \right)^2 \frac{(D_1 - D_2)^2}{2D_1 D_2} < 0, \quad (6.37)$$

such that the cardiac wall can be considered a negatively curved space. This implies that initially parallel geodesics diverge, and it was shown that the wave spreading occurs faster than in a positively curved space [198]. Still, we consider it more likely that the myocardial fiber rotation rate evolved due to mechanical stability and efficiency reasons, and the effect on wave spreading is a consequence rather than a driving factor.

6.4.4. Some results on the dynamics of activation patterns in the anisotropic medium

The curved-space viewpoint alone does not presuppose that one is working with a cardiac model, only that the system obeys a reaction-diffusion equation of the type (5.10) with no-flux boundary conditions, in which stable wave fronts or rotating spiral patterns can arise. Under these circumstances, one finds the following results:

1. **Wellner's minimal principle.** In [153], Wellner et al. not only introduced Eq. (6.35), but also postulated, on the basis of numerical simulations, that for rotational anisotropy, the stationary configuration of a scroll wave filament with

endpoints pinned to opposite sides of the cardiac wall is a geodesic of the space with metric (6.35). This was a significant breakthrough, since this result demonstrated that the apparent “compromise between following the myofiber direction or the transmural direction” could be made concrete and assigned a proper geometric interpretation. In hindsight, the minimal principle follows from applying the positive tension property in a curved space. From the local equivalence principle, a filament with positive tension will strive to minimize its length along every section of it. Therefore, it will only stop doing so once it has become the curve of minimal length between fixed endpoints, i.e. a geodesic.

2. **Anisotropic filament dynamics.** By considering a scroll wave as a stack of 2D spiral waves, Verschelde et al. have derived the anisotropic generalization of Eq. (6.30). Given the discussion above, it should come as no surprise that their final result comes down to replacing ordinary derivatives with covariant derivatives in Eq. (6.30) [169]:

$$\partial_t \vec{X} = \gamma_1 \mathcal{D}_s^2 \vec{X} + \gamma_2 \mathcal{D}_s \vec{X} \times \mathcal{D}_s^2 \vec{X} + \mathcal{O}(k^3). \quad (6.38)$$

From this law of motion, it follows in the given order that $\partial_t \vec{X} = 0$ if and only if the geodesic equation (A.5) is satisfied. Therefore, Eq. (6.38) immediately proves Wellner’s minimal principle for rotor filaments.

3. **Anisotropic velocity–curvature relation.** In a similar manner, the velocity–curvature relation (2.26) can be generalized to the following expression [170]:

$$\partial_t \vec{X} = c \vec{N} - \gamma \mathcal{D}^2 \vec{X} + \mathcal{O}(K^2) \quad (6.39)$$

with $\mathcal{D}^2 = \sum_{a=1}^2 \mathcal{D}_a \mathcal{D}_a$ being the covariant Laplacian in the wave front surface; furthermore one has that $\mathcal{D}^2 \vec{X} = \text{Tr}(\mathbf{K}) \vec{N}$ such that formally Eq. (2.26) is still valid, if one measures distances (and velocity) using (6.35) and replaces the ‘mean curvature’ by the trace of the extrinsic curvature tensor in the curved space.

4. **Rotor drift on a curved anisotropic surface.** Spiral waves are known to change their frequency on a spherical surface [199], and to drift on non-uniformly curved surfaces [200,201]. Additionally imposing anisotropic diffusion on the surface only has the effect that its metric g_{ij} will contain the effects of both extrinsic curvature (i.e. the shape of the surface) and intrinsic curvature (i.e. Gaussian curvature of the surface plus anisotropy). By mapping an unperturbed spiral solution $\mathbf{u}_0$ to Riemann normal coordinates on this curved surface, it is possible to derive for circular-core spirals that [132]

$$\partial_t \Phi = \omega + q_0 \mathcal{R}^2, \quad \partial_t \vec{X} = -q_1 \vec{\nabla} \mathcal{R} - q_2 \vec{N} \times \vec{\nabla} \mathcal{R}. \quad (6.40)$$

That is, expecting that $q_0 > 0$, spiral waves will rotate faster on a positively curved surface due to the angular deficit in a curved space. E.g. on a sphere the circumference of a circle with radius a is slightly smaller than $2\pi a$, see Fig. 24.

The second equation governs spiral wave drift, and states that cardiac rotors are stationary in regions of extremal curvature. That is, if $q_1 > 0$, $\mathcal{R}$ acts as a (pseudo)potential and spiral waves will drift to regions with the most negative curvature, i.e. saddle points of the (intrinsically curved) surface. For $q_1 < 0$, spiral a wave will drift towards positively curved zones (e.g. the atrial appendages).

7. Discussion

7.1. Transition processes in physiology

7.1.1. Multiple equilibria in physiological systems

Many physiological systems manifest bistability or multistability. This means that the system can have more than one stable equilibrium in normal or in pathological conditions. Various diseases can “create” a new stable equilibrium of the system, while the original disease-free equilibrium either remains stable or it becomes unstable. The development of the disease is often a gradual process where the transition from the disease-free equilibrium to the disease equilibrium occurs due to a spatial propagation of the transition zone in the tissues concerned by this process.

Several examples of such processes were considered in this review. Atherosclerosis provides a typical example of such spreading disease which develops as a reaction–diffusion wave between the normal and pathological states. We discussed above that the stable atherosclerotic equilibrium can coexist either with a stable or with an unstable disease-free equilibrium. The existence of the disease equilibrium and stability of the disease-free equilibrium are determined by the level of cholesterol (LDL) in blood. Summarizing the properties of this system, we can identify the following cases. If the LDL level is sufficiently low, then the disease-free state is unique and it is globally asymptotically stable. If the LDL level is sufficiently high, then the disease-free state is unstable, the disease state is stable (monostable case), and the disease will necessarily develop. In the intermediate case, both states are stable (bistable case), and the disease will develop if the wave speed is positive and if the initial perturbation is sufficiently strong to overcome some threshold level.

We can speculate that such self-amplifying positive feedback inflammation – immune response – inflammation is specific also for other inflammatory diseases and for the diseases involving inflammation. In particular, for cancer, neurodegenerative diseases, rheumatoid arthritis, and so on. Let us recall that cancer itself is a spreading disease in the sense that tumor growth can be considered as a reaction–diffusion wave where malignant cells replace normal cells. In the simplest case, it can be modeled by the scalar equation (2.6) with the logistic term $F(u) = u(1 - u)$ for the normalized density of tumor cells u . In a more complete model, it can be described by a system of competition of normal and malignant cells [202] similar to the corresponding models in population dynamics.

From the point of view of multistability, thrombosis is different in comparison with atherosclerosis since blood coagulation kinetics contains two stable equilibria in its normal physiological condition. The balance between them can be shifted to either side resulting in bleeding disorders in one case and to thrombosis in the other case. Inflammation, cancer, chemotherapy treatment move this system towards the prothrombotic equilibrium. External perturbation (injury) creates the initial conditions for the transition from one equilibrium to another one.

Multistability is also present in the heart. It can occur at the level of excitation of single cells. In certain conditions after excitation the heart cell will not recover back to the resting state and will remain at the depolarized state for a very long time. As shown by Tran et al. [203], such multistability is associated with a saddle–node bifurcation, changing the number of equilibria from 1 to 3. Interestingly, other bifurcations also occur when the recovery processes in cardiac cells are suppressed. One of them is a supercritical Hopf bifurcation, beyond which point there is no recovery to the steady state. Instead, the transmembrane voltage exhibits several additional small amplitude spikes (early afterdepolarizations) or small-amplitude oscillations [203].

Fig. 24. Surface curvature and angular deficit. (a) in a positively curved space, the circumference C of a circle of radius r is smaller than $2\pi r$, and it can be expected that a spiral wave thus rotates faster than in the plane. (b) On a negatively curved surface (e.g. a saddle or hyperboloid), the converse is true.

Early afterdepolarizations, oscillations and non-recovery after excitation are well known in physiology and are called triggered activity. Triggered activity is considered as one of the fundamentally important mechanisms of cardiac arrhythmias [204] and is actively investigated both in experimental and modeling studies. A detailed description of such regimes is beyond the scope of our paper. For a recent review on this topic please see e.g. [205].

The presence of depolarized regions can also produce a pathological local periodic electrical activity which was recently confirmed in experimental and modeling studies [206]. There are also many other types of multi-stability at the level of the excitation patterns in the heart. For example, it is well known that by setting different boundary conditions, e.g., by applying an additional electrical stimulus, ventricular fibrillation can be induced in an absolutely normal heart [207]. However, induction of fibrillation in normal heart requires a strong perturbation, while in case of the heart disease it is much simpler to achieve and can even occur spontaneously.

Thus, the presence of more than one equilibrium is specific for physiological systems. The normal rest state can be perturbed by a disease or by an external stimulus providing transition to another equilibrium. If the original state remains stable, then the perturbation should be sufficiently strong. If it becomes unstable, then the transition occurs for any small initial perturbation.

7.1.2. Similar models for different processes

It is a common situation in modeling that similar models can be used to describe different processes. For example, although the logistic equation as well as the system of competition of species were originally suggested for some problems in population dynamics, they are now widely used to describe physiological processes. We will discuss here another interesting similarity between blood coagulation and combustion. This similarity is not formal; it is based on the physical mechanisms of these processes where the rate of chemical reaction strongly increases as a function of temperature in combustion waves and as a function of thrombin concentration in coagulation waves.

A classical model in combustion represents a reaction-diffusion system for the temperature T and for the concentration of reactant P :

$$\frac{\partial T}{\partial t} = \kappa \frac{\partial^2 T}{\partial x^2} + K(T)P - \sigma T, \quad (7.41)$$

$$\frac{\partial P}{\partial t} = d \frac{\partial^2 P}{\partial x^2} - K(T)P. \quad (7.42)$$

Here κ is the coefficient of thermal diffusivity, d the coefficient of mass diffusion, $K(T)$ is the temperature dependence of the

reaction rate, and σ is the normalized rate of heat loss. The function $K(T)$ is usually considered in the form of the Arrhenius exponential, $K(T) = k \exp(-E/RT)$, where E is the activation energy of the chemical reaction, R is the universal gas constant, and k is a factor.

If $\kappa = d$ and $\sigma = 0$, then this system can be reduced to a single equation. For small σ , it can be approximated by the single equation

$$\frac{\partial \theta}{\partial t} = d \frac{\partial^2 \theta}{\partial x^2} + e^{Z\theta}(1 - \theta) - \sigma \theta \quad (7.43)$$

for the dimensionless temperature θ . This equation is similar to Eq. (4.11) for the dimensionless thrombin concentration with the difference that the exponential in (7.43) is replaced by the polynomial in (4.11). Zeldovich and Frank-Kamenetskii developed the method of narrow reaction zone for Eq. (7.43) in order to determine the speed of combustion waves [208,209]. This method gives a good approximation of the wave speed for large values of Z for which the exponential is rapidly increasing. Though the polynomial function θ^3 has a weaker growth rate, this method still gives an acceptable approximation of the wave speed of blood coagulation [70].

The system of Eqs. (4.1)–(4.6) of blood coagulation considered above is obtained under some approximations including the assumption that the concentration of prothrombin can be expressed through the concentration of thrombin, as it is done above for the transition from system (7.41), (7.42) to Eq. (7.43). Without this assumption, the reduced model of blood coagulation for thrombin T and prothrombin P corresponds exactly to system (7.41), (7.42) with the polynomial function instead of exponential. Similar to combustion theory where the method of narrow reaction zone is generalized for the system of equations, it can also be used for this model of blood coagulation [76].

Another interesting remark concerns stability of combustion waves [3,209]. Plane waves lose their stability for large values of Z resulting in appearance of cellular flames or of various oscillating regimes. Such effects are not observed for coagulation waves (though we cannot affirm that they do not exist), probably because the dependence of the reaction rate on thrombin concentration is not sufficiently strong.

7.2. Integration of models for cardiovascular diseases

In terms of cardiac modeling, we have reviewed mainly the electrical function of the heart. One of the trends in the field is a multiphysics approach, where in addition to the electrical activation also mechanical contraction and blood flow inside the

heart is studied [210]. Such approach uses not only the reaction–diffusion systems, but also includes approaches from continuum mechanics and fluid dynamics. All equations can currently be coded in the framework of the finite element method and effectively solved on modern high-performance computers. Of course, verification of such multiphysics approach will require years of work and close collaboration between modeling and experimental studies. However, we think that such methodology can be put even further. For example, it is feasible to extend the existing multiphysics approach and also include blood coagulation processes. This problem is of great practical interest, as the main complication of atrial fibrillation (AF) is clot formation in the atria which can result in stroke. Approximately one-third of all strokes are caused by AF and these strokes are often more severe than those occurrences of stroke without AF [2]. Adding blood coagulation processes to the multiphysics modeling framework is straightforward. Indeed, simulation of excitation patterns during atrial fibrillation can be done by using the corresponding equations for atrial cardiac cells and the description of the substrate for AF (e.g., as in [211]). Electrical excitation will produce contraction pattern and deformation of atria, which can be simulated using one of the approaches of [210,212,213]. The deformation of the atrial wall can be linked to blood flow using e.g. a Windkessel model with the elastic myocardial wall [213]. Finally, the simulated flow can be linked to formation of clots using e.g. dissipative particle dynamics or a spatial adsorption equation approach [45, 214]. Of course, this is just a schematic description of steps and its realization requires a lot of practical work to implement and, more importantly, to verify the hierarchy of the models. First results on this novel integrated line of research have been published [215] and due to the extreme importance of the subject and exciting scientific nature of the problem we believe that work in that direction is or will be undertaken soon in many cardiac modeling centers.

Acknowledgments

VV was partially supported by the “RUDN University Program 5-100, Russia”. AVP was partially supported by the “URFU University Program 5-100, Russia”.

Appendix. Elements from differential geometry

A.1. Metric tensor

$$ds^2 = g_{ab} dx^a dx^b. \quad (\text{A.1})$$

The matrix inverse of g_{ab} is denoted g^{ab} , and $g = \det(\bar{g})$. The Einstein summation convention is used throughout, i.e. a summation over pairs of upper and lower indices is implied without writing the summation symbol. In Cartesian coordinates in the 3D cardiac wall, $g^{ab} = D^a b$, such that $\bar{g}$ is positive-definite and the resulting curved space is called a Riemannian manifold (as opposed to the Minkowski-type metrics which appear in the theory of relativity).

A.1.1. Covariant differentiation

When working on manifolds, one should take into account that basis vectors of the coordinate system can also vary from point to point, leading to the notion of ‘covariant differentiation’:

$$\mathcal{D}_a V^b = \partial_a V^b + \Gamma_{ac}^b V^c \mathcal{D}_a T^{bc} = \partial_a T^{bc} + \Gamma_{ad}^b T^{dc} + \Gamma_{ac}^b T^{bd} \quad (\text{A.2})$$

where the connection coefficients $\Gamma_{ac}^b = \bar{e}^b \cdot \mathcal{D}_a \bar{e}_c$ take into account that the basis vectors for the coordinates also vary from point to

point in the curved space or surface. For a space with metric g_{ij} whose matrix inverse is denoted as g^{ij} one has

$$\Gamma_{bc}^a = g^{ad} \Gamma_{d, bc}, \quad \Gamma_{d, bc} = \frac{1}{2} (\partial_b g_{cd} + \partial_c g_{bd} - \partial_d g_{bc}). \quad (\text{A.3})$$

Another convention is that upper indices may be changed to lower indices by matrix product with g or its inverse:

$$V^a = g^{ab} V_b, \quad V_b = g_{ba} V^a. \quad (\text{A.4})$$

This implies that $g^a_b = \delta^a_b$. Raising and lowering of indices commutes with the covariant derivative, since $\mathcal{D}_c g_{ab} = 0$.

A.2. Geodesics

In a curved space or surface, the equivalent of a straight line is a locally straight line or geodesic. These have the defining property that the covariant derivative of the tangent vector along the curve should vanish:

$$\mathcal{D}_s \vec{T} = \mathcal{D}_s^2 \vec{X} = \vec{0}. \quad (\text{A.5})$$

Note that, in Euclidean space, $g_{ij} = \delta_{ij}$ whence $\Gamma_{ac}^b = 0$ such that covariant derivatives reduce to ‘ordinary’ derivatives and Eq. (A.5) is satisfied by any straight line: $\vec{X}(s) = \vec{X}(0) + s\vec{T}$.

A.3. Riemann normal coordinates

Geodesics can be used to construct around a given origin the coordinates which best approximate Euclidean coordinates; these are known as Riemann normal coordinates. They are constructed by starting geodesics from a given point and labeling them as polar or spherical coordinates. After returning back to Cartesian coordinates, one finds that the metric in Riemann normal coordinates ρ^a can be expanded as

$$g_{ab} = \delta_{ab} - \frac{1}{3} \sum_{c,d} (R_{acbd} + R_{adbc}) \rho^c \rho^d + \dots \quad (\text{A.6})$$

Here, R_{abcd} is the Riemann-tensor, given by

$$R^a_{bcd} = \partial_c \Gamma_{bd}^a - \partial_d \Gamma_{bc}^a + \Gamma_{ce}^a \Gamma_{bd}^e - \Gamma_{de}^a \Gamma_{bc}^e \quad (\text{A.7})$$

which measures intrinsic curvature, i.e. the local deviation from Euclidean space.

Due to symmetries in its indices, it has $n^2(n^2 - 1)/12$ independent components in n spatial dimensions. E.g. for a surface, $n = 2$ such that the Riemann tensor $R_{abcd} = \mathcal{R} \epsilon_{ab} \epsilon_{cd}$ where the Riemann scalar curvature $\mathcal{R}$ equals twice the Gaussian curvature in that point of the surface.

A.4. Intrinsic curvature

Note that it is possible to compute the Riemann tensor based on knowledge of the metric tensor g_{ab} , i.e. by measuring mutual distances within the curved space (submanifold) one is considering. Therefore, this type of curvature is called ‘intrinsic’, opposed to extrinsic curvature (discussed below), which relates to the embedding in ambient space.

Since the Riemann tensor is a quite complex object even in three-dimensional space, one often takes the trace of the Riemann curvature tensor to find the Ricci scalar curvature

$$\mathcal{R} = \sum_{a,b,c,d} R_{abcd} g^{ac} g^{bd}. \quad (\text{A.8})$$

which states whether a space is positively or negatively curved or Euclidean ($\mathcal{R} = 0$).

A.5. Extrinsic curvature

The *extrinsic* curvature measures how an object is embedded in the surrounding space. It does so by considering deviations of the normal vector to the curve, surface or hypersurface. Formally, one defines the extrinsic curvature tensor as

$$K^a_b = \vec{e}^a \cdot D_b \vec{N}. \quad (\text{A.9})$$

In 1 dimension, i.e. along a curve with arc length s , tangent $\vec{T}$ and normal $\vec{N}$, $K^1_1 = \vec{T} \cdot \partial_s \vec{N} = -\vec{N} \cdot \partial_s \vec{T} = -k$ and the extrinsic curvature tensor reduces to the curve's curvature k .

For a surface, the eigenvalues of the rank-2 tensor K^a_b are the intrinsic principal radii of the surface, R_1, R_2 . Its trace and determinant are known as the mean and Gaussian curvature, respectively:

$$H = \text{Tr}(\mathbf{K}) = \frac{1}{R_1} + \frac{1}{R_2},$$

$$K_G = \det(\mathbf{K}) = \frac{1}{R_1 R_2}. \quad (\text{A.10})$$

The mean curvature states how a surface is embedded in the ambient space. E.g. a sheet of paper can be rolled to a cylinder of radius r , such that $R_1 = r$, $R_2 = +\infty$ and $H = 1/r$.

However, Gauss showed in his Theorema Egregium that the product of principal radii is an intrinsic property of the surface, not of the embedding. In modern terms, one states that

$$2K_G = \frac{2}{R_1 R_2} = R_{1212}. \quad (\text{A.11})$$

E.g. a sheet of paper has no intrinsic curvature, whence $K_G = 0$. Still, it can be rolled to a cylinder of radius r , in which case $H = 1/r$ but K_G remains zero.

References

- <http://www.who.int/news-room/fact-sheets/detail/the-top-10-causes-of-death>.
- Kathrin Hahne, Gerold Moennig, Alexander Samol, Atrial fibrillation and silent stroke: links, risks, and challenges, in: *Vascular Health and Risk Management*, 2016.
- A. Volpert, Vit. Volpert, V.I. Volpert, Traveling wave solutions of parabolic systems, in: *Translation of Mathematical Monographs*, vol. 140, Amer. Math. Society, Providence, 1994.
- P.C. Fife, J.B. McLeod, The approach to solutions of nonlinear diffusion equations to traveling front solutions, *Arch. Ration. Mech. Anal.* 65 (1977) 335–361.
- P.C. Fife, J.B. McLeod, A phase plane discussion of convergence to travelling fronts for nonlinear diffusion, *Arch. Ration. Mech. Anal.* 75 (1981) 281–314.
- V. Volpert, Asymptotic behavior of solutions of a nonlinear diffusion equation with a source term of general form, *Sib. Math. J.* 30 (1) (1989) 25–36.
- V. Volpert, Convergence to a wave of solutions of a nonlinear diffusion equation with source of general type, *Sib. Math. J.* 30 (2) (1989) 203–210.
- V. Volpert, A. Volpert, Location of spectrum and stability of solutions for monotone parabolic systems, *Adv. Difference Equ.* 2 (5) (1997) 811–830.
- V. Volpert, A. Volpert, Spectrum of elliptic operators and stability of travelling waves, *Asymptot. Anal.* 23 (2000) 111–134.
- M. Marion, V. Volpert, Existence of pulses for a monotone reaction-diffusion system, *Pure Appl. Funct. Anal.* 1 (1) (2016) 97–122.
- M. Marion, V. Volpert, Existence of pulses for the system of competition of species, *J. Dyn. Diff. Equat.* (2017) <http://dx.doi.org/10.1007/s10884-017-9582-6>.
- T. Galochkina, M. Marion, V. Volpert, Initiation of reaction-diffusion waves of blood coagulation, *Physica D* 376 (2018) 160–170.
- H. Berestycki, L. Nirenberg, Travelling fronts in cylinders, *Annales de l'HP. Analyse non linéaire* 9 (5) (1992) 497–572.
- A. Volpert, V. Volpert, Existence of multidimensional travelling waves and systems of waves, *Comm. Partial Differential Equations* 26 (3–4) (2001) 421–459.
- N. Apreutesei, A. Tosenberger, V. Volpert, Existence of reaction-diffusion waves with nonlinear boundary conditions, *Math. Model. Nat. Phenom.* 8 (4) (2013) 2–17.
- N. Apreutesei, V. Volpert, Reaction-diffusion waves with nonlinear boundary conditions, *Nonlinear Heterog. Medium* 8 (1) (2013) 23–35.
- N. Apreutesei, V. Volpert, Travelling waves for reaction-diffusion problems with nonlinear boundary conditions, in: *Application to a Model of Atherosclerosis*, in: *Pure Appl. Funct. Anal.*, vol. 3, (no. 1) 2018, pp. 43–56.
- A.I. Volpert, V.A. Volpert, Applications of the rotation theory of vector fields to the study of wave solutions of parabolic equations, *Trans. Moscow Math. Soc.* 52 (1990) 59–108.
- H. Dierckx, O. Bernus, H. Verschelde, Accurate eikonal-curvature relation for wave fronts in locally anisotropic reaction-diffusion systems, *Phys. Rev. Lett.* 107 (2011) 108101.
- N. El Khatib, S. Genieys, B. Kazmierczak, V. Volpert, Reaction-diffusion model of atherosclerosis development, *J. Math. Biol.* 65 (2) (2012) 349–374.
- R.L. Bratzler, C.K. Colton, K.M. Smith, Endothelium and permeability: theoretical models for transport of low-density lipoproteins in the arterial wall, in: G.W. Manning, M. Daria-Haust (Eds.), *Atherosclerosis*, Plenum, New York, 1977, pp. 943–951.
- D.K. Stangeby, C.R. Ethier, Coupled computational analysis of arterial LDL transport – effects of hypertension, *Comput. Methods Biomech. Biomed. Eng.* 5 (3) (2002) 233–241.
- C.A. Cobbold, J.A. Sherratt, S.R.J. Maxwell, Lipoprotein oxidation and its significance for atherosclerosis: a mathematical approach, *Bull. Math. Biol.* 64 (2002) 65–95.
- A. Friedman, W. Hao, A mathematical model of atherosclerosis and associated risk factors, *Bull. Math. Biol.* (2017) 1–23.
- N. El Khatib, S. Genieys, V. Volpert, Atherosclerosis initiation modeled as an inflammatory process, *MMNP* (2) (2007) 126–141.
- N. El Khatib, S. Genieys, B. Kazmierczak, V. Volpert, Mathematical modelling of atherosclerosis as an inflammatory disease, *Phil. Trans. R. Soc. A* 367 (2009) 4877–4886.
- R.N. Poston, D.R.M. Poston, Typical atherosclerotic plaque morphology produced in silico by an atherogenesis model based on self-perpetuating propagating macrophage recruitment, *Math. Model. Nat. Phenom.* 2 (2) (2007) 142–149.
- Z.Y. Li, S. Howarth, T. Tang, J.H. Gillard, How critical is fibrous cap thickness to carotid plaque stability? A flow plaque interaction model, *Stroke* 37 (2006) 1195–1196.
- Z.Y. Li, S. Howarth, R.A. Trivedi, J.M. U-King-Im, M.J. Graves, A. Brown, L. Wang, J.H. Gillard, Stress analysis of carotid plaque rupture based on in vivo high resolution MRI, *J. Biomech.* 39 (2006) 2611–2622.
- D. Tang, C. Yang, J. Zheng, P.K. Woodard, G.A. Sicard, J.E. Saffitz, Chun Yuan, 3D MRI-based multicomponent FSI models for atherosclerotic plaques, *Ann. Biomed. Eng.* 32 (2004) 947–960.
- T. Ishikawa, L.F.R. Guimaraes, S. Oshima, R. Yamane, Effect of non-Newtonian property of blood on flow through a stenosed tube, *Fluid Dyn. Res.* 22 (1998) 251–264.
- S. Boujena, O. Kafi, N. El Khatib, A 2D mathematical model of blood flow and its interactions in an atherosclerotic artery, *Math. Model. Nat. Phenom.* 6 (2014) 32–54.
- S. Butenas, K.G. Mann, Blood coagulation, *Biochem. (Moscow)* 61 (3) (2001) 3–12.
- T. Orfeo, S. Butenas, K.E. Brummel-Ziedins, K.G. Mann, The tissue factor requirement in blood coagulation, *J. Biol. Chem.* 280 (52) (2005) 42887–42896.
- H.C. Hemker, Thrombin generation, an essential step in haemostasis and thrombosis, *Haemost. Thromb.* 3 (1993) 477–491.
- H.C. Hemker, S. Béguin, Thrombin generation in plasma: its assessment via the endogenous thrombin potential, *Thromb. Haemost.* 74 (1) (1995) 134–138.
- T. Orfeo, K.E. Brummel-Ziedins, M. Gissel, S. Butenas, K.G. Mann, The nature of the stable blood clot procoagulant activities, *J. Biol. Chem.* 283 (15) (2008) 9776–9786.
- D. Gailani, G.J. Broze, Factor XI activation in a revised model of blood coagulation, *Science* 253 (5022) (1991) 909–912.
- M.A. Panteleev, M.V. Ovanesov, D.A. Kireev, A.M. Shibeko, E.I. Sinauridze, N.M. Ananyeva, A.A. Butylin, E.L. Saenko, F.I. Ataullakhanov, Spatial propagation and localization of blood coagulation are regulated by intrinsic and protein c pathways, respectively, *Biophys. J.* 90 (5) (2006) 1489–1500.
- S.P. Jackson, W.S. Nesbitt, E. Westein, Dynamics of platelet thrombus formation, *J. Thromb. Haemostasis* 7 (1) (2009) 17–20.
- B. Furie, B.C. Furie, Thrombus formation in vivo, *J. Clin. Invest.* 115 (2005) 3355–3362.
- T.J. Stalker, E.A. Traxler, J. Wu, K.M. Wannemacher, S.L. Cermignano, R. Voronov, S.L. Diamond, L.F. Brass, Hierarchical organization in the hemostatic response and its relationship to the platelet signaling network, *Blood* 121 (2013) 1875–1885.

- [43] M.J.E. Kuijpers, K. Gilio, S. Reitsma, R. Nergiz-Unal, L. Prinzen, S. Heeneman, E. Lutgens, M. van Zanvoort, B. Neiswandt, M.O. Egbrink, J. Heemskerk, Complementary roles of platelets and coagulation in thrombus formation on plaques acutely ruptured by targeted ultrasound treatment: a novel intravital model, *J. Thromb. Haemost.* 7 (2009) 152–161.
- [44] A.V. Belyaev, J.L. Dunster, J.M. Gibbins, M.A. Panteleev, V. Volpert, Modeling thrombosis in silico: Frontiers, challenges, unresolved problems and milestones, *Phys. Life Rev.* (2018).
- [45] A. Tosenberger, F. Ataullakhanov, N. Bessonov, M. Panteleev, A. Tokarev, V. Volpert, Modelling of thrombus growth in flow with a DPD-PDE method, *J. Theoret. Biol.* 337 (2013) 30–41.
- [46] A. Tosenberger, F. Ataullakhanov, N. Bessonov, M. Panteleev, A. Tokarev, V. Volpert, Modelling of platelet-fibrin clot formation in flow with a DPD-PDE method, *J. Math. Biol.* 72 (3) (2016) 649–681.
- [47] M.F. Hockin, A model for the stoichiometric regulation of blood coagulation, *J. Biol. Chem.* 277 (21) (2002) 18,322–18,333.
- [48] K.C. Jones, K.G. Mann, A model for the tissue factor pathway to thrombin, *J. Biol. Chem.* 269 (37) (1994) 23,367–23,378.
- [49] S. Jordan, E. Chaikof, Simulated surface-induced thrombin generation in a flow field, *Biophys. J.* 101 (2) (2011) 276–286.
- [50] M.A. Khanin, D.V. Rakov, A.E. Kogan, Mathematical model for the blood coagulation prothrombin time test, *Thromb. Res.* 89 (5) (1998) 222–232.
- [51] A.L. Kuharsky, A.L. Fogelson, Surface-mediated control of blood coagulation: the role of binding site densities and platelet deposition, *Biophys. J.* 80 (3) (2001) 1050–1074.
- [52] D. LaCroix, A reduced equation mathematical model for blood coagulation and fibrinolysis in quiescent plasma, *Int. J. Struct. Chang. Solids* 4 (2012) 23–35.
- [53] D. Luan, M. Zai, J.D. Varner, Computationally derived points of fragility of a human cascade are consistent with current therapeutic strategies, *PLoS Comput. Biol.* 3 (7) (2007) e142, <http://dx.doi.org/10.1371/journal.pcbi.0030142>.
- [54] M.A. Panteleev, A.N. Balandina, E.N. Lipets, M.V. Ovanesov, F.I. Ataullakhanov, Task-oriented modular decomposition of biological networks: Trigger mechanism in blood coagulation, *Biophys. J.* 98 (9) (2010) 1751–1761.
- [55] T. Wajima, G.K. Isbister, S.B. Duffull, A comprehensive model for the humoral coagulation network in humans, *Clin. Pharmacol. Ther.* 86 (3) (2009) 290–298.
- [56] G. Willems, T. Lindhout, W. Hermens, H. Hemker, Simulation model for thrombin generation in plasma, *Haemostasis* 21 (1991) 197–207.
- [57] V.I. Zarnitsina, F.I. Ataullakhanov, A.I. Lobanov, O.L. Morozova, Dynamics of spatially nonuniform patterning in the model of blood coagulation, *Chaos* 11 (2001) 57.
- [58] A.I. Lobanov, T.K. Starozhilova, The effect of convective flows on blood coagulation processes, *Pathophysiol. Haemostasis Thromb.* 34 (2–3) (2005) 121–134.
- [59] A. Sequeira, T. Bodnar, Blood coagulation simulations using a viscoelastic model, *Math. Model. Nat. Phenom.* 9 (6) (2014) 34–45.
- [60] R.D. Guy, A.L. Fogelson, J.P. Keener, Fibrin gel formation in a shear flow, *Math. Med. Biol.* (2006) <http://dx.doi.org/10.1093/imammb/dql022>.
- [61] S. Gir, S. Slack, V.T. Turito, A numerical analysis of Factor X activation in the presence of tissue factor-factor VIIa complex in a flow reactor, *Ann. Biomed. Eng.* 24 (1996) 394–399.
- [62] L.M. Haynes, Y.C. Dubief, K.G. Mann, Membrane binding events in the initiation and propagation phases of tissue factor-initiated zymogen activation under flow, *J. Biol. Chem.* 287 (8) (2012) 5225–5234.
- [63] S.W. Jordan, E.L. Chaikof, Simulated surface-induced thrombin generation in a flow field, *Biophys. J.* 101 (2) (2011) 276–286.
- [64] O.S. Rukhlenko, O.A. Dudchenko, K.E. Zlobina, G.T. Guria, Mathematical modeling of intravascular blood coagulation under wall shear stress, *PLoS One* 10 (7) (2015) e0134028.
- [65] V.I. Zarnitsina, A.V. Pokhilko, F.I. Ataullakhanov, A mathematical model for the spatio-temporal dynamics of intrinsic pathway of blood coagulation. I. The model description, *Thromb. Res.* 84 (1996) 225.
- [66] V.I. Zarnitsina, A.V. Pokhilko, F.I. Ataullakhanov, A mathematical model for the spatio-temporal dynamics of intrinsic pathway of blood coagulation. II. Results, *Thromb. Res.* 84 (5) (1996) 333–344.
- [67] F.I. Ataullakhanov, G.T. Guria, V.I. Sarbash, R.I. Volkova, Spatiotemporal dynamics of clotting and pattern formation in human blood, *Biochim. Biophys. Acta* 1425 (1998) 453–468.
- [68] K. Guria, G. Guria, Spatial aspects of blood coagulation: Two decades of research on the self-sustained traveling wave of thrombin, *Thromb. Res.* 135 (3) (2015) 423–433.
- [69] T. Galochkina, H. Ouzzane, A. Bouchnita, V. Volpert, Traveling wave solutions in the mathematical model of blood coagulation, *Appl. Anal.* 96 (16) (2017) 2891–2905.
- [70] T. Galochkina, A. Bouchnita, P. Kurbatova, V. Volpert, Reaction-diffusion waves of blood coagulation, *Math. Biosci.* 288 (2017) 130–139.
- [71] V. Volpert, Elliptic partial differential equations, in: *Reaction-Diffusion Equations*, vol. 2, Birkhäuser, Basel, 2014.
- [72] A. Tokarev, Y. Krasotkina, M. Ovanesov, M. Panteleev, M. Azhigirova, Spatial dynamics of contact-activated fibrin clot formation in vitro and in silico in haemophilia B: Effects of severity and a hemophil B treatment, *Math. Model. Nat. Phenom.* 1 (2006) 124.
- [73] A. Bouchnita, A. Tosenberger, V. Volpert, On the regimes of blood coagulation, *Appl. Math. Lett.* 51 (3) (2016) 74–79.
- [74] A.R. Wufsus, N.E. Macera, K.B. Neeves, The hydraulic permeability of blood clots as a function of fibrin and platelet density, *Biophys. J.* 104 (8) (2013) 1812–1823.
- [75] A. Bouchnita, T. Galochkina, P. Kurbatova, P. Nony, V. Volpert, Conditions of microvessel occlusion for blood coagulation in flow, *Int. J. Numer. Methods Biomed. Eng.* 33 (9) (2017) e2850.
- [76] A. Bouchnita, T. Galochkina, V. Volpert, Influence of antithrombin on the regimes of blood coagulation: insights from the mathematical model, *Acta Biotheor.* 64 (4) (2016) 327–342.
- [77] V. Volpert, Elliptic partial differential equations, in: *Fredholm Theory of Elliptic Problems in Unbounded Domains*, vol. 1, Birkhäuser, Basel, 2011.
- [78] A.T. Winfree, Electrical turbulence in three-dimensional heart muscle, *Science* 266 (1994) 1003–1006.
- [79] T. Ashihara, N.A. Trayanova, Cell and tissue responses to electric shocks, *EP Eur.* 7 (s2) (2005) S155–S165.
- [80] S.F. Sears, J.D. Hauf, K. Kirian, G. Hazelton, J.B. Conti, Posttraumatic stress and the implantable cardioverter-defibrillator patient: what the electrophysiologist needs to know, *Circ. Arrhythm. Electrophysiol.* 4 (2) (2011) 242–250.
- [81] D. Noble, Modeling the heart—from genes to cells to the whole organ, *Science* 295 (2002) 1678–1682.
- [82] Paulus Kirchhof, Stefano Benussi, Dipak Kotecha, Anders Ahlsson, Dan Atar, Barbara Casadei, Manuel Castella, Hans-Christoph Diener, Hein Heidbuchel, Jeroen Hendriks, Gerhard Hindricks, Antonis S. Manolis, Jonas Oldgren, Bogdan Alexandru Popescu, Ulrich Schotten, Bart Van Putte, Panagiotis Vardas, Stefan Agewall, John Camm, Gonzalo Baron Esquivias, Werner Budts, Scipione Carerj, Filip Casselman, Antonio Coca, Raffaele De Caterina, Spiridon Deftereos, Dobromir Dobrev, José M. Ferro, Gerasimos Filippatos, Donna Fitzsimons, Bulent Gorenek, Maxine Guenoun, Stefan H. Hohnloser, Philippe Kolh, Gregory Y.H. Lip, Athanasios Manolis, John McMurray, Piotr Ponikowski, Raphael Rosenhek, Frank Ruschitzka, Irina Savelieva, Sanjay Sharma, Piotr Suwalski, Juan Luis Tamargo, Clare J. Taylor, Isabelle C. Van Gelder, Adriaan A. Voors, Stephan Windecker, Jose Luis Zamorano, Katja Zeppenfeld, ESC guidelines for the management of atrial fibrillation developed in collaboration with eacts, *EP Eur.* 18 (11) (2016) 1609–1678.
- [83] G.R. Mines, On dynamic equilibrium of the heart, *J. Physiol.* 46 (1913) 349–482.
- [84] A.L. Hodgkin, A.F. Huxley, A quantitative description of membrane current and its application to conduction and excitation in nerve, *J. Physiol.* 117 (1952) 500–544.
- [85] Yoram Rudy, Jonathan R. Silva, Computational biology in the study of cardiac ion channels and cell electrophysiology, *Q. Rev. Biophys.* 39 (1) (2006) 57–116.
- [86] Catherine M. Lloyd, James R. Lawson, Peter J. Hunter, Poul F. Nielsen, The cellml model repository, *Bioinformatics* 24 (18) (2008) 2122–2123.
- [87] R. FitzHugh, Impulses and physiological states in theoretical models of nerve membrane, *Biophys. J.* 1 (1961) 445–465.
- [88] John Rinzel, Joseph B. Keller, Traveling wave solutions of a nerve conduction equation, *Biophys. J.* 13 (12) (1973) 1313–1337.
- [89] D. Barkley, A model for fast computer simulation of waves in excitable media, *Physica D* 49 (1991) 61–70.
- [90] R.R. Aliev, A.V. Panfilov, A simple two-variable model of cardiac excitation, *Chaos Solitons Fractals* 7 (1996) 293–301.
- [91] Colleen C. Mitchell, David G. Schaeffer, A two-current model for the dynamics of cardiac membrane, *Bull. Math. Biol.* 65 (5) (2003) 767–793.
- [92] A. Karma, Spiral breakup in model equations of action potential propagation in cardiac tissue, *Phys. Rev. Lett.* 71 (1993) 1103–1106.
- [93] F.H. Fenton, A. Karma, Vortex dynamics in three-dimensional continuous myocardium with fiber rotation: filament instability and fibrillation, *Chaos* 8 (1998) 20–47.
- [94] A. Bueno-Orovio, E.M. Cherry, F.H. Fenton, Minimal model for human ventricular action potentials in tissue, *J. Theoret. Biol.* 253 (2008) 544–560.
- [95] J.D. Bayer, R.C. Blake, G. Plank, N.A. Trayanova, A novel rule-based algorithm for assigning myocardial fiber orientation to computational heart models, *Ann. Biomed. Eng.* 40 (10) (2012) 2243–2254.
- [96] Sergey F. Pravdin, Vitaly I. Berdyshev, Alexander V. Panfilov, Leonid B. Katsnelson, Olga Solovyova, Vladimir S. Markhasin, Mathematical model of the anatomy and fibre orientation field of the left ventricle of the heart, *Biomed. Eng. Online* 12 (1) (2013) 54.

- [97] Sergey F. Pravdin, Hans Dierckx, Leonid B. Katsnelson, Olga Solovyova, Vladimir S. Markhasin, Alexander V. Panfilov, Electrical wave propagation in an anisotropic model of the left ventricle based on analytical description of cardiac architecture, *PLoS One* 9 (5) (2014) e93617.
- [98] I.J. LeGrice, B.H. Smail, L.Z. Chai, S.G. Edgar, J.B. Gavin, P.J. Hunter, Laminar structure of the heart. II. Mathematical model, *Am. J. Physiol.* 269 (1995) H571–H582.
- [99] Bryan J. Caldwell, Mark L. Trew, Gregory B. Sands, Darren A. Hooks, Ian J. LeGrice, Bruce H. Smail, Three distinct directions of intramural activation reveal nonuniform side-to-side electrical coupling of ventricular myocytes, *Circ. Arrhythm. Electrophysiol.* 2 (4) (2009) 433–440.
- [100] Siew Yen Ho, Robert H. Anderson, Damián Sánchez-Quintana, Atrial structure and fibres: morphologic bases of atrial conduction, *Cardiovasc. Res.* 54 (2) (2002) 325–336.
- [101] A.L. Muler, V.S. Markin, *Biofizika* 22 (1977) 307.
- [102] L. Tung, A Bi-Domain Model for Describing Ischemic Myocardial D-C Potentials, (Ph.D. thesis), 1977.
- [103] A. Peskoff, Electric potential in three-dimensional electrically syncytial tissues, *Bull. Math. Biol.* 41 163.
- [104] C.S. Henriquez, Simulating the electrical behavior of cardiac tissue using the bidomain model, *Crit. Revs. Biomed. Eng.* 21 (1993) 1–77.
- [105] R.H. Clayton, O. Bernus, E.M. Cherry, H. Dierckx, F.H. Fenton, L. Mirabella, A.V. Panfilov, F. Sachse, G. Seemann, H. Zhang, Models of cardiac tissue electrophysiology: Progress, challenges and open questions, *Prog. Biophys. Mol. Biol.* 104 (2011) 22–48.
- [106] P. Colli-Franzone, L.F. Pavarion, B. Taccardi, Simulating patterns of excitation, repolarization and action potential duration with cardiac bidomain and monodomain models, *Math. Biosci.* 197 (2005) 35–66.
- [107] M. Potse, B. Dube, J. Richer, A. Vinet, R.M. Gulrajani, A comparison of monodomain and bidomain reaction–diffusion models for action potential propagation in the human heart, *IEEE Trans. Biomed. Eng.* 53 (12) (2006) 2425–2435.
- [108] B.J. Roth, How the anisotropy of the intracellular and extracellular conductivities influences stimulation of cardiac muscle, *J. Math. Biol.* 30 (1992) 633–646.
- [109] P.A. Helm, H.-J. Tseng, L. Younes, E.R. McVeigh, R.L. Winslow, Ex vivo 3D diffusion tensor imaging and quantification of cardiac laminar structure, *Magn. Reson. Med.* 54 (2005) 850–859.
- [110] Myofiber architecture of the human atria as revealed by submillimeter diffusion tensor imaging, *Circ. Arrhythm. Electrophysiol.* 9 (2016) e004133.
- [111] S.H. Gilbert, A.P. Benson, P. Li, A.V. Holden, Regional localisation of left ventricular sheet architecture: integration with current models of cardiac fibre, sheet and band structure, *Eur. J. Cardiothorac. Surg.* 32 (2007) 231–249.
- [112] A.J. Pope, G.B. Sands, B.H. Smail, I.J. LeGrice, Three-dimensional transmural organization of perimysial collagen in the heart, *Am. J. Physiol. Heart Circ. Physiol.* 295 (2008) 1243–1252.
- [113] F. Sachse, A.P. Moreno, G. Seemann, J.A. Abildskov, A model of electrical conduction in cardiac tissue including fibroblasts, *Ann. Biomed. Eng.* 37 (2009) 874.
- [114] D.E. Hurtado, S. Castro, A. Gizzi, Computational modeling of non-linear diffusion in cardiac electrophysiology: A novel porous-medium approach, *Comput. Methods Appl. Mech. Engrg.* 300 (2016) 70–83.
- [115] Alfonso Bueno-Orovio, David Kay, Vicente Grau, Blanca Rodriguez, Kevin Burrage, Fractional diffusion models of cardiac electrical propagation: role of structural heterogeneity in dispersion of repolarization, *J. R. Soc. Interface* 11 (97) (2014) 20140352.
- [116] Simone Rossi, Boyce E. Griffith, Incorporating inductances in tissue-scale models of cardiac electrophysiology, *Chaos* 27 (9) (2017) 093926.
- [117] Peter Kohl, Peter Hunter, Denis Noble, Stretch-induced changes in heart rate and rhythm: clinical observations, experiments and mathematical models, *Prog. Biophys. Mol. Biol.* 71 (1) (1999) 91–138.
- [118] L.D. Weise, A.V. Panfilov, A discrete electromechanical model for human cardiac tissue: effects of stretch-activated currents and stretch conditions on restitution properties and spiral wave dynamics, *PLoS One* 8 (2013) e59317.
- [119] Martyn P. Nash, Peter J. Hunter, Computational mechanics of the heart, *J. Elasticity* 61 (1–3) (2000) 113–141.
- [120] Serdar Göktepe, Ellen Kuhl, Electromechanics of the heart: a unified approach to the strongly coupled excitation–contraction problem, *Comput. Mech.* 45 (2–3) (2010) 227–243.
- [121] Viatcheslav Gurev, Ted Lee, Jason Constantino, Hermenegild Arevalo, Natalia A. Trayanova, Models of cardiac electromechanics based on individual hearts imaging data, *Biomech. Model. Mechanobiol.* 10 (3) (2011) 295–306.
- [122] Thomas Fritz, Christian Wieners, Gunnar Seemann, Henning Steen, Olaf Dössel, Simulation of the contraction of the ventricles in a human heart model including atria and pericardium, *Biomech. Model. Mechanobiol.* 13 (3) (2014) 627.
- [123] Sander Arens, Hans Dierckx, Alexander V. Panfilov, GEMS: A fully integrated PETSc-based solver for coupled cardiac electromechanics and bidomain simulations, *Frontiers Physiol.* 9 (2018).
- [124] V.S. Zykov, *Simulation of Wave Processes in Excitable Media*, Manchester University Press, Manchester, 1987.
- [125] I.R. Efimov, V.I. Krinsky, J. Jalife, Dynamics of rotating vortices in the Beeler–Reuter model of cardiac tissue, *Chaos Solitons Fractals* 5 (1995) 513–526.
- [126] Z. Qu, F. Xie, A. Garfinkel, J.N. Weiss, Origins of spiral wave meander and breakup in a two-dimensional cardiac tissue model, *Ann. Biomed. Eng.* 28 (2000) 755–771.
- [127] R.L. Burton, S.M. Cobbe, Dispersion of ventricular repolarization and refractory period, *Cardiovasc. Res.* 50 (2001) 10–23.
- [128] A.N. Rudenko, A.V. Panfilov, Drift and interaction of vortices in two-dimensional heterogeneous active medium, *Studia Biophys.* 98 (1983) 183–188.
- [129] A.V. Panfilov, B.N. Vasiev, Vortex initiation in a heterogeneous excitable medium, *Physica D* 49 (1991) 107–113.
- [130] K.H.W.J. Ten Tusscher, A.V. Panfilov, Reentry in heterogeneous cardiac tissue described by the Luo–Rudy ventricular action potential model, *Am. J. Physiol. Heart Circ. Physiol.* 284 (2003) H542–H548.
- [131] A.V. Panfilov, J.P. Keener, Re-entry generation in anisotropic twisted myocardium, *J. Cardiovasc. Electrophys.* 4 (1993) 412–421.
- [132] H. Dierckx, E. Brisard, H. Vershelde, A.V. Panfilov, Drift laws for spiral waves on curved anisotropic surfaces, *Phys. Rev. E* 88 (1) (2013) 012908.
- [133] A.Yu. Abramychyev, V. Davydov, V. Zykov, Drift of spiral waves on nonuniformly curved surfaces, *J. Exp. Theor. Phys.* 70 (1990) 666.
- [134] J.A. Abildskov, R.L. Lux, The mechanism of simulated torsade de pointes in a computer model of propagated excitation, *J. Cardiovasc. Electrophysiol.* 2 (1991) 224–237.
- [135] R.A. Gray, J. Jalife, A. Panfilov, W.T. Baxter, C. Cabo, A.M. Pertsov, Non-stationary vortex-like reentrant activity as a mechanism of polymorphic ventricular tachycardia in the isolated rabbit heart, *Circulation* 91 (1995) 2454–2469.
- [136] A. Garfinkel, Z. Qu, Nonlinear dynamics of excitation and propagation in cardiac tissue, in: D.P. Zipes, J. Jalife (Eds.), *Cardiac Electrophysiology. From Cell to Bedside*, third ed., Saunders, Philadelphia, 2000, pp. 315–320.
- [137] A.V. Panfilov, H. Dierckx, *Theory of Cardiac Arrhythmias*, Elsevier, London, 2017, pp. 325–335.
- [138] A.V. Panfilov, A.N. Rudenko, Two regimes of the scroll ring drift in the three dimensional active media, *Physica D* 28 (1987) 215–218.
- [139] V.N. Biktashev, A.V. Holden, H. Zhang, Tension of organizing filaments of scroll waves, *Philos. Trans. R. Soc. Lond. A Math. Phys. Eng. Sci.* 347 (1994) 611–630.
- [140] S. Alonso, F. Sagues, A.S. Mikhailov, Taming winfree turbulence of scroll waves in excitable media, *Science* 299 (2003) 1722–1725.
- [141] H. Dierckx, O. Selsil, H. Vershelde, V.N. Biktashev, Buckling of scroll waves, *Phys. Rev. Lett.* 109 (2012) 174102.
- [142] Hermenegild J. Arevalo, Fijoy Vadakkumpadan, Eliseo Guallar, Alexander Jebb, Peter Malamas, Katherine C. Wu, Natalia A. Trayanova, Arrhythmia risk stratification of patients after myocardial infarction using personalized heart models, *Nature Commun.* 7 (2016) 11437.
- [143] Adityo Prakosa, Hermenegild J. Arevalo, Dongdong Deng, Patrick M. Boyle, Plamen P. Nikolov, Hiroshi Ashikaga, Joshua J.E. Blauer, Elyar Ghafoori, Carolyn J. Park, Robert C. Blake, et al., Personalized virtual-heart technology for guiding the ablation of infarct-related ventricular tachycardia, *Nat. Biomed. Eng.* 2 (10) (2018) 732.
- [144] A.V. Panfilov, A.V. Holden, Self-generation of turbulent vortices in a two-dimensional model of cardiac tissue, *Phys. Lett. A* 151 (1990) 23–26.
- [145] D. Noble, A modification of the Hodgkin–Huxley equation applicable to purkinje fiber action and pacemaker potential, *J. Physiol.* 160 (1962) 317–352.
- [146] A.V. Panfilov, P. Hogeweg, Spiral break-up in a modified FitzHugh–Nagumo model, *Phys. Lett. A* 176 (1993) 295–299.
- [147] M. Baer, M. Eiswirth, Turbulence due to spiral breakup in a continuous excitable medium, *Phys. Rev. E* 48 (1993) R1635–R1637.
- [148] A. Karma, Spiral breakup in model equations of action potential propagation in cardiac tissue, *Phys. Rev. Lett.* 71 (1993) 1103–1106.
- [149] G.R. Mines, On dynamics equilibrium in the heart, *J. Physiol. (Lond)* 46 (1913) 349–382.
- [150] V.I. Arnold, *Geometrical Methods in the Theory of Ordinary Differential Equations*, Springer-Verlag, New York, Heidelberg, Berlin, 1983.
- [151] M.R. Guevara, A. Ward, A. Shrier, L. Glass, Electrical alternans and period doubling bifurcations, *IEEE Comp. Cardiol.* 562 (1984) 167–170.
- [152] J.B. Nolasco, R.W. Dahlen, A graphic method for the study of alternation in cardiac action potentials, *J. Appl. Physiol.* 25 (1968) 191–196.
- [153] M.L. Riccio, M.L. Koller, R.F. Gilmour Jr., Electrical restitution and spatiotemporal organization during ventricular fibrillation, *Circ. Res.* 84 (1999) 955–963.

- [154] A. Garfinkel, Y.H. Kim, O. Voroshilovsky, Z. Qu, J.R. Kil, M.H. Lee, H.S. Karagueuzian, J.N. Weiss, P.S. Chen, Preventing ventricular fibrillation by flattening cardiac restitution, *Proc. Natl. Acad. Sci. USA* 97 (2000) 6061–6066.
- [155] M. Courtemanche, L. Glass, J.P. Keener, Instabilities of a propagating pulse in a ring of excitable media, *Phys. Rev. Lett.* 70 (1993) 2182–2185.
- [156] Eric Cytrynbaum, James P. Keener, Stability conditions for the traveling pulse: Modifying the restitution hypothesis, *Chaos* 12 (3) (2002) 788–799.
- [157] Kyung Chan Park, Michael J. Shattock, James Winter, Kenneth T. MacLeod, Hsiang-Yu Yang, Angela W.C. Lee, Steven Niederer, Restitution slope is principally determined by steady-state action potential duration, *Cardiovasc. Res.* 113 (7) (2017) 817–828.
- [158] E.G. Tolkacheva, D.G. Schaeffer, Daniel J. Gauthier, W. Krassowska, Condition for alternans and stability of the 1:1 response pattern in a memory model of paced cardiac dynamics, *Phys. Rev. E* 67 (2003) 031904.
- [159] S.S. Kalb, H.M. Dobrovolny, E.G. Tolkacheva, S.F. Idriss, W. Krassowska, D.J. Gauthier, The restitution portrait: A new method for investigating rate-dependent restitution, *J. Cardiovasc. Electrophysiol.* 15 (6) (2004) 698–709.
- [160] Alan Garfinkel, Eight (or more) kinds of alternans, *J. Electrocardiol.* 40 (6) (2007) S70–S74.
- [161] Z. Qu, A. Garfinkel, P.S. Chen, J.N. Weiss, Mechanisms of discordant alternans and induction of reentry in simulated cardiac tissue, *Circulation* 102 (2000) 1664–1670.
- [162] M.E. Diaz, S.C. O'Neill, D.A. Eisner, Sarcoplasmic reticulum calcium content fluctuation is the key to cardiac alternans, *Circ. Res.* 94 (2004) 650.
- [163] Elizabeth M. Cherry, Flavio H. Fenton, Suppression of alternans and conduction blocks despite steep apd restitution: electrotonic, memory, and conduction velocity restitution effects, *Am. J. Physiol.-Heart Circ. Physiol.* 286 (6) (2004) H2332–H2341, PMID: 14751863.
- [164] R.H. Keldermann, K.H.W.J. ten Tusscher, M.P. Nash, R. Hren, P. Taggart, A.V. Panfilov, Effect of heterogeneous apd restitution on vf organization in a model of the human ventricles, *Am. J. Physiol.-Heart Circ. Physiol.* 294 (2) (2008) H764–H774, PMID: 18055526.
- [165] R.H. Keldermann, K.H.W.J. ten Tusscher, M.P. Nash, C.P. Bradley, R. Hren, P. Taggart, A.V. Panfilov, A computational study of mother rotor vf in the human ventricles, *Am. J. Physiol.-Heart Circ. Physiol.* 296 (2) (2009) H370–H379, PMID: 19060124.
- [166] Alexandre Bossu, Rosanne Varkevisser, Henriette D.M. Beekman, Marien J.C. Houtman, Marcel A.G. Van Der Heyden, Marc A. Vos, Short-term variability of repolarization is superior to other repolarization parameters in the evaluation of diverse antiarrhythmic interventions in the chronic atrioventricular block dog, *J. Cardiovasc. Pharmacol.* 69 (6) (2017) 398.
- [167] J.P. Keener, A geometrical theory for spiral waves in excitable media, *SIAM J. Appl. Math.* 46 (1986) 1039–1056.
- [168] J.P. Keener, The dynamics of three-dimensional scroll waves in excitable media, *Physica D* 31 (1988) 269–276.
- [169] H. Verschelde, H. Dierckx, O. Bernus, Covariant stringlike dynamics of scroll wave filaments in anisotropic cardiac tissue, *Phys. Rev. Lett.* 99 (2007) 168104.
- [170] H. Dierckx, Dynamics of Wave Fronts and Filaments in Anisotropic Cardiac Tissue (Ph.D. thesis), Ghent University, 2010.
- [171] R.M. Zariwsky, A.M. Pertsov, Stable spiral structures and their interaction in two-dimensional excitable media, *Phys. Rev. E* 66 (2002) 066120.
- [172] Christian W. Zemlin, Karthik Mukund, Vadim N. Biktashev, Arkady M. Pertsov, Dynamics of bound states of same-chirality spiral waves, *Phys. Rev. E* 74 (1, 2) (2006).
- [173] C.D. Marcotte, R.O. Grigoriev, Unstable spiral waves and local Euclidean symmetry in a model of cardiac tissue, *Chaos* 25 (6) (2015) 063116.
- [174] H. Dierckx, H. Verschelde, A.V. Panfilov, I.V. Biktasheva, V.N. Biktashev, Filament tension and phase-locking of meandering scroll waves, *Phys. Rev. Lett.* 119 (2017) 258101, 1–6.
- [175] V.N. Biktashev, A.V. Holden, Resonant drift of autowave vortices in two dimensions and the effect of boundaries and inhomogeneities, *Chaos Solitons Fractals* 5 (1995) 575–622.
- [176] D. Barkley, Euclidean symmetry and the dynamics of rotating spiral waves, *Phys. Rev. Lett.* 72 (1994) 164–167.
- [177] V. Hakim, H. Henry, Scroll waves in isotropic excitable media: linear instabilities, bifurcations and restabilized states, *Phys. Rev. E* 65 (2002) 046235.
- [178] I.V. Biktasheva, V.N. Biktashev, Wave-particle dualism of spiral wave dynamics, *Phys. Rev. E* 67 (2003) 026221.
- [179] H. Dierckx, H. Verschelde, A.V. Panfilov, V.N. Biktashev, I.V. Biktasheva, A response function framework for the dynamics of meandering or large-core spiral waves and modulated traveling waves, *Phys. Rev. E* 99 (2019) 022217, 1–20.
- [180] O.A. Mornev, I.M. Tsyganov, O.V. Aslanidi, M.A. Tsyganov, Beyond the Kuramoto-Zel'dovich theory: steadily rotating concave spiral waves and their relation to the echo phenomenon, *JETP Lett.* 77 (2003) 270.
- [181] V.N. Biktashev,
- [182] R.R. Aliev, A.V. Panfilov, A simple two-variable model of cardiac excitation, *Chaos Solitons Fractals* 7 (1996) 293–301.
- [183] I.V. Biktasheva, D. Barkley, V.N. Biktashev, G.V. Borydugov, A.J. Foulkes, Computation of the response functions of spiral waves in active media, *Phys. Rev. E* 79 (2009) 056702.
- [184] D. Barkley, V.N. Biktashev, I.V. Biktasheva, G.V. Borydugov, A.J. Foulkes, DXspiral: A Code for Studying Spiral Waves on a Disk, 2010, <http://cgi.csc.liv.ac.uk/~ivb/SOFTWARE/DXspiral.html>.
- [185] H. Henry, Spiral wave drift in an electrical field and scroll wave instabilities, *Phys. Rev. E* 70 (2004) 026204.
- [186] H. Dierckx, O. Bernus, H. Verschelde, A geometric theory for scroll wave filaments in anisotropic excitable tissue, *Physica D* 238 (2009) 941–950.
- [187] I.V. Biktasheva, H. Dierckx, V.N. Biktashev, Drift of scroll waves in thin layers caused by thickness features: asymptotic theory and numerical simulations, *Phys. Rev. Lett.* 114 (2015) 068302.
- [188] H. Dierckx, H. Verschelde, Effective dynamics of twisted and curved scroll waves using virtual filaments, *Phys. Rev. E* 88 (2013) 062907.
- [189] D. Barkley, Linear stability analysis of rotating spiral waves in excitable media, *Phys. Rev. Lett.* 68 (1992) 2090–2093.
- [190] N.F. Otani, A primary mechanism for spiral wave meandering, *Chaos* 12 (3) (2002) 829–842.
- [191] A.S. Mikhailov, V.A. Davydov, V.S. Zykov, Complex dynamics of spiral waves and motion of curves, *Physica D* 70 (1994) 1–39.
- [192] V. Hakim, A. Karma, Theory of spiral wave dynamics in weakly excitable media: asymptotic reduction to a kinematic model and applications, *Phys. Rev. E* 60 (1999) 5073–5105.
- [193] V. Hakim, A. Karma, Spiral wave meander in excitable media: the large core limit, *Phys. Rev. Lett.* 79 (4) (1997) 665–668.
- [194] J.P. Keener, The effects of discrete gap junction coupling on propagation in myocardium, *J. Theoret. Biol.* 148 (1991) 49–82.
- [195] M. Wellner, O.M. Berenfeld, J. Jalife, A.M. Pertsov, Minimal principle for rotor filaments, *Proc. Natl. Acad. Sci. USA* 99 (2002) 8015–8018.
- [196] K.H.W. ten Tusscher, A.V. Panfilov, Eikonal formulation of the minimal principle for scroll wave filaments, *Phys. Rev. Lett.* 93 (2004) 108106.
- [197] R.J. Young, A.V. Panfilov, Anisotropy of wave propagation in the heart can be modeled by a Riemannian electrophysiological metric, *Proc. Natl. Acad. Sci. USA* 107 (2010) 15063–15068.
- [198] Tristan Aumentado-Armstrong, Amir Kadivar, Peter Savadjiev, Steven Zucker, Kaleem Siddiqi, Conduction in the heart wall: Helicoidal fibers minimize diffusion bias, *Sci. Rep.* 8 (2018) 12.
- [199] V.S. Zykov, S.C. Müller, Spiral waves on circular and spherical domains of excitable medium, *Physica D* 97 (1996) 322–332.
- [200] V.A. Davydov, V.S. Zykov, Kinematics of spiral waves on nonuniformly curved surfaces, *Physica D* 49 (1991) 71–74.
- [201] V.A. Davydov, V.S. Zykov, T. Yamaguchi, Drift of spiral waves on nonuniformly curved surfaces, *Macromol. Symp.* 160 (2000) 99–106.
- [202] N. Bessonov, F. Crauste, I. Demin, V. Volpert, Dynamics of erythroid progenitors and erythroleukemia, *Math. Model. Nat. Phenom.* 4 (3) (2009) 210–232.
- [203] Diana X. Tran, Daisuke Sato, Arik Yochelis, James N. Weiss, Alan Garfinkel, Zhilin Qu, Bifurcation and chaos in a model of cardiac early afterdepolarizations, *Phys. Rev. Lett.* 102 (25) (2009).
- [204] Charles Antzelevitch, Alexander Burashnikov, Overview of basic mechanisms of cardiac arrhythmia, *Cardiac. Electrophysiol. Clin.* 3 (1) (2011) 23–45.
- [205] Nele Vandersickel, Enid Van Nieuwenhuysse, Gunnar Seemann, Alexander V. Panfilov, Spatial patterns of excitation at tissue and whole organ level due to early afterdepolarizations, *Front. Physiol.* 8 (2017).
- [206] A. Teplinin, H. Dierckx, A.S. de Vries, D. Pijnappels, A.V. Panfilov, Paradoxical onset of arrhythmic waves from depolarized areas in cardiac tissue due to curvature-dependent instability, *Phys. Rev. X* 8 (2018) 021077, 1–14.
- [207] P.S. Chen, P. Wolf, E.G. Dixon, N.D. Daniele, D.W. Frazier, W.M. Smith, R.E. Ideker, Mechanism of ventricular vulnerability to single premature stimuli in open-chest dogs, *Circ. Res.* 62 (1988) 1191–1209.
- [208] Y.B. Zeldovich, D.A. Frank-Kamenetskii, A theory of thermal propagation of flame, *Acta Physicochim. USSR* 9 (1938) 341–350.
- [209] Y.B. Zeldovich, G.I. Bareblatt, V.B. Librovich, G.M. Makhviladze, *The Mathematical Theory of Combustion and Explosion*, Plenum, New York, 1985.
- [210] D.A. Nordsletten, S.A. Niederer, M.P. Nash, P.J. Hunter, N.P. Smith, Coupling multi-physics models to cardiac mechanics, *Prog. Biophys. Mol. Biol.* 104 (1–3) (2011) 77–88.
- [211] Michel Haissaguerre, Ashok J. Shah, Hubert Cochet, Meleze Hocini, Remi Dubois, Igor Efimov, Edward Vigmond, Olivier Bernus, Natalia Trayanova, Intermittent drivers anchoring to structural heterogeneities as a major pathophysiological mechanism of human persistent atrial fibrillation, *J. Phys.* 594 (9) (2016) 2387–2398.

- [212] Rikkert H. Keldermann, Martyn P. Nash, Hanneke Gelderblom, Vicky Y. Wang, Alexander V. Panfilov, Electromechanical wavebreak in a model of the human left ventricle, *Am. J. Physiol.-Heart Circ. Physiol.* 299 (1) (2010) H134–H143.
- [213] Julia Hoermann, Cristóbal. Bertoglio, Andreas Nagler, Martin R. Pfaller, Felix Bourier, M. Hadamitzky, Isabel Deisenhofer, Wolfgang A. Wall, Multiphysics modeling of the atrial systole under standard ablation strategies, *Cardiovascular Eng. Technol.* 8 (2) (2017) 205–218.
- [214] Evgenia S. Babushkina, Nikolay M. Bessonov, Fazoil I. Ataullakhanov, Mikhail A. Panteleev, Continuous modeling of arterial platelet thrombus formation using a spatial adsorption equation, *PLoS One* 10 (10) (2015) e0141068.
- [215] Alessandro Masci, Martino Alessandrini, Davide Forti, Filippo Menghini, Luca Dedé, Corrado Tommasi, Alfio Quarteroni, Cristiana Corsi, A patient-specific computational fluid dynamics model of the left atrium in atrial fibrillation: Development and initial evaluation, in: Mihaela Pop, Graham A. Wright (Eds.), *Functional Imaging and Modelling of the Heart*, Springer International Publishing, Cham, 2017, pp. 392–400.