**DR-10. DETECTION OF NITROAROMATIC EXPLOSIVES
BY 2-AMINO-3-ETHOXYCARBONYL-6-(1-METHYLINDOL-3-YL)-5-
(4-CHLOROPHENYL)-PYRAZINE AND ITS DERIVATIVES**

O. S. Taniya¹, L. K. Sadieva¹, I. S. Kovalev¹, G. V. Zyryanov^{1,2},
D. S. Kopchuk^{1,2}, V. L. Rusinov^{1,2}, O. N. Chupakhin^{1,2}

¹Ural Federal University of the first President of Russia B. N. Yeltsin,
Mira St., 19, Yekaterinburg, 620002, Russia

²I. Ya. Postovsky Institute of Organic Synthesis UB RAS,
S. Kovalevskoy/Akademicheskaya St., 20/22, Yekaterinburg, 620990, Russia

E-mail: olgatanya@yandex.ru

Indolyl derivatives of aminopyrazine are widely distributed in nature as compounds involved in the processes of biochemiluminescence. Their structure is part of luciferin – one of the most famous bioluminescent systems. We have previously synthesized isosteric analogues of etioluciferin cypridine. In this article we wish to report our studies on using 2-amino-3-ethoxycarbonyl-6-(1-methylindol-3-yl)-5-(4-chlorophenyl)-pyrazines for the detection of nitroaromatic compounds.

This work was supported by the Russian Science Foundation (Ref. № 18-73-10119).