

**DR-31. PREPARATION OF MONOETHANOLAMINE
AND 5-PHENYL-2,2'-BIPYRIDINE DERIVATIVES
AND THEIR SUBSEQUENT TOSYLATION REACTIONS**


D. S. Kopchuk^{1,2}, I. S. Kovalev¹, S. Santra¹, G. V. Zyryanov^{1,2}, V. L. Rusinov^{1,2},
O. N. Chupakhin^{1,2}, V. N. Charushin^{1,2}

¹Ural Federal University of the first President of Russia B. N. Yeltsin,
Mira St., 19, Yekaterinburg, 620002, Russia

²I. Ya. Postovsky Institute of Organic Synthesis UB RAS,
S. Kovalevskoy / Akademicheskaya St., 20/22, Yekaterinburg, 620990, Russia.

E-mail: dkopchuk@mail.ru

Oxazolidine derivatives are of interest as biologically active compounds, as well as catalysts for enantioselective reactions. Separately, in this aspect, it should be noted that a fragment of N-tosyloxazolidine is presented in a number of biologically active structures. To form the oxazolidine cycle, we used a reaction of monoethanolamine-based Schiff base with tosyl chloride. To date, there are very few examples of such syntheses. In this publication, we propose an approach to N-tosyloxazolidine derivatives having a 2,2'-bipyridine fragment.



This work was supported by the Russian Science Foundation (Ref. № 18-73-00301).