

## DR-29. PREPARATION OF INDOLE-CONTAINING 3-(2-PYRIDYL)-1,2,4-TRIAZINES AS TRYPTAMINE DERIVATIVES


M. I. Savchuk<sup>1</sup>, D. S. Kopchuk<sup>1,2</sup>, A. P. Krinochkin<sup>1,2</sup>, I. S. Kovalev<sup>1</sup>,  
G. V. Zyryanov<sup>1,2</sup>, V. L. Rusinov<sup>1,2</sup>, O. N. Chupakhin<sup>1,2</sup>

<sup>1</sup>Ural Federal University of the first President of Russia B. N. Yeltsin,  
Mira St., 19, Yekaterinburg, 620002, Russia

<sup>2</sup>I. Ya. Postovsky Institute of Organic Synthesis UB RAS,  
S. Kovalevskoy/Akademicheskaya St., 20/22, Yekaterinburg, 620990, Russia

E-mail: savchuk.chuk@yandex.ru

Indole derivatives, including 1,2,4-triazine containing ones, are of interest from the point of view of their biological activity. In this publication, we propose a method for obtaining a Schiff base, comprising indole and 3-(2-pyridyl)-1,2,4-triazine fragments.



*This work was supported by the Russian Science Foundation (Ref. № 18-73-10119).*