


ISSN 2072-5981
doi: 10.26907/mrsej




***magnetic
Resonance
in Solids***

Electronic Journal

*Volume 21
Special Issue 4
Paper No 19409
1-10 pages
2019*

doi: 10.26907/mrsej-19409

<http://mrsej.kpfu.ru>
<http://mrsej.ksu.ru>



Established and published by Kazan University
Endorsed by International Society of Magnetic Resonance (ISMAR)
Registered by Russian Federation Committee on Press (#015140),
August 2, 1996
First Issue appeared on July 25, 1997

© Kazan Federal University (KFU)*

"Magnetic Resonance in Solids. Electronic Journal" (MRSej) is a peer-reviewed, all electronic journal, publishing articles which meet the highest standards of scientific quality in the field of basic research of a magnetic resonance in solids and related phenomena.

Indexed and abstracted by
Web of Science (ESCI, Clarivate Analytics, from 2015), Scopus (Elsevier, from 2012), RusIndexSC (eLibrary, from 2006), Google Scholar, DOAJ, ROAD, CyberLeninka (from 2006), SCImago Journal & Country Rank, etc.

Editor-in-Chief

Boris **Kochelaev** (KFU, Kazan)

Honorary Editors

Jean **Jeener** (Universite Libre de Bruxelles, Brussels)

Raymond **Orbach** (University of California, Riverside)

Executive Editor

Yurii **Proshin** (KFU, Kazan)

mrsej@kpfu.ru



This work is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-sa/4.0/).



This is an open access journal which means that all content is freely available without charge to the user or his/her institution. This is in accordance with the [BOAI definition of open access](https://www.boai.ru/).

Special Editor of Issue

Eduard **Baibekov** (KFU)

Editors

Vadim **Atsarkin** (Institute of Radio Engineering and Electronics, Moscow)

Yurij **Bunkov** (CNRS, Grenoble)

Mikhail **Eremin** (KFU, Kazan)

David **Fushman** (University of Maryland, College Park)

Hugo **Keller** (University of Zürich, Zürich)

Yoshio **Kitaoka** (Osaka University, Osaka)

Boris **Malkin** (KFU, Kazan)

Alexander **Shengelaya** (Tbilisi State University, Tbilisi)

Jörg **Sichelschmidt** (Max Planck Institute for Chemical Physics of Solids, Dresden)

Haruhiko **Suzuki** (Kanazawa University, Kanazawa)

Murat **Tagirov** (KFU, Kazan)

Dmitrii **Tayurskii** (KFU, Kazan)

Valentine **Zhikharev** (KNRTU, Kazan)

Technical Editors of Issue

Nurbulat **Abishev** (KFU)

Maxim **Avdeev** (KFU)

Eduard **Baibekov** (KFU)

Alexander **Kutuzov** (KFU)

* In Kazan University the Electron Paramagnetic Resonance (EPR) was discovered by Zavoisky E.K. in 1944.

The Löwdin orthogonalization and magnetoelectric coupling for noncentrosymmetric ions

A.S. Moskvina

Ural Federal University, Lenin str. 51, Ekaterinburg 620083, Russia

E-mail: alexander.moskvina@urfu.ru

(Received May 9, 2019; accepted May 28, 2019; published June 6, 2019)

The Löwdin orthogonalization procedure being the well-known technique, particularly in quantum chemistry, however, gives rise to novel effects missed in earlier studies. Making use of the technique of irreducible tensorial operators, we have developed a regular procedure for the account of the orthogonalization effects. For illustration, we address the emergence of a specific magnetoelectric coupling for noncentrosymmetric $3d$ or $4f$ ions.

PACS: 31.10.+z, 75.85.+t.

Keywords: Löwdin orthogonalization, irreducible tensorial operators, magnetoelectric coupling.

Dedicated to Boris Malkin, on the occasion of his 80th birthday

1. Introduction

More than 50 years ago P.-O. Löwdin [1] suggested a regular procedure for the orthogonalization of the atomic functions localized at different sites. The orthogonalization problem was in the focus of the electron structure calculations in 1960-70ies, though later on it was undeservedly forgotten in the shade of the so-called “*ab-initio*” calculations. However, many interesting points have been missed being overboard the orthogonalization procedure. In this paper, we generalize the Löwdin technique for many-electron atoms and demonstrate that the irreducible tensorial operator method, which is well-known in theory of free atoms, can be successfully applied to “orthogonalized” electrons. As an urgent issue, we address the orthogonalization contribution to the orbital magnetoelectric coupling.

The paper is organized as follows. In Sec. II we demonstrate the conservation of the effective rotation symmetry under the Löwdin orthogonalization technique. In Sec. III and IV we address some novel properties of different operators acting on the orthogonalized basis set. In Sec. V and VI we calculate the electric dipole moment of the noncentrosymmetric quasi-atoms and the orthogonalization contribution to the orbital magnetoelectric coupling.

2. Löwdin orthogonalization procedure

Making use of the one-electron $\varphi_{fnlm}(\mathbf{r})$ wave functions of a free atom as a basis set for description of the electronic structure in crystals is restricted because of their nonorthogonality for different sites/atoms. One of the more practicable techniques for solution of the problem was suggested by Löwdin [1]. Let us start with a set of standard one-electron functions $\varphi_{\kappa}(\mathbf{r}) = \varphi_{fnlm}(\mathbf{r})$, where f labels a site, and introduce a Hermitian overlap matrix $\hat{\Delta} = \hat{\mathbf{I}} + \hat{\mathbf{S}}$ as follows:

$$\hat{\Delta}_{\kappa\kappa'} = \langle \varphi_{\kappa}(\mathbf{r}) | \varphi_{\kappa'}(\mathbf{r}) \rangle = \delta_{\kappa\kappa'} + S_{\kappa\kappa'}. \quad (1)$$

Linear transformation $\varphi \rightarrow \psi = \varphi \hat{\mathbf{A}}$

$$\psi_{\kappa}(\mathbf{r}) = \sum_{\kappa'} \varphi_{\kappa'}(\mathbf{r}) A_{\kappa'\kappa}, \quad (2)$$

yields a set of orthogonalized functions $\psi_{\kappa}(\mathbf{r})$:

$$\langle \psi_{\kappa}(\mathbf{r}) | \psi_{\kappa'}(\mathbf{r}) \rangle = \delta_{\kappa\kappa'}.$$

It is easy to see that matrix elements $\hat{\mathbf{A}}$ obey a matrix equation

$$\hat{\mathbf{A}}^\dagger \hat{\Delta} \hat{\mathbf{A}} = \hat{\mathbf{1}}, \quad (3)$$

whose solution can be represented as follows

$$\hat{\mathbf{A}} = \hat{\Delta}^{-1/2} \hat{\mathbf{B}}, \quad (4)$$

where $\hat{\mathbf{B}}^\dagger \hat{\mathbf{B}} = \hat{\mathbf{1}}$, given the existence of the matrix $\hat{\Delta}^{-1/2} = (\hat{\mathbf{1}} + \hat{\mathbf{S}})^{-1/2}$. Obviously, the choice of the unitary $\hat{\mathbf{B}}$ matrix, hence the certain orthogonalization procedure, is an ambiguous one. The most physically clear and practicable procedure of a so-called symmetric orthogonalization, in the case when $\hat{\mathbf{B}} = \hat{\mathbf{1}}$ and $\hat{\mathbf{A}} = \hat{\Delta}^{-1/2} = (\hat{\mathbf{1}} + \hat{\mathbf{S}})^{-1/2}$, was suggested by Löwdin [1]. Given a small overlap, when $\sum_{\kappa'} |S_{\kappa'\kappa}| \leq q < 1$ for all κ , the series

$$(\hat{\mathbf{1}} + \hat{\mathbf{S}})^{-1/2} = \hat{\mathbf{1}} - \frac{1}{2} \hat{\mathbf{S}} + \frac{3}{8} \hat{\mathbf{S}}^2 + \dots \quad (5)$$

will converge, and that enables to represent the orthogonalized functions ψ_κ as follows

$$\psi_\kappa(\mathbf{r}) = \sum_{\kappa'} \left((1 + \hat{S})^{-1/2} \right)_{\kappa\kappa'} \varphi_{\kappa'}(\mathbf{r}),$$

or

$$\begin{aligned} \psi_{fnlm}(\mathbf{r}) = & \varphi_{fnlm}(\mathbf{r}) - \frac{1}{2} \sum_{f'n'l'm'} S_{f'n'l'm';fnlm} \varphi_{f'n'l'm'}(\mathbf{r} - \mathbf{R}_{ff'}) + \\ & + \frac{3}{8} \sum_{f'n'l'm'} \sum_{f''n''l''m''} S_{f'n'l'm';f''n''l''m''} S_{f''n''l''m'';fnlm} \varphi_{f'n'l'm'}(\mathbf{r} - \mathbf{R}_{ff'}) + \dots \end{aligned} \quad (6)$$

Hereafter the orthogonalized functions ψ_{fnlm} can be termed as “quasi-atomic”, at variance with nonorthogonalized atomic functions φ_{fnlm} .

The two-site overlap integrals obey an analog of the Wigner-Eckart theorem [2]:

$$\begin{aligned} \int \phi_{l_1 m_1}^*(\mathbf{r}) \phi_{l_2 m_2}(\mathbf{r} - \mathbf{R}_{12}) d\mathbf{r} &= \langle l_1 m_1 | l_2 m_2 \rangle = \\ &= \sum_{kq} (-1)^{l_1 - m_1} \begin{pmatrix} l_1 & k & l_2 \\ -m_1 & q & m_2 \end{pmatrix} S_k(l_1 l_2) C_q^{k*}(\mathbf{R}_{12}), \end{aligned} \quad (7)$$

where $\begin{pmatrix} l_1 & k & l_2 \\ m_1 & q & m_2 \end{pmatrix}$ is a Wigner coefficient [2], k obeys the triangle rule $|l_1 - l_2| \leq k \leq l_1 + l_2$ given even values of $l_1 + k + l_2$. In particular, for $l_1 = 1$, $l_2 = 2$ (pd -overlap), k can have only two values: $k = 1, 3$. In the coordinate system with O_z directed along the radius-vector \mathbf{R}_{12}

$$\begin{aligned} \langle 10 | 20 \rangle = S_{pd\sigma} &= -\sqrt{\frac{2}{15}} S_1(pd) + \sqrt{\frac{3}{35}} S_3(pd), \\ \langle 1 \pm 1 | 2 \pm 1 \rangle = S_{pd\pi} &= -\sqrt{\frac{1}{10}} S_1(pd) - \sqrt{\frac{1}{35}} S_3(pd), \end{aligned}$$

where $S_{pd\sigma}$ and $S_{pd\pi}$ are overlap integrals on the σ - and π -bonds, respectively.

Thus, taking into account that the bare atomic functions $\varphi_{fnlm}(\mathbf{r})$ form a basis of the irreducible representation $D^{(l)}$ of the rotation group, we can rewrite $\psi_{fnlm}(\mathbf{r})$ as follows:

$$\begin{aligned} \psi_{fnlm}(\mathbf{r}) = & \varphi_{fnlm}(\mathbf{r}) - \frac{1}{2[l]^{1/2}} \sum_k \sum_{f'n'l'} S_k(f'n'l'; fnl) \left[C^k(\mathbf{R}_{ff'}) \times \varphi_{f'n'l'}(\mathbf{r} - \mathbf{R}_{ff'}) \right]_m^l + \\ & + \frac{3}{8[l]^{1/2}} \sum_{k_1 k_2} \sum_{f'n'l'} \sum_{f''n''l''} [l'']^{-1/2} S_{k_1}(f'n'l'; f''n''l'') S_{k_2}(f''n''l''; fnl) \times \\ & \times \left[C^{k_1}(\mathbf{R}_{ff''}) \times [C^{k_2}(\mathbf{R}_{ff'}) \times \varphi_{f'n'l'}(\mathbf{r} - \mathbf{R}_{ff'})]^{l''} \right]_m^l + \dots, \end{aligned} \quad (8)$$

for even values of $(k + l' + l)$, $(k_1 + k_2 + l' + l)$, ... Here we made use of direct products $[\cdot \times \cdot]$ of irreducible tensorial operators [2], $[l] = 2l + 1$. Unified tensorial form of different terms in the right-hand side unambiguously points to the same transformational properties of the bare atomic function $\varphi_{fnlm}(\mathbf{r})$ and its orthogonalized counterpart $\psi_{fnlm}(\mathbf{r})$ (Slater-Koster theorem [3]), however, with regard to rotations of the laboratory system, rather than the physical one. Indeed, the former transformation involves coordinates both of the electron (\mathbf{r}) and the lattice (\mathbf{R}_f), while the latter concerns only the electron coordinates (\mathbf{r}). We should emphasize here the preservation of the transformational properties with regard to any rotations rather than certain rotations from the local point group.

Within a linear approximation, the orthogonalization gives rise to a mixing of atomic functions centered at different sites (a nonlocal hybridization), while within a quadratic approximation the on-site mixing of different atomic nlm -functions appears (a local hybridization). The quadratic correction can be written as follows:

$$\Delta\psi_{fnlm}(\mathbf{r}) = \sum_k \sum_{f''} \sum_{n'l'} S_k(ff''n'l'; ff''nl) \left[C^k(\mathbf{R}_{ff''}) \times \varphi_{fn'l'}(\mathbf{r}) \right]_m^l, \quad (9)$$

where

$$S_k(ff''n'l'; ff''nl) = \frac{3}{8[l]^{1/2}} \sum_{k_1 k_2} \sum_{n''l''} (-1)^{k_2} [k]^{1/2} \begin{bmatrix} k_1 & k_2 & k \\ 0 & 0 & 0 \end{bmatrix} \left\{ \begin{matrix} k_1 & k_2 & k \\ l' & l & l'' \end{matrix} \right\} \times \quad (10)$$

$$\times S_{k_1}(fn'l'; f''n''l'') S_{k_2}(f''n''l''; fnl),$$

given even values of $k + l' + l$, $\begin{bmatrix} k_1 & k_2 & k \\ 0 & 0 & 0 \end{bmatrix}$ and $\left\{ \begin{matrix} k_1 & k_2 & k \\ l' & l & l'' \end{matrix} \right\}$ are the Clebsch-Gordan coefficient and $6j$ -symbol, respectively [2]. Structure of $S_k(ff''n'l'; ff''nl)$, in particular, summation on $n''l''$ implies significant troubles with its reliable estimates. Usually, this quantity is of the order of $S_{pd}^2 \sim 0.01$ for typical $3d$ oxides. It is worth noting that the orthogonalization does enhance the role of high-energy unfilled excited states which are characterized by a strong overlap with atomic functions of the filled orbitals of neighboring atoms.

Application of the irreducible tensor technique to the orthogonalized $fnlm$ -functions provides a regular procedure for a revision of different effects which are typical for many-electron atomic systems. Below we address different properties of quasi-atoms which are composed of “orthogonalized” electrons.

3. “Delocalization” of the orbital atomic operators

Delocalization of atomic orbitals in orthogonalized ψ_{fnlm} functions gives rise to an unconventional effect of a “delocalization” of the orbital atomic operators. Indeed, a local irreducible tensorial operator $\hat{V}_\alpha^a(fnl)$, working within the conventional nonorthogonalized basis set of the $(2l + 1)$ nlm -states for the f -atom according to the Wigner-Eckart theorem

$$\langle nlm | \hat{V}_\alpha^a(fnl) | nlm' \rangle = (-1)^{l-m} \begin{pmatrix} l & a & l \\ -m & \alpha & m' \end{pmatrix} \langle nl | \hat{V}^a(fnl) | nl \rangle, \quad (11)$$

will be described within the basis set of the orthogonalized counterparts ψ_{fnlm} by an equivalent operator

$$\begin{aligned}
 & \hat{V}_\alpha^a(fnl) + 2 \sum_{kb} (-1)^b [l, a]^{1/2} S_k(f f'' nl; f f'' nl) \left\{ \begin{matrix} k & a & b \\ l & l & l \end{matrix} \right\} \left[\hat{V}^b(fnl) \times C^k(\mathbf{R}_{ff''}) \right]_\alpha^a \\
 & + \frac{1}{4[a]^{1/2}} \sum_{k_1 k_2 k b} \sum_{f' n' l'} [b][k]^{1/2} S_{k_1}(f' n' l'; fnl) S_{k_2}(f' n' l'; fnl) \langle n' l' | \hat{V}^a(f' n' l') | n' l' \rangle \\
 & \left[\begin{matrix} k_1 & k_2 & k \\ 0 & 0 & 0 \end{matrix} \right] \left\{ \begin{matrix} k_1 & l' & l \\ k_2 & l' & l \\ k & a & b \end{matrix} \right\} \left[\hat{v}^b(fnl) \times C^k(\mathbf{R}_{ff'}) \right]_\alpha^a \\
 & \text{given even values of } k \text{ and } (a+b), \left\{ \begin{matrix} k_1 & l' & l \\ k_2 & l' & l \\ k & a & b \end{matrix} \right\} \text{ is the } 9j\text{-symbol [2], } [a, l] = (2a+1)(2l+1).
 \end{aligned} \tag{12}$$

Here $\hat{v}^b(fnl)$ is a b -rank irreducible tensorial operator whose submatrix element equals one. The Exp. (12) describes the delocalization effect for different quantities related to tensorial operators. Note the emergence of tensorial operators $\hat{v}_\beta^b(fnl)$ whose rank differs from that of (a) of the bare operator.

4. Effective orbital momentum

Obviously that, at variance with the $\varphi_{fnlm}(\mathbf{r})$ functions, the $\psi_{fnlm}(\mathbf{r})$ functions are not the eigenfunctions for operators $\hat{\mathbf{I}}^2$ and \hat{l}_z (the square and z -component of the orbital momentum operator), though they form a basis of the irreducible representation $D^{(l)}$ of the rotation group. Nonetheless, one can introduce an effective orbital momentum, or quasi-momentum $\hat{\mathbf{I}}$ as follows:

$$\hat{\mathbf{I}} = -i\hbar \left([\mathbf{r} \times \nabla_r] + \sum_{\mathbf{R}_f} [\mathbf{R}_f \times \nabla_{R_f}] \right). \tag{13}$$

By analogy with a free atom we can introduce an addition of the orbital quasi-momenta thus forming many-electron configurations and wave functions $|\tilde{L}SM_{\tilde{L}}M_S\rangle$ for the $^{2S+1}\tilde{L}$ terms of the many-electron quasi-free atom, which incorporates all the effects of the orthogonalization of one-electron states for different sites.

The orthogonalization procedure preserves many (though not all) advantages of the free atom theory based on the application of the theory of the rotation group. Thus the Wigner-Eckart theorem for conventional irreducible tensorial operators, such as spherical harmonic $C_q^k(\mathbf{r})$, orbital momentum $\hat{\mathbf{I}}$ and other orbital operators acting in the conventional \mathbf{r} space, does not work on $\psi_{fnlm}(\mathbf{r})$ -functions, because matrix elements will depend parametrically on the lattice vectors.

However, in practice, e.g., for Zeeman coupling one should address a true orbital momentum whose relations with effective orbital momentum will have a nontrivial form. Within a $|\tilde{L}M_{\tilde{L}}\rangle$ multiplet, these relations are as follows:

$$\hat{L}_i = a_{ij} \tilde{L}_j + a_{ijkl} \tilde{L}_j \tilde{L}_k \tilde{L}_l + \dots, \tag{14}$$

where we meet unconventional tensorial linear and different nonlinear terms.

The second-rank tensor a_{ij} for the systems such as a $3d$ ion in oxides can be written as follows:

$$a_{ij} = a\delta_{ij} + \Delta a_{ij}, \tag{15}$$

where $a \leq 1$, but $\Delta a_{ij} \sim S^2$ (S is a cation-anion overlap integral), and the difference between a and 1 is of the order of S^2 . Point symmetry puts distinct limitations on the a_{ij} tensor, e.g., for a cubic symmetry, $\Delta a_{ij} = 0$.

Complex nonlinear relations between true and effective orbital momenta result in a nontrivial form of Zeeman coupling

$$\hat{V}_Z = \mu_B(\hat{\mathbf{L}} \cdot \mathbf{H}) = \mu_B \left\{ \hat{\mathbf{L}}\hat{a}^{(2)}\mathbf{H} + \hat{\mathbf{L}}\hat{\mathbf{L}}\hat{\mathbf{L}}\hat{a}^{(4)}\mathbf{H} + \dots \right\}, \quad (16)$$

where we make use of a symbolic form of a tensorial product of vector operators. Thus the Zeeman coupling acquires novel features due to the orthogonalization procedure: i) a reduction of the effective orbital g -factor as compared with its bare value $g_l=1$ with its anisotropy for low-symmetry sites; ii) emergence of a nonlinear Zeeman coupling. These features evolve from the orthogonalization procedure as a zero order perturbation effect prior to effects of crystal field and covalency.

5. Electric dipole moment of a “free quasi-atom”

Orthogonalized $\psi_{fnlm}(\mathbf{r})$ functions for noncentrosymmetric sites f , at variance with bare $\varphi_{fnlm}(\mathbf{r})$ functions, do not have a definite parity. This immediately gives rise to a nonzero electric dipole polarization of such a “free quasi-atom”.

Taking into account the local hybridization effects (10), we can write one-electron matrix element of electric dipole moment as follows

$$\begin{aligned} \langle \psi_{fnlm} | \hat{d}_q | \psi_{fnlm} \rangle &= \frac{2e}{\sqrt{3}} [l]^{\frac{1}{2}} \sum_{a,k} \sum_{f''} \sum_{n'l'} [a](l||C^1||l') r_{nl;n'l'} S_k(ff''n'l'; ff''nl) \times \\ &\times \left\{ \begin{matrix} 1 & k & a \\ l & l & l' \end{matrix} \right\} (-1)^{l-m} \begin{pmatrix} l & a & l \\ -m & \alpha & m' \end{pmatrix} \begin{bmatrix} a & k & 1 \\ \alpha & q' & q \end{bmatrix} C_{q'}^k(\mathbf{R}_{ff''}), \end{aligned} \quad (17)$$

where $a = 0, 2, \dots, 2l$ and $k = a \pm 1$ are even and odd numbers, respectively. Here $(l||C^1||l')$ is a submatrix element of a tensorial harmonic [2],

$$r_{nl;n'l'} = \int_0^\infty R_{nl}(r) r^3 R_{n'l'}(r) dr$$

is a dipole radial integral. Thus within the ψ_{fnlm} basis set with a certain fnl the dipole moment operator can be replaced by an effective (equivalent) operator as follows

$$\hat{d}_q = \sum_{a,k,f'} d_{ak}(R_{ff'}) \left[\hat{v}_\alpha^a(\mathbf{1}) \times C^k(\mathbf{R}_{ff'}) \right]_q^1, \quad (18)$$

where $\hat{v}_\alpha^a(l)$ is an orbital a -rank irreducible tensorial operator with the unit submatrix. In Cartesian coordinates the Exp. (18) can be written as follows

$$\hat{d}_i = d_i^{(0)} + \frac{1}{2} d_{ijk}^{(2)} \{ \hat{l}_j, \hat{l}_k \} + \text{terms with } a = 4, \dots, \quad (19)$$

where $\{ \hat{l}_j, \hat{l}_k \} = \hat{l}_j \hat{l}_k + \hat{l}_k \hat{l}_j$. Of a particular interest are the terms with nonzero values $a = 2, 4, \dots$, which directly relate the electric polarization to the degenerated or quasi-degenerated orbital state, in particular, to quadrupole ($a = 2$), octupole ($a = 4$) electronic momenta of quasi-atoms and give rise to an orbital magnetoelectric coupling. Strictly speaking, the electric polarization induced by the orthogonalization can amount to big values due to both big overlap and radial integrals $r_{nl;n'l'} \sim 1 \text{ \AA}$.

The expression (18) can be easily generalized for many-electron quasi-atomic configurations. For certain terms ^{2S+1}L for the nl^N shell of equivalent electrons one should replace the one-electron operator $\hat{v}_\alpha^a(\mathbf{1})$ in (18) by the equivalent many-electron orbital operator $\hat{V}_\alpha^a(\mathbf{L})$ which

acts on the $|LM_L\rangle$ basis set according to the Wigner-Eckart theorem

$$\langle LM_L|\hat{V}_\alpha^a(\mathbf{L})|L'M_{L'}\rangle = (-1)^{L-M_L} \begin{pmatrix} L & a & L' \\ -M_L & \alpha & M_{L'} \end{pmatrix} U_{SL;SL'}^{(a)}, \quad (20)$$

where $U_{SL;SL'}^{(a)}$ is the spectroscopic Racah coefficient [4]. For the $^{2S+1}L_J$ multiplets the $\hat{V}_\alpha^a(\mathbf{L})$ operator in (18) should be replaced by the equivalent many-electron orbital operator $\hat{V}_\alpha^a(\mathbf{J})$ which acts on the $|SLJM_J\rangle$ basis set according to the Wigner-Eckart theorem

$$\langle SLJM_J|\hat{V}_\alpha^a(\mathbf{J})|SL'J'M_{J'}\rangle = (-1)^{J-M_J} \begin{pmatrix} J & a & J' \\ -M_J & \alpha & M_{J'} \end{pmatrix} U_{SLJ;SL'J'}^{(a)}, \quad (21)$$

where [4]

$$U_{SLJ;SL'J'}^{(a)} = (-1)^{S+a+L'+J} \sqrt{(2J+1)(2J'+1)} \begin{Bmatrix} L & J & S \\ J' & L' & a \end{Bmatrix} U_{SL;SL'}^{(a)}. \quad (22)$$

A consideration of the linear overlap effects needs the knowledge of two-site dipole matrix elements which can be written as follows:

$$\begin{aligned} \langle \varphi_{fnlm}|\hat{d}_q|\varphi_{f'n'l'm'}\rangle &= \sum_{a,\alpha,k,q'} (-1)^{l-m} \begin{pmatrix} l & a & l' \\ -m & \alpha & m' \end{pmatrix} \\ \langle fnl||d^{ak}||f'n'l'\rangle &\begin{bmatrix} a & k & 1 \\ \alpha & q' & q \end{bmatrix} C_{q'}^k(\mathbf{R}_{ff'}). \end{aligned} \quad (23)$$

Accordingly, the contribution to dipole matrix on the ψ_{fnlm} basis with a certain set of quantum numbers fnl acquires a tensorial form which is similar to (18). As an obvious practical implication of the Exp. (18), we should point on the calculation of the probabilities of the intra-configurational electro-dipole transitions, e.g., $d-d$ transitions for $3d$ compounds or $f-f$ transitions for $4f$ compounds, which are dipole-allowed for noncentrosymmetric quasi-atoms.

Judd-Ofelt theory of effective electric dipole moment for noncentrosymmetric ions

The Judd-Ofelt theory of effective electric dipole moment for noncentrosymmetric ions [5] takes into account the admixing of configurations of opposite parity due to the odd-parity crystal field

$$\hat{V}_{cf} = \sum_{kq} A_{kq}^* r^k C_q^k(\theta, \phi), \quad (24)$$

where A_{kq} are crystal field parameters. The effective dipole moment operator can be written out similarly to (18) as follows

$$\hat{d}_q = \sum_{a,k} \tilde{d}_{ak} \left[\hat{v}^a(\mathbf{1}) \times A^k \right]_q^1, \quad (25)$$

where

$$\tilde{d}_{ak} = -\frac{1}{\sqrt{3}}(2a+1)\Xi(ka), \quad (26)$$

and

$$\begin{aligned} \Xi(ka) &= 2(2a+1) \sum_{l'} (2l'+1) (-1)^{l+l'} \begin{Bmatrix} 1 & a & k \\ l & l' & l \end{Bmatrix} \\ &\begin{pmatrix} l & 1 & l' \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l' & k & l \\ 0 & 0 & 0 \end{pmatrix} \frac{\langle 4f|r|n'l'\rangle \langle 4f|r^k|n'l'\rangle}{E_{n'l'} - E_{nl}}. \end{aligned} \quad (27)$$

6. The overlap contribution to orbital magnetoelectric coupling

For paramagnetic ions with an orbital (quasi)degeneracy, the thermal expectation value $\langle \hat{V}_\alpha^a(l) \rangle$ in (18) can strongly depend on the magnetic field, either internal molecular or the external one, thus providing an effective magnetic control of electric polarization. Indeed, in the absence of crystal-field and spin-orbital effects, the thermal expectation value can be represented as follows:

$$\langle \hat{V}_\alpha^a(l) \rangle_T = \langle \hat{V}_0^a(l) \rangle_T C_\alpha^a(\mathbf{L}), \quad \langle V_0^a \rangle_T = \langle V_0^a \rangle_0 \rho_a(H, T), \quad (28)$$

where $C_\alpha^a(\mathbf{L})$ is tensorial spherical harmonic with a classical vector \mathbf{L} as an argument, $\langle V_\alpha^a \rangle$ the thermal expectation value with $\langle V_0^a \rangle_0 = \begin{pmatrix} L & a & L \\ -L & 0 & L \end{pmatrix}$, and $\rho_a(H, T)$ is a temperature factor, e.g.:

$$\rho_0 = 1; \quad \rho_2 = \frac{\langle 3L_z^2 - L(L+1) \rangle}{L(2L-1)}. \quad (29)$$

Within a molecular field approximation $C_\alpha^a(\mathbf{L}) \equiv C_\alpha^a(\mathbf{H})$. Thus in frames of our simplifications we arrive at a very interesting expression

$$\langle \hat{d}_q \rangle = \sum_{a,k,f'} d_{ak}(R_{ff'}) \langle V_0^a(L) \rangle_0 \rho_a(H, T) \left[C^a(\mathbf{H}) \times C^k(\mathbf{R}_{ff'}) \right]_q^1, \quad (30)$$

that demonstrates a magnetic control of the electric polarization in the most clear way. In Cartesian coordinates the Exp. (18) can be written as follows

$$\hat{d}_i = d_i^{(0)} + d_{ijk}^{(2)}(H, T) h_j h_k + \text{terms with } a = 4, \dots, \quad (31)$$

where $d_{ijk}^{(2)}(H, T) \propto \rho_2(H, T)$, and $\mathbf{h} = \mathbf{H}/H$.

For rare-earth ions with $4f^n$ shell and strong spin-orbital coupling, the electric polarization for the SLJ -multiplet can be easily written as follows:

$$\begin{aligned} \langle \hat{d}_q \rangle &= \sum_{a,k,f'} (-1)^{S+a+L+J} [J] \begin{Bmatrix} L & J & S \\ J & L & k \end{Bmatrix} \times \\ &\times d_{ak}(R_{ff'}) \langle V_0^a(J) \rangle_0 \rho_a(T) \left[C^a(\mathbf{H}) \times C^k(\mathbf{R}_{ff'}) \right]_q^1. \end{aligned} \quad (32)$$

It should be noted that the expressions (30) and (32) define both the field and temperature dependence of the electric dipole moment, however, given a complete neglect of the the crystal field quenching effects. These effects are usually much stronger than Zeeman coupling, in particular, for $3d$ ions. However, for some rare earth ions with so-called quasi-doublets in ground state the Zeeman and crystal field effects can compete with each other, and our Exps. (18) and (32) predict a rather strong magnetoelectric coupling. Such a situation is realized, e.g., for Tb^{3+} ion in the multiferroic terbium manganate TbMn_2O_5 (see, e.g., Ref. [6]).

For the $3d$ ions, the crystal field quenches both the orbital momenta and orbital magnetoelectric coupling. However, due to the spin-orbital coupling/mixing, the orbital effects give rise to different effective spin interactions, e.g., the emergence of an orbital contribution to effective spin g -tensor and single-ion spin anisotropy. Furthermore, the orbital operator $\hat{V}_\alpha^a(\mathbf{L})$ given $a = 2, 4, \dots$ for orbitally nondegenerated ground state can be replaced by an effective spin operator. This is relatively easy to perform for so-called S -type $3d$ ions, which have an orbitally nondegenerate ground state of the A_1 or A_2 type for a high-symmetry octahedral, cubic or tetrahedral crystal field. These are ions, e.g., with $3d^3$ (Cr^{3+} , Mn^{4+}), $3d^5$ (Fe^{3+} , Mn^{2+}),

$3d^8$ (Ni^{2+} , Cu^{3+}) configurations with an octahedral surroundings. Strictly speaking, first we should replace the $\hat{V}_\alpha^a(\mathbf{L})$ by a linear combination

$$\hat{V}_\nu^{a\gamma}(\mathbf{L}) = \sum_\alpha c_{a\alpha}^{\gamma\nu} \hat{V}_\alpha^a(\mathbf{L}), \quad (33)$$

which form a basis of the irreducible representation D^γ (line ν) of the point symmetry group, O_h or T_d . Here $\gamma = E$ or T_2 given $a = 2$, $\gamma = A_1, E, T_1$ or T_2 given $a = 4$. Next we can replace the orbital operator by a spin equivalent as follows

$$\langle \hat{V}_\nu^{a\gamma}(\mathbf{L}) \rangle_{\text{GS}} = \eta_{a\gamma} \hat{V}_\nu^{a\gamma}(\mathbf{S}), \quad (34)$$

where the $\langle \dots \rangle_{\text{GS}}$ means a mapping on the ground state. The $\eta_{a\gamma}$ parameters depend on the type of the 3d ion: $\eta_{2\gamma} \sim (\zeta_{3d}/\Delta)^2 \leq 10^{-3}$, here ζ_{3d} is a spin-orbital constant, Δ is a mean energy of excited T -terms. Thus for S -type ions we can transform the orbital dipole moment operator (18) into an effective spin operator which acts within the SM_S -multiplet of the ground state as follows

$$\hat{d}_q = \sum_{a\gamma\nu} R_{q\nu}^{(a\gamma)} \hat{V}_\nu^{a\gamma}(\mathbf{S}), \quad (35)$$

where

$$R_{q\nu}^{(a\gamma)} = \sum_{a,k,f'} d_{ak}(R_{ff'}) \eta_{a\gamma} \sum_{\alpha q'} c_{a\alpha}^{\gamma\nu*} \begin{bmatrix} a & k & 1 \\ \alpha & q' & q \end{bmatrix} C_{q'}^k(\mathbf{R}_{ff'}). \quad (36)$$

As in (33), $\hat{V}_\nu^{a\gamma}(\mathbf{S}) = \sum_\alpha c_{a\alpha}^{\gamma\nu} \hat{V}_\alpha^a(\mathbf{S})$, however, with a more simple spin matrix for $\hat{V}_\alpha^a(\mathbf{S})$:

$$\langle SM_S | \hat{V}_\alpha^a(\mathbf{S}) | SM_{S'} \rangle = (-1)^{S-M_S} \begin{pmatrix} S & a & S \\ -M_S & \alpha & M_{S'} \end{pmatrix}. \quad (37)$$

The spin irreducible tensorial operators $\hat{V}_\nu^{a\gamma}(\mathbf{S})$ can be transformed into a Cartesian form, e.g.

$$\hat{V}_0^{2E}(\mathbf{S}) = 2 \frac{3\hat{S}_z^2 - S(S+1)}{\sqrt{(2S+3)^{(5)}}}; \quad \hat{V}_2^{2E}(\mathbf{S}) = 2\sqrt{3} \frac{(\hat{S}_x^2 - \hat{S}_y^2)}{\sqrt{(2S+3)^{(5)}}}, \dots, \quad (38)$$

where $(2S+3)^{(5)} = (2S+3)(2S+2)\dots(2S-1)$. In Cartesian coordinates the Exp. (35) can be reduced to a standard form

$$\hat{d}_i = d_i^{(0)} - \frac{1}{2} R_{ijk} \{ \hat{S}_j, \hat{S}_k \} + \text{terms with } a = 4, \dots, \quad (39)$$

widely adopted in the theory of the electric field effects in electron spin resonance (ESR) [7].

The magnetoelectric effect due to local noncentrosymmetry was addressed recently in Ref. [8]. The authors started with the nonorthogonalized basis set of $3d$ orbitals, next considered an odd-parity crystal field and spin-orbital coupling as perturbations. Finally, they arrived at an expression which is similar to (39). Furthermore, in fact, the microscopic consideration in Ref. [8] reproduces, however, with strong simplification the well-known paper by Kiel and Mims [9] on electric field effect in ESR for Mn^{2+} ions in CaWO_4 .

It is interesting that the experimental findings of this and many other papers on the electric field effect in ESR [7] can be used for a direct estimation of the single-ion contribution to the magnetoelectric coupling in different multiferroics. For instance, for well known multiferroic MnWO_4 [10] one might use the parameters R_{ijk} measured for Mn^{2+} ions in CaWO_4 [9]: being normalized to unit cell volume of MnWO_4 [10] ($V_c \approx 138 \text{ \AA}^3$), these are as follows: $|R_{123}| \approx 1.8$, $|R_{113}| \approx 1.5$, $|R_{311}| \approx 0.3$, $|R_{312}| \approx 5.0 \mu\text{C m}^{-2}$ ($R_{223} = -R_{113}$, $R_{213} = R_{123}$). It should be noted

that for Mn^{2+} ions in SrWO_4 these parameters are nearly two times larger [11]. Taking account of two Mn^{2+} ions in unit cell and a rather large value (≤ 6) of the spin factor in (39), one may conclude that the single-ion mechanism can be a significant contributor to the ferroelectric polarization observed in MnWO_4 ($P_b \sim 50 \mu\text{C m}^{-2}$) [10]. It is interesting that very recent quantitative estimates of the spin-dependent ferroelectric polarization in MnWO_4 based on the low-energy model, derived from the first-principles electronic structure calculations [12] showed values which are an order of magnitude less than the experimental ones.

It is worth noting that the single-ion term (39) does not produce the magnetoelectric coupling for quantum spins $S = \frac{1}{2}$ (e.g., Cu^{2+}) due to a kinematic constraint: $0 \leq a \leq 2S$.

7. Summary

Making use of the Löwdin orthogonalization for one-electron atomic $\varphi_{fnlm}(\mathbf{r})$ wave functions, we arrive at a basis set of orthogonalized $\psi_{fnlm}(\mathbf{r})$ orbitals which formally preserve symmetric properties of the bare $\varphi_{fnlm}(\mathbf{r})$ orbitals. Instead of many-electron atomic configurations composed of non-orthogonalized nlm -orbitals, we arrive at quasi-atoms composed of the orthogonalized nlm -counterparts. Formal conservation of the rotational symmetry allows to apply the powerful technique of irreducible tensorial operators [2] and Racah algebra [4] to description of the quasi-atoms, in particular, to the overlap contribution in the crystal field parameters [13].

As an illustration, we addressed a single-ion contribution to magnetoelectric coupling which is usually missed in current studies on multiferroics. A regular procedure has been developed for calculation of the overlap contribution to the single-ion orbital magnetoelectric coupling both for $3d$ - and $4f$ -ions. In a sense, the overlap contribution resembles the point charge contribution to a crystal field, this correctly describes both the lattice symmetry and the symmetry of electronic states, and provides reasonable semiquantitative estimates. Furthermore, making use of experimental data for the electric field effect in ESR, we have shown that the single-ion magnetoelectric coupling can be a leading mechanism of multiferroicity, e.g., in MnWO_4 .

Acknowledgments

Supported by Act 211 Government of the Russian Federation, agreement No 02.A03.21.0006 and by the Ministry of Education and Science, projects No 2277 and No 5719.

References

1. Löwdin P.-O. *Adv. Phys.* **5**, 1 (1966)
2. Varshalovich D.A., Moskalev A.N., Khersonskii V.K. *Quantum Theory of Angular Momentum*, World Scientific (1988)
3. Slater J.C., Koster G.F. *Phys. Rev.* **94**, 1498 (1954)
4. Sobelman I.I. *Introduction to the Theory of Atomic Spectra*, Pergamon Press, Oxford, New York (1972)
5. Judd B. *Phys. Rev.* **127**, 750 (1962)
6. Chang L.J., Su Y., Schweika W., Bruckel T., Chen Y.Y., Jang D. S., Liu R.S. *Physica B* **404**, 2517 (2009)
7. Mims W.B. *The Linear Electric Field Effect in Paramagnetic Resonance*, Oxford University Press, UK (1976)
8. Sakhnenko V.P., Ter-Oganessian N.V. *J. Phys.: Cond. Mat.* **24**, 266002 (2012)

9. Kiel A., Mims W.B. *Phys. Rev.* **153**, 378 (1967)
10. Taniguchi K., Abe N., Takenobu T., Iwasa Y., Arima T. *Phys. Rev. Lett.* **97**, 097203 (2006)
11. Kiel A., Mims W.B. *Phys. Rev. B* **3**, 2878 (1971)
12. Solovyov I.V. *Phys. Rev.* **87**, 144403 (2013)
13. Malkin B.Z., Kaminskii A.A., Agamalyan N.R., Bumagina L.A., Butaeva T.I. *Phys. Stat. Sol. B* **110**, 417 (1982)