DR-4

AN EFFICIENT METHOD FOR THE SYNTHESIS OF DIHYDROPYRIMIDINES USING BRØNSTED ACIDIC IONIC LIQUID: A SOLVENT AND HEATING FREE REACTION

Satyajit Samanta,¹ Rana Chatterjee,¹ Sougata Santra,² Adinath Majee¹¹Department of Chemistry; Visva-Bharati (A Central University), Santiniketan 731235, India.²Ural Federal University, 19 Mira St., Yekaterinburg-620002, Russian Federation.

E-mail: satyasamanta527@gmail.com

Abstract. Dihydropyrimidinones occupy a considerable position in the dominion of natural and synthetic organic chemistry, due to their various therapeutic and pharmacological properties, such as antiviral, antibacterial, antihypertensive and antitumor effects.¹ Dihydropyrimidinones scaffold rose as mainstay of several drugs which are used as calcium channel blockers.² Recently, isolated marine alkaloids have been connected with dihydropyrimidine-5-carboxylate core and found to be potent to HIV-gp-120 CD4 inhibitors.³ Years back, Biginelli reported the most simple and straight forward procedure for the synthesis of DHPMs.⁴ However, From bingeing, many methods have been developed for the synthesis of DHPMs and its derivative using various types of catalytic reagents.⁵ Considering the synthetic importance of Dihydropyrimidinones we developed an efficient and greener method for the synthesis of these compounds using Brønsted acidic ionic liquid (BAILs) {1-Butane sulfonic acid-3-methylimidazolium tosylate, [BSMIM]OTs} under solvent free condition and at room temperature (Scheme 1).

Scheme 1. Synthesis of Dihydropyrimidinones using BAILs

References

1. 100 years of the Biginelli dihydropyrimidine synthesis / O. Kappe // Tetrahedron. – 1993. – Vol. 49. – P. 6937-6963.
2. Dihydropyrimidine calcium channel blockers. 4. Basic 3-substituted-4-aryl-1,4-dihydropyrimidine-5-carboxylic acid esters. Potent antihypertensive agents / G. C. Rovnyak, K. S. Atwal, A. Hedberg, S. D. Kimball, S. Moreland, J. Z. Gougoutas, B. C. O'Reilly, M. F. Malley // J. Med. Chem. – 1992. – Vol. 35. – P. 3254.
3. Biomimetic synthesis of (.,+.)-crambines A, B, C1, and C2. Revision of the structure of crambines B and C1 / B. B. Snider, Z. Shi // J. Org. Chem. – 1993. – Vol. 58 – P. 3828. 4. Biginelli Reaction / P. Biginelli // Ber. Dtsch. Chem. Ges. – 1893. – Vol. 26. – P. 447.
4. Lanthanide Triflate Catalyzed Biginelli Reaction. One-Pot Synthesis of Dihydropyrimidinones under Solvent-Free Conditions / Y. Ma, C. Qian, L. Wang, M. Yang // J. Org. Chem. – 2000, – Vol. 65. – P. 3864.

This work was funded by Russian Science Foundation (Ref. No. 18-73-00301).