





2-(4-Oxo-1,3-thiazolidin-2-ylidene)acetamid as promising scaffold for designing new antifungal compounds

Konstantin L. Obydenov* , Tatiana A. Kalinina , Daria V. Ryabova, Maria F. Kosterina, Tatiana V. Glukhareva 

Institute of Chemical Engineering, Ural Federal University, Ekaterinburg 620009, Russia

* Corresponding author: k.l.obydenov@urfu.ru



This paper belongs to the MOSM2022 Special Issue.

Abstract

1,3-Thiazolidin-4-one derivatives with an exocyclic C=C double bond in position 2 of the hetero ring have a wide spectrum of biological activity, but their fungicidal activity has not been studied as much as it should be. This paper presents a simple and convenient approach for obtaining potential antifungal agents based on 2-(4-oxo-1,3-thiazolidin-2-ylidene)acetamides. The first examples of evaluating the fungicidal activity of 8 obtained compounds on 8 strains of phytopathogenic fungi are presented. A highly active compound **4e** with EC₅₀ of 0.85 and 2.29 µg/mL against *A. solani* and *P. lingam*, respectively, was found to be promising for further study.

Keywords

1,3-thiazolidine
cyanoacetamide
exocyclic double bond
fungicide
biological activity

Received: 08.12.22

Revised: 21.12.22

Accepted: 22.12.22

Available online: 29.12.22

© 2022, the Authors. This article is published in open access under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).




1. Introduction

Derivatives of 1,3-thiazolidin-4-one with a double C=O, C=N, C=S, C=C exocyclic bond in position 2 of the hetero ring have a wide range of biological activity: antituberculosis [1], antioxidant [2], anticancer [3, 4], antiinflammatory [5], anticonvulsant [6], antiviral [7, 8], trypanocidal [9], antiarrhythmic [10], antibacterial [11, 12] and fungicidal [12, 13]. Although 1,3-thiazolidines and 1,3-thiazolines have centers for polar interactions and hydrogen bonds [14], these heterocycles often act as a scaffold for substituents interacting with biotargets. Thus, the analysis of the crystal structure of the complex of succinate dehydrogenase with the inhibitor thiapronil **I** (pdb id: 6MYR) showed that thiapronil **I** does not form strong non-covalent interactions (hydrogen bonds and π-π interactions) with the enzyme due to the 1,3-thiazol-2-ylidene fragment [15]. Anticancer compounds **II** [16] and **III** [17] containing a 3,4,5-trimethoxyphenyl substituent are prominent examples as well. Due to this substituting group, compounds **II** and **III** bind to the bioreceptor tubulin (Scheme 1) [18].


It is known that 1,3-thiazolidine derivatives with C=O [2, 19], C=N [20–22], C=S [23] exocyclic bonds in position 2 of the hetero ring show fungicidal properties. However, information on the study of the antifungal properties of 1,3-thiazolidine with a C=C double bond in position 2 of the ring

is limited [12]. For example, 1,3-thiazolidin-2-ylidene **IV** is known to exhibit antifungal activity against human pathogens *C. albicans* and *C. neoformans* [24].

One of the convenient methods for the synthesis of 1,3-thiazolidin-2-ylidene derivatives is the condensation of cyanoacetamides **1** with thioglycolic acid **2** (Scheme 2) [25]. In this case, 1,3-thiazolidin-2-ylidenes **3** can be modified both at the NH and CH₂ groups of the hetero ring.



Scheme 1 Examples of active 1,3-thiazol-2-ylidene and 1,3-thiazolidin-2-ylidene derivatives.



Scheme 2 Synthesis of 1,3-thiazolidin-2-ylidenes.

As a result of varying the substituents in the acetamide fragment and the hetero ring, a wide range of 1,3-thiazolidin-2-ylidenes can be obtained for the search for biologically active compounds.

In this paper, we present the first results of using this approach to search for antifungal compounds.

2. Experimental

2.1. Synthesis of target compounds

¹H and ¹³C NMR spectra were recorded with a Bruker Avance II (Karlsruhe, Germany) spectrometer (400 MHz for ¹H, 100 MHz for ¹³C) using Me₄Si as an internal standard. The NMR spectra of all compounds are demonstrated in the Supporting Information (Figures S1–S10). Mass spectra were recorded with a Shimadzu GCMS-QP 2010 “Ultra” (Kyoto, Japan) in electron ionization (EI) mode (electron energy 70 eV). The Fourier transform infrared (FT-IR) spectra were obtained using a Bruker Alpha (ATR, ZnSe) spectrometer (Ettlingen, Germany). Elemental analyses were performed with a Perkin-Elmer 2400 Series II CHNS/O analyzer (Shelton, CT USA). Melting points were determined using a Stuart SMP 3 apparatus (Staffordshire, ST15 OSA, UK). The progress of the reactions and the purity of the compounds were monitored by thin-layer chromatography (TLC, Merck KGaA) in an ethyl acetate-hexane system.

The synthesis of compounds **3a–c** was carried out according to the published method [25].

Alkylation reaction of thiazolidines **3a–c** with the formation of products **4a–e** was carried out according to the published method [26].

(2Z)-N-benzyl-2-(4-oxo-1,3-thiazolidin-2-ylidene)acetamide (3a). Yield 0.37 g (72%), white powder, mp 207–209 °C (lit. 208–209 °C [25]).

(2Z)-2-(4-oxo-1,3-thiazolidin-2-ylidene)-N-phenylacetamide (3b). Yield 0.44 g (67%), white powder, mp 290–293 °C (decomp.) (lit. 282–285 °C [25]).

(2Z)-N-(2-methylphenyl)-2-(4-oxo-1,3-thiazolidin-2-ylidene)acetamide (3c). Yield 0.23 g (75%), white powder, mp 235–236 °C. IR spectrum, ν , cm⁻¹: 3205 (NH), 2969, 2594, 1702 (C=O), 1623 (C=O), 1600, 1576, 1549, 1492, 1456, 1427, 1399, 1367, 1319. ¹H NMR spectrum (400 MHz, DMSO-*d*₆), δ , ppm, (*J*, Hz): 2.19 (3H, *s*, CH₃); 3.67 (2H, *s*, CH₂); 5.91 (1H, *s*, CH=); 7.02 (1H, *t*, *J* = 7.2, Ar H); 7.00 (1H, *t*, *J* = 7.3, Ph *p*-H); 7.08–7.24 (2H, *m*, Ph H); 7.49 (1H, *d*, *J* = 7.4, Ph H); 9.08 (1H, *s*, NH); 11.46 (1H, *s*, NH). ¹³C NMR spectrum (100 MHz, DMSO-*d*₆), δ , ppm: 17.99 (CH₃); 32.00 (CH₂); 92.37 (C-2’); 124.36 (2C Ar); 125.78 (C Ar); 130.18 (C Ar); 130.77 (C Ar); 136.79 (C Ar), 153.92 (C-2);

165.40 (C=O), 174.18 (C=O). Found, %: C 58.05; H 4.87; N 11.28. C₁₂H₁₂N₂O₂S. Calculated, %: C 57.96; H 4.66; N 11.36.

(2Z)-N-benzyl-2-(3-methyl-4-oxo-1,3-thiazolidin-2-ylidene)acetamide (4a). Yield 0.50 g (64%), white powder, mp 183–186 °C. IR spectrum, ν , cm⁻¹: 3295 (NH), 1705 (C=O), 1635 (C=O), 1623, 1553, 1534, 1495, 1453, 1431, 1411, 1386, 1334. ¹H NMR spectrum (400 MHz, DMSO-*d*₆), δ , ppm, (*J*, Hz): 3.04 (3H, *s*, CH₃); 3.72 (2H, *s*, CH₂); 4.32 (2H, *d*, *J* = 5.4, CH₂N); 5.67 (1H, *s*, CH=); 7.22–7.34 (5H, *m*, Ph H); 8.22 (1H, *t*, *J* = 5.1, NH). ¹³C NMR spectrum (100 MHz, DMSO-*d*₆), δ , ppm: 29.47 (CH₂); 31.10 (CH₃); 93.13 (C-2’); 126.70 (C Ar); 127.24 (C Ar); 128.25 (C Ar); 139.81 (C Ar); 153.08 (C-2), 166.19 (C=O), 172.12 (C=O). EI-MS *m/z* (%): 262 [M]⁺ (90). Found, %: C 59.52; H 5.38; N 10.68. C₁₃H₁₄N₂O₂S. Calculated, %: C 59.32; H 5.37; N 10.56.

(2Z)-N-benzyl-2-(3-benzyl-4-oxo-1,3-thiazolidin-2-ylidene)acetamide (4b). Yield 0.31 g (76%), white powder, mp 144–147 °C. IR spectrum, ν , cm⁻¹: 3307 (NH), 1720 (C=O), 1649 (C=O), 1631, 1572, 1546, 1495, 1454, 1430, 1389, 1364, 1321. ¹H NMR spectrum (400 MHz, DMSO-*d*₆), δ , ppm, (*J*, Hz): 3.86 (2H, *s*, CH₂); 4.25 (2H, *d*, *J* = 5.5, CH₂N); 4.79 (2H, *s*, CH₂N); 5.65 (1H, *s*, CH=); 7.21–7.37 (10H, *m*, Ph H); 8.17 (1H, *t*, *J* = 5.2, NH). ¹³C NMR spectrum (100 MHz, DMSO-*d*₆), δ , ppm: 31.45 (CH₂); 42.49 (CH₂N); 46.38 (CH₂N); 94.32 (C-2’); 127.26 (C Ar); 127.90 (C Ar); 127.97 (C Ar); 128.75 (C Ar); 129.11 (C Ar); 135.56 (C Ar); 140.15 (C Ar); 152.47 (C-2), 166.53 (C=O), 173.21 (C=O). EI-MS *m/z* (%): 338 [M]⁺ (23.79). Found, %: C 67.43; H 5.36; N 8.28. C₁₉H₁₈N₂O₂S. Calculated, %: C 67.38; H 5.35; N 8.41.

(2Z)-2-(3-methyl-4-oxo-1,3-thiazolidin-2-ylidene)-N-phenylacetamide (4c). Yield 0.22 g (79%), white powder, mp 209–211 °C (lit. 205–208 °C [26,27]). IR spectrum, ν , cm⁻¹: 3447 (NH), 1714 (C=O), 1650 (C=O), 1617, 1608, 1596, 1567, 1494, 1443, 1421, 1405, 1346, 1311. ¹H NMR spectrum (400 MHz, DMSO-*d*₆), δ , ppm, (*J*, Hz): 3.09 (3H, *s*, CH₃); 3.78 (2H, *s*, CH₂); 5.80 (1H, *s*, CH=); 7.00 (1H, *t*, *J* = 7.3, Ph *p*-H); 7.28 (2H, *t*, *J* = 7.7, Ph H); 7.60 (2H, *t*, *J* = 7.9, Ph H); 9.87 (1H, *s*, NH). ¹³C NMR spectrum (100 MHz, DMSO-*d*₆), δ , ppm: 29.43 (CH₃); 31.00 (CH₃); 93.37 (C-2’); 118.50 (C Ar); 122.38 (C Ar); 128.45 (C Ar); 139.49 (C Ar); 154.76 (C-2), 164.79 (C=O), 171.95 (C=O). EI-MS *m/z* (%): 248 [M]⁺ (31). Found, %: C 58.05; H 4.87; N 11.28. C₁₂H₁₂N₂O₂S. Calculated, %: C 58.27; H 5.07; N 11.45.

(2Z)-2-(3-benzyl-4-oxo-1,3-thiazolidin-2-ylidene)-N-phenylacetamide (4d). Yield 0.54 g (80%), white powder, mp 214–217 °C (lit. 216–218 °C [27]).

(2Z)-2-(3-benzyl-4-oxo-1,3-thiazolidin-2-ylidene)-N-(2-methylphenyl)acetamide (4e). Yield 0.37 g (76%), white powder, mp 166–169 °C. IR spectrum, ν , cm⁻¹: 3274.81, 3062.51, 1713.51 (C=O), 1653.91 (C=O), 1639.00, 1578.96, 1566.82, 1535.97, 1494.73, 1455.75, 1397.69, 1364.59, 1316.31. ¹H NMR spectrum (400 MHz, DMSO-*d*₆), δ , ppm, (*J*, Hz): 2.16 (3H, *s*, CH₃); 3.90 (2H, *s*, CH₂); 4.85 (2H, *s*, CH₂N); 5.96 (1H, *s*, CH=); 7.02–7.48 (9H, *m*, Ar H); 9.06 (1H, *s*, NH). ¹³C NMR spectrum (100 MHz, DMSO-*d*₆), δ , ppm: 17.93 (CH₃);

31.02 (CH₂); 45.98 (CH₂N); 93.81 (C-2'); 124.08 (C Ar); 124.42 (C Ar); 125.84 (C Ar); 126.72 (C Ar); 127.51 (C Ar); 128.66 (C Ar); 130.22 (C Ar); 130.41 (C Ar); 135.04 (C Ar); 136.53 (C Ar); 153.86 (C-2); 165.00 (C=O); 172.78 (C=O). EI-MS *m/z* (%): 338 [M]⁺ (8.83). Found, %: C 67.43; H 5.36; N 8.28. C₁₉H₁₈N₂O₂S. Calculated, %: C 67.58; H 5.20; N 8.19.

2.2. Study of fungicidal activity

The fungicidal activity of compounds **3a-c** and **4a-e** was tested *in vitro* on *Alternaria solani* Sorauer MFP601021, *Botrytis cinerea* Pers. MFG 60449, *Colletotrichum coccodes* JS 161-1, *Fusarium solani* (Mart.) Sacc. MFG 70523, *Phytophthora infestans* (Mont.) de Bary, *Plenodomus lingam* (Tode:Fr.) Höhn. MF Br17-044, *Rhizoctonia solani* RCAM01785 and *Sclerotinia sclerotiorum* using the agar block method. *P. infestans* was isolated at Nankai University (Tianjin, China). *C. coccodes* was purchased from the All-Russian Collection of Industrial Microorganisms (Moscow, Russia). The remaining strains of fungi were purchased from the Russian Collection of Agricultural Microorganisms (St. Petersburg, Russia).

Solutions of the tested compounds were prepared at a concentration of 0.5 mg/mL by dissolving 5 mg of the compound in 1 mL of DMSO with the addition of 9 mL of water. A total 1 mL of the test solutions was added to sterile Petri dishes containing 9 mL of heated (60 °C) nutrient medium and then mixed in a laminar flow cabinet. Fungal discs (4 mm in diameter) were cup under aseptic conditions from a 7-day-old culture of the test fungus using a sterile cork borer. The mycelium discs were placed in the center of Petri dishes containing the culture medium at room temperature. A negative control was prepared with the culture medium and DMSO. The fungi were incubated at 25 °C (48 h for *R. solani*, 120 h for *A. solani*, *P. lingam*, 72 h for other fungi). After incubation, the diameter of fungal colonies was measured. The percentage of fungus growth inhibition was determined by the formula [28]:

$$I (\%) = [(C - T)/(C - 4 \text{ mm})] \cdot 100, \quad (1)$$

where *I* (%) – the degree of inhibition of mycelial growth, *T* (mm) – the mean value of the diameter of the colonies in the presence of a given concentration of each compound, and *C* (mm) – the mean diameter of the colonies in the absence of the compound under the same conditions. All the

experiments were carried out in triplicate. The standard deviation was calculated.

The half maximal effective concentration for compound **4e** was determined by linear regression of the probability of the corresponding percentage of fungus radial growth from the logarithm of the concentration [29] in the GraphPad Prism 9.4.1 program. The commercial fungicide carbendazim, which is highly active against *P. lingam*, was used as a comparator agent.


3. Results and Discussion

Compounds **3a-c** were obtained by reacting the corresponding cyanoacetamides **1a-c** with thioglycolic acid **2** in pyridine with the addition of dimethylaminopyridine as a catalyst according to a previously published procedure (Scheme 3 [25]). Further alkylation with methyl iodide or benzyl chloride in dimethylformamide in the presence of K₂CO₃ gave products **4a-e** in 64–80% yields.

The fungicidal activity of obtained compounds **3a-c** and **4a-e** was studied against 8 strains of phytopathogenic fungi that are widespread and cause significant damage to agriculture, such as *A. solani*, *B. cinerea*, *C. coccodes*, *F. solani*, *P. infestans*, *P. lingam*, *R. solani* and *S. sclerotiorum*.

An *in vitro* study of antifungal activity showed that most of the obtained compounds exhibit low (*I* < 50%) or moderate (50% < *I* < 70%) activity against phytopathogenic fungi, inhibiting the growth of mycelium by 65% or less (Table 1).

However, compound **4e**, containing 2-methylphenyl at the nitrogen atom of the acetamide fragment and benzyl at the nitrogen atom of the thiazolidine ring, showed high activity against *A. solani* (causative agent of early blight of nightshade crops) and *P. lingam* (causative agent of cruciferous plants phomosis) with the inhibition degree of the fungi radial growth 75.77 and 85.94%, respectively.



Scheme 3 Synthesis of compounds **3a-c** and **4a-e**.

Table 1 Results of antifungal activity study *in vitro* for compounds **3a-c** and **4a-e** at a concentration of 50 µg/mL*.

| Com- pounds | Degree of inhibition of mycelial growth (I±SD, %) | | | | | | | |
|----------------|---|-------------------|--------------------|------------------|------------------|---------------------|------------------|---------------------|
| | <i>A. solani</i> | <i>B. cinerea</i> | <i>C. coccodes</i> | <i>F. solani</i> | <i>R. solani</i> | <i>P. infestans</i> | <i>P. lingam</i> | <i>S. scleroti-</i> |
| 3a | 29.60±0.53 | 29.95±0.96 | 9.34±0.65 | 7.28±1.17 | 13.38±0.72 | 8.77±0.35 | 28.95±2.26 | 31.58±0.31 |
| 3b | 37.80±0.60 | 32.50±1.21 | 4.20±0.77 | 9.63±0.99 | 9.75±0.68 | 5.48±0.60 | 1.95±0.16 | 6.06±0.94 |
| 3c | 54.95±2.02 | 6.70±0.23 | 39.21±0.22 | 4.86±0.63 | 2.40±0.79 | 6.33±0.50 | 26.12±3.30 | 10.98±3.11 |
| 4a | 37.22±2.63 | 12.25±0.67 | 58.65±0.18 | 1.08±1.12 | 9.76±0.77 | 10.25±0.21 | 43.45±2.48 | 17.01±1.04 |
| 4b | 54.11±0.61 | 10.84±2.19 | 36.18±1.04 | 5.09±0.28 | 9.59±0.18 | 9.90±0.88 | 31.83±0.63 | 25.63±0.34 |
| 4c | 51.73±1.56 | 8.21±0.76 | 47.04±0.52 | 3.42±0.38 | 0 | 12.33±1.12 | 30.59±1.15 | 64.45±2.10 |
| 4d | 47.37±0.47 | 7.92±0.61 | 19.43±0.57 | 6.83±0.66 | 9.84±0.81 | 8.62±0.91 | 11.83±0.34 | 7.58±1.71 |
| 4e | 75.77±0.39 | 22.89±1.00 | 54.47±0.11 | 8.74±0.96 | 3.27±0.32 | 10.23±1.38 | 85.94±0.06 | 34.09±0.91 |

*SD – standard deviation, *I* = 100 – active compound, *I* = 0 – in active compound.

Half maximal effective concentration values of 1,3-thiazolidin-2-ylidene **4e** for *A. solani* and *P. lingam* were determined (Table 2).

Thus, promising results were obtained. Compound **4e** was found to have low EC₅₀ values at the level of 1–2 µg/mL, comparable with commercial fungicides [30].

4. Limitations

In this paper, we present the first data on the antifungal activity of 2-(4-oxo-1,3-thiazolidin-2-ylidene)acetamide derivatives. The proposed approach for the preparation of potential fungicides is a one-step method for the synthesis of 2-(4-oxo-1,3-thiazolidin-2-ylidene)acetamide scaffold. This approach makes it easy to modify the resulting compounds. Using this strategy makes it possible to obtain a wide range of compounds. Thus, this direction in the future will be followed so as to discover new compounds that are highly active against fungi. For targeted design of the potential fungicide structures based on the 2-(4-oxo-1,3-thiazolidin-2-ylidene)acetamides, additional studies of the mode of their biological action and identification of biological targets are required.

Also, it is necessary to perform further biological studies *in vivo* of the highly active compounds to evaluate their selectivity, toxicity to humans, animals and beneficial microorganisms, etc.

5. Conclusions

Thus, as a result of the work, we synthesized 8 derivatives of 2-(4-oxo-1,3-thiazolidin-2-ylidene)acetamide and studied their fungicidal activity *in vitro*. Compound **4e** was highly active against *A. solani* and *P. lingam*. It was shown that 2-(4-oxo-1,3-thiazolidin-2-ylidene)acetamides are promising base compounds for designing new fungicides.

• Supplementary materials

This article contains supplementary materials with copies of ¹H, and ¹³C spectra, which are available on the corresponding online page.

• Funding

This research was supported by the Russian Science Foundation and Government of Sverdlovsk region, Joint Grant No 22-26-20124, <https://rscf.ru/en/project/22-26-20124>.



• Acknowledgments

The authors are grateful to the Laboratory of Integrated Research and Expert Evaluation of Organic Materials of Ural Federal University for the registration of NMR spectra of compounds.

Table 2 Antifungal EC₅₀ values of compound **4e**.

| Fungi | Compounds | Regression equation | R ² | EC ₅₀ , µg/mL |
|------------------|-----------|-------------------------|----------------|--------------------------|
| <i>A. solani</i> | 4e | $y = 1.4767x + 9.5355$ | 0.9654 | 0.85 |
| | CZ* | $y = 0.5110x + 2.3270$ | 0.9794 | >10000 |
| <i>P. lingam</i> | 4e | $y = 2.3053x + 11.0900$ | 0.9881 | 2.29 |
| | CZ | $y = 3.649x + 15.7200$ | 0.9880 | 1.15 |

*CZ – commercial fungicide carbendazim.

• Author contributions

Conceptualization: K.L.O., T.V.G.

Data curation: K.L.O., T.A.K.

Formal Analysis: T.A.K.

Funding acquisition: T.A.K.

Investigation: T.A.K., D.V.R., M.F.K.

Methodology: T.V.G., K.L.O.

Project administration: T.A.K.

Resources: T.A.K., T.V.G.

Supervision: T.V.G.

Validation: T.A.K., T.V.G.

Visualization: K.L.O.

Writing – original draft: K.L.O., T.A.K.

Writing – review & editing: T.V.G.

• Conflict of interest

The authors declare no conflict of interest.

• Additional information

Author IDs:

Konstantin L. Obydenov, Scopus ID [54684683000](https://orcid.org/0000-0002-5468-4683);

Tatiana A. Kalinina, Scopus ID [54684032600](https://orcid.org/0000-0002-5468-4032600);

Maria F. Kosterina, Scopus ID [6506644209](https://orcid.org/0000-0002-6506-644209);

Tatiana V. Glukhareva, Scopus ID [6602582999](https://orcid.org/0000-0002-6602-582999).

Website:

Ural Federal University, <https://urfu.ru/en>.

References

- Trotsko N. Antitubercular properties of thiazolidin-4-one. Eur J Med Chem. 2021;215:113266. doi:[10.1016/j.ejmech.2021.113266](https://doi.org/10.1016/j.ejmech.2021.113266)
- Kumar H, Deep A, Kumar Marwaha R. Design, synthesis, in silico studies and biological evaluation of 5-((E)-4-((substituted aryl/alkyl)methyl)benzylidene)thiazolidine-2,4-dione derivatives. BMC Chem. 2020;14:25. doi:[10.1186/s13065-020-00678-2](https://doi.org/10.1186/s13065-020-00678-2)
- Chandrappa S, Kavitha CV, Shahabuddin MS, Vinaya K, Ananda Kumar CS, Ranganatha SR, Raghavan SC, Rangappa KS. Synthesis of 2-(5-((5-(4-chlorophenyl)furan-2-yl)methylidene)-4-oxo-2-thioxothiazolidin-3-yl)acetic acid derivatives and evaluation of their cytotoxicity and induction of apoptosis in human leukemia cells. Bioorg Med Chem. 2009;17(6):2576–2584. doi:[10.1016/j.bmc.2009.01.016](https://doi.org/10.1016/j.bmc.2009.01.016)
- Zhou H, Wu S, Zhai S, Liu A, Sun Y, Li R, Zhang Y, Ekins S, Swaan PW, Fang B, Zhang B, Yan B. Design, synthesis, cytotoxicity, structure-activity relationships, and pharmacophore of thiazolidinone derivatives targeting drug-resistant

- lung cancer cells. *J Med Chem.* 2008;51(5):1242–1251. doi:[10.1021/jm7012024](https://doi.org/10.1021/jm7012024)
5. Uchôa FDT, da Silva TG, de Lima M do CA, Galdino SL, Pitta I da R, Costa TD. Preclinical pharmacokinetic and pharmacodynamic evaluation of thiazolidinone PG15: An anti-inflammatory candidate. *J Pharm Pharmacol.* 2009;61(3):339–345. doi:[10.1211/jpp.61.03.0008](https://doi.org/10.1211/jpp.61.03.0008)
 6. Agarwal A, Lata S, Saxena KK, Srivastava VK, Kumar A. Synthesis and anticonvulsant activity of some potential thiazolidinonyl 2-oxo/thiobarbituric acids. *Eur J Med Chem.* 2006;41(10): 1223–1229. doi:[10.1016/j.ejmech.2006.03.029](https://doi.org/10.1016/j.ejmech.2006.03.029)
 7. Terzioğlu N, Karalı N, Gürsoy A, Pannecouque C, Leysen P, Paeshuysse J, Neyts J, De Clercq E. Synthesis and primary antiviral activity evaluation of 3-hydrazono-5-nitro-2-indolinone derivatives. *Arxivoc.* 2006;1:109–118.
 8. Powers JP, Piper DE, Li Y, Mayorga V, Anzola J, Chen JM, Jaen JC, Lee G, Liu J, Peterson MG, Tonn GR, Ye Q, Walker PC, Wang, Z. SAR and Mode of Action of Novel Non-Nucleoside Inhibitors of Hepatitis C NS5b RNA Polymerase. *J Med Chem.* 2006;49:1034–1046. doi:[10.1021/jm050859x](https://doi.org/10.1021/jm050859x)
 9. Pizzo C, Saiz C, Talevi A, Gavernet L, Palestro P, Bellera C, Cazzulo JJ, Chidichimo A, Wipf P, Mahler SG. Synthesis of 2-Hydrazolyl-4-Thiazolidinones Based on Multicomponent Reactions and Biological Evaluation Against *Trypanosoma Cruzi*. *Chem Biol Drug Des.* 2011;77(3):166–172. doi:[10.1111/j.1747-0285.2010.01071.x](https://doi.org/10.1111/j.1747-0285.2010.01071.x)
 10. Bhandari SV, Bothara KG, Patil AA, Chitre TS, Sarkate AP, Gore ST, Dangre SC, Khachane CV. Design, Synthesis and Pharmacological Screening of Novel Antihypertensive Agents Using Hybrid Approach. *Bioorg Med Chem.* 2009;17(1),390–400. doi:[10.1016/j.bmc.2008.10.032](https://doi.org/10.1016/j.bmc.2008.10.032)
 11. Vicini P, Geronikaki A, Incerti M, Zani F, Dearden J, Hewitt M. 2-Heteroarylrimino-5-benzylidene-4-thiazolidinones analogues of 2-thiazolylimino-5-benzylidene-4-thiazolidinones with antimicrobial activity: Synthesis and structure-activity relationship. *Bioorg Med Chem.* 2008;16(7):3714–3724. doi:[10.1016/j.bmc.2008.02.001](https://doi.org/10.1016/j.bmc.2008.02.001)
 12. Salem MA. Synthesis of New Thiazole, Bithiazolidinone and Pyrano[2,3-d]thiazole Derivatives as Potential Antimicrobial Agents. *Croat Chem Acta.* 2017;90(1):7–15. doi:[10.5562/cca2955](https://doi.org/10.5562/cca2955)
 13. Siddiqui IR, Singh PK, Singh J, Singh J. Facile synthesis and fungicidal activity of novel 4,4'-bis[2''-(5'''-substituted rhodanin-3'''-yl)thiazol-4''-yl]bibenzyls. *Indian J. Chem., Sect B [Internet].* 2005[cited 2016];44: 2102–6. English. Available from: <http://nopr.niscpr.res.in/handle/123456789/9197>, Accessed on 07 December 2022.
 14. Mendgen T, Steuer C, Klein CD. Privileged scaffolds or promiscuous binders: A comparative study on rhodanines and related heterocycles in medicinal chemistry. *J Med Chem.* 2011;55(2):743–753. doi:[10.1021/jm201243p](https://doi.org/10.1021/jm201243p)
 15. Huang L, Lümmen P, Berry EA. Crystallographic investigation of the ubiquinone binding site of respiratory Complex II and its inhibitors. *Biochim Biophys Acta – Proteins and Proteom.* 2021;1869:140679. doi:[10.1016/j.bbapap](https://doi.org/10.1016/j.bbapap)
 16. Noha RM, El hamid MK, Ismail MM, Manal RM, Salwa E. Design, synthesis and screening of benzimidazole containing compounds with methoxylated aryl radicals as cytotoxic molecules on (HCT-116) colon cancer cells. *Eur J Med Chem.* 2091:112870. doi:[10.1016/j.ejmech.2020.112870](https://doi.org/10.1016/j.ejmech.2020.112870)
 17. Wang S, Zhao Y, Zhu W, Liu Y, Guo K, Gong P. Synthesis and Anticancer Activity of Indolin-2-one Derivatives Bearing the 4-Thiazolidinone Moiety. *Arch Pharm Chem Life Sci.* 2012;345:73–80. doi:[10.1002/ardp.201100082](https://doi.org/10.1002/ardp.201100082)
 18. Ranjan P, Kumar SP, Kari V, Jha PC. Exploration of interaction zones of b-tubulin colchicine binding domain of helminths and binding mechanism of anthelmintics. *Comput Biol Chem.* 2017;68:78–91. doi:[10.1016/j.compbiolchem.2017.02.008](https://doi.org/10.1016/j.compbiolchem.2017.02.008)
 19. Avupati VR, Yejella RP, Akula A, Guntuku GS, Doddi BR, Vutla VR, Anagani SR, Adimulam LS, Vyricharla AK. Synthesis, characterization and biological evaluation of some novel 2,4-thiazolidinediones as potential cytotoxic, antimicrobial and antihyperglycemic agents. *Bioorg Med Chem Lett.* 2012;22(20):6442–6450. doi:[10.1016/j.bmcl.2012.08.052](https://doi.org/10.1016/j.bmcl.2012.08.052)
 20. Li YX, Wang SH, Li ZM, Su N, Zhao WG. Synthesis of novel 2-phenylsulfonylhydrazono-3-(2',3',4',6'-tetra-O-acetyl-β-d-glucopyranosyl)thiazolidine-4-ones from thiosemicarbazide precursors. *Carbohydr Res.* 2006;341(17):2867–2870. doi:[10.1016/j.carres.2006.09.010](https://doi.org/10.1016/j.carres.2006.09.010)
 21. Haroun M, Tratrart C, Kolokotroni A, Petrou A, Geronikaki A, Ivanovic M, Kostic M, Sokovic M, Carazo A, Mlad ěnka P, Sreeharsha N, Venugopala KN, Nair AB, Elsewedy HS. 5-Benzylidene-2-(5-methylthiazol-2-ylimino)thiazolidin-4-ones as Antimicrobial Agents. Design, Synthesis, Biological Evaluation and Molecular Docking Studies. *Antibiot.* 2021;10(3):309. doi:[10.3390/antibiotics10030309](https://doi.org/10.3390/antibiotics10030309)
 22. Abo-Bakr AM, Hassan EA, Mahdy A-HS, Zayed SE. Synthetic and biological studies on some new camphor thiazolidinones. *J Iran Chem Soc.* 2021;18(10):2757–2769. doi:[10.1007/s13738-021-02228-6](https://doi.org/10.1007/s13738-021-02228-6)
 23. Orchard MG, Neuss JC, Galley CM, Carr A, Porter DW, Smith P, Scopes DIC, Haydon D, Voudsen K, Stubberfield CR, Young K, Page M. Rhodanine-3-acetic acid derivatives as inhibitors of fungal protein mannosyl transferase 1 (PMT1). *Bioorg Med Chem Lett.* 2004;14(15):3975–3978. doi:[10.1016/j.bmcl.2004.05.050](https://doi.org/10.1016/j.bmcl.2004.05.050)
 24. Chaban T, Matiichuk Y, Chulovska Z, Tymoshuk O, Chaban I, Matiychuk V. Synthesis and biological evaluation of new 4-oxo-thiazolidin-2-ylidene derivatives as antimicrobial agents. *Arch Pharm.* 2021;354(7):2100037. doi:[10.1002/ardp.202100037](https://doi.org/10.1002/ardp.202100037)
 25. Obydenov KL, Galushchinskiy AN, Kosterina MF, Glukhareva TV, Morzherin YY. Synthesis of 4-(4-oxo-1,3-thiazolidin-2-ylidene)-pyrrolidine-2,3,5-triones. *Chem Heterocycl Compd.* 2017;53(5):622–625. doi:[10.1007/s10593-017-2102-0](https://doi.org/10.1007/s10593-017-2102-0)
 26. Kolarević A, Ilić BS, Kocić G, Džambaski Z, Šmelcerović A, Bondžić BP. Synthesis and DNase I inhibitory properties of some 4-thiazolidinone derivatives. *J Cell Biochem.* 2019;120(1):264–274. doi:[10.1002/jcb.27339](https://doi.org/10.1002/jcb.27339)
 27. Džambaski Z, Marković R, Kleinpeter E, Baranac-Stojanović M. 2-Alkylidene-4-oxothiazolidine S-oxides: synthesis and stereochemistry. *Tetrahedron.* 2013;69(31):6436–6447. doi:[10.1016/j.tet.2013.05.087](https://doi.org/10.1016/j.tet.2013.05.087)
 28. Obydenov KL, Kalinina TA, Galieva NA, Beryozkina TV, Zhang Y, Fan Z, Glukhareva TV, Bakulev VA. Synthesis, fungicidal activity, and molecular docking of 2-acylamino and 2-thioacylamino derivatives of 1H-benzo[d]imidazoles as anti-tubulin agents. *J Agric Food Chem.* 2021;69(40):12048–12062. doi:[10.1021/acs.jafc.1c03325](https://doi.org/10.1021/acs.jafc.1c03325)
 29. Obydenov KL, Khamidullina LA, Galushchinskiy AN, Shatunova SA, Kosterina MF, Kalinina TA, Fan Zh, Glukhareva TV, Morzherin YuYu. Discovery of Methyl (5Z)-[2-(2,4,5-Trioxopyrrolidin-3-ylidene)-4-oxo-1,3-thiazolidin-5-ylidene]acetates as Antifungal Agents against Potato Diseases. *J Agric Food Chem.* 2018;66(24):6239–6245. doi:[10.1021/acs.jafc.8b02151](https://doi.org/10.1021/acs.jafc.8b02151)
 30. Yu X, Teng P, Zhang Y-L, Xu Z-J, Zhang M-Z, Zhang W-H. Design, synthesis and antifungal activity evaluation of coumarin-3-carboxamide derivatives. *Fitoterapia.* 2018;127:387–395. doi:[10.1016/j.fitote.2018.03.013](https://doi.org/10.1016/j.fitote.2018.03.013)