

PR-12

METHOD DEVELOPMENT AND VALIDATION OF A REVERSED PHASE HPLC METHOD FOR DETERMINATION OF ANASTRAZOLE AND TEMOZOLOMIDE IN PHARMACEUTICAL DOSAGE FORM**N. MD. Akram,^{1*} N. Madana Gopal,² A. Balakrishna,³ N. Bakthavatchala Reddy⁴**¹*Santhiram Engineering College, Nandyal, Kurnool (Dt), Andhra Pradesh, India.*²*Santhiram College of pharmacy, Nandyal, Kurnool(Dt), Andhra Pradesh, India.*³*Rajeev Gandhi Memorial College of Engineering and Technology (Autonomous), Nandyal 518501, Andhra Pradesh, India.*⁴*Ural Federal University, Chemical Engineering Institute, Yekaterinburg, 620002, Russian Federation.*

*Corresponding author, E-mail: mdakram.chem@gmail.com

Abstract. A new simple assay method has been developed and validated for the determination of Anastrozole and Temozolomide using reverse-phase high performance liquid chromatography in their pharmaceutical dosage form. The chromatographic separation was performed on an Inertsil ODS (4.6 x 150mm, 5 μ m) using mobile phase phosphate buffer pH 3.0 and methanol of 30:70% v/v at a flow rate of 0.8 mL/min. Analytes were detected at 260 nm. The method was found to be linear in the concentration range of 1-5 μ g/mL for both medicaments with the coefficient value (R²) of >0.999. The accuracy was measured via recovery studies and found to be acceptable and the percentage recoveries were found in the range of 98.81-100.72 and 99.29-100.70%. The proposed method was successfully validated and applied for the quantitative estimation of these drugs in both bulk and tablet dosage forms.