

The Variational Calculation of Bulk Moduli for Liquid Binary Alloys of Alkali Metals

Acta Physica Polonica A 131, 237 (2019), ERRATUM

N.E. DUBININ^{a,b,*}

^aUral Federal University, 19 Mira Str., 620002 Ekaterinburg, Russia

^bInstitute of Metallurgy of the Ural Branch of the Russian Academy of Sciences,
101 Amundsen Str., 620016 Ekaterinburg, Russia

(Received April 11, 2016; in final form November 27, 2016)

The bulk moduli of the Na–K, Na–Cs, and K–Rb liquid equiatomic alloys at $T = 373$ K are calculated by the variational method with the hard-sphere reference system. The local Animalu–Heine model pseudopotential and Toigo–Woodruff exchange-correlation function are used for the calculation. A good agreement with experimental data is achieved.

original DOI: [10.12693/APhysPolA.131.237](https://doi.org/10.12693/APhysPolA.131.237)

actual DOI: [10.12693/APhysPolA.136.225](https://doi.org/10.12693/APhysPolA.136.225)

PACS/topics: 05.70.Ce, 61.20.Ne, 71.22.+i, 62.10.+s

This article was originally published on February 2017 with some mistakes in Eqs. (4) and (7). In corrected form these equations must be rewritten as follows:

$$\phi_{\text{NFE}}(r) = \frac{z^2}{r} + \frac{\Omega}{\pi^2} \int_0^{\infty} F(q) \frac{\sin(qr)}{qr} q^2 dq, \quad (4)$$

$$\phi_{\text{NFE}_{ij}}(r) = \frac{z_i z_j}{r} + \frac{\Omega^{\text{bin}}}{\pi^2} \int_0^{\infty} F_{ij}(q) \frac{\sin(qr)}{qr} q^2 dq. \quad (7)$$

Author apologizes for this error.

*e-mail: ned67@mail.ru